

ACCELERON[®]

D-109

Fungicide Seed Treatment

For commercial and on-farm use

Seed treatment for disease control and plant health in cotton and soybean.

Active Ingredient*:

pyraclostrobin: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-

yl]oxy]methyl]phenyl]methoxy-, methyl ester) 18.4%

Other Ingredients: 81.6%

Total: 100.0%

*Equivalent to 1.67 pounds of pyraclostrobin per gallon formulated as a flowable concentrate for seed treatment.

EPA Reg. No. 7969-266-264

EPA Est. No. 51036-GA-001

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See full label for **complete First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577.

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937).

Net Contents: 1.25 GAL

Product of Germany

Article No: 81142952

MON99999984A 220926A 03/23

**PRODUCED FOR: Bayer CropScience LP • 800 N. Lindbergh Blvd.
St. Louis, Missouri 63167**

FIRST AID

If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• DO NOT give anything to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.• Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Avoid contact with skin or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber* \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils.
- Shoes plus socks

*Includes natural rubber blends and laminates

In addition to the PPE listed in other portions of the Personal Protective Equipment (PPE) section, handlers involved in multiple commercial seed treatment activities must wear:

- Dust/mist respirator with MSHA/NIOSH approval number prefix TC-21 or any N, R, P, or HE filter

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of pyraclostrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Groundwater Advisory

Pyraclostrobin is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in possession of the user at the time of fungicide application.

DO NOT apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions, and **Conditions of Sale and Warranty** are to be followed.

AGRICULTURAL USE REQUIREMENTS

Use in this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and the restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

EXCEPTION: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Socks and shoes
- Chemical-resistant gloves made of any waterproof material

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use. **DO NOT** store near food or feed.

Pesticide Disposal: To avoid pesticide waste, use all material in this container by application according to label directions. If pesticide waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity < 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

In Case of Emergency

In case of a large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BAYER CROPSCIENCE 1-800-334-7577

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

Steps to take if this material is released into the environment or spilled:

- Wear **Personal Protective Equipment (PPE)** and avoid exposure when managing a spill. (See **Precautionary Statements** section of this label for required PPE.)
- Dike and contain the spill with inert, absorbent material (e.g., sand, earth) and transfer liquid and solid diking material to separate containers for disposal. Small-scale spills of this product (that can be cleaned up with a typical spill kit) may be applied to labeled sites.
- Remove contaminated clothing, and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep spill out of all sewers and open bodies of water.

Product Information

Acceleron® D-109 Fungicide Seed Treatment is a broad spectrum liquid fungicide that protects seed and seedlings from certain seed-borne and soil-borne diseases resulting in improved plant health. **Acceleron® D-109 Fungicide Seed Treatment** increases emergence and plant stands by controlling or suppressing these diseases. See **Acceleron® D-109 Fungicide Seed Treatment Crop-specific Use Directions** for crop uses and diseases controlled or suppressed by **Acceleron® D-109 Fungicide Seed Treatment**. Disease symptoms such as seed rot, seedling rot, seedling blight, root rot, and damping-off (preemergence or postemergence) can be associated with the actions of various fungal pathogens. These symptoms are referred to in **Acceleron® D-109 Fungicide Seed Treatment Crop-specific Use Directions** as "seed and seedling disease".

Seed treatment applications with **Acceleron® D-109 Fungicide Seed Treatment** have been associated with more rapid and increased emergence of seedlings under certain cold conditions.

Use the higher rates of **Acceleron® D-109 Fungicide Seed Treatment** when disease pressure is expected to be high.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapco.org/metals.htm>.

Mode of Action

Pyraclostrobin, the active ingredient of **Acceleron® D-109 Fungicide Seed Treatment**, belongs to the group of respiration inhibitors classified by the US EPA and Canada PMRA as quinone outside inhibitors (QoI), or target-site-of-action **Group 11** fungicides.

Resistance Management

Acceleron® D-109 Fungicide Seed Treatment contains a **Group 11** (pyraclostrobin) fungicide. Any fungal population may contain individuals naturally resistant to **Acceleron® D-109 Fungicide Seed Treatment** and other **Group 11** fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, including enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Acceleron® D-109 Fungicide Seed Treatment** or other **Group 11** fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated pest management (IPM) program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development. Monitor the efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/ or disease development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance, contact your BAYER CROPS SCIENCE representative, local extension specialist, or certified crop advisor.

Application Instructions

Acceleron® D-109 Fungicide Seed Treatment can be used in commercial seed treatment facilities and on-farm systems. For use with commercial or professional seed treatment application equipment only. **DO NOT** apply with planting application equipment, e.g., hopper box, planter box, seed treaters, etc.

Seed must be of a good quality, well conditioned, and free of chaff before treatment.

The purchaser of **Acceleron® D-109 Fungicide Seed Treatment** is responsible for ensuring all seed treated with this product is adequately dyed with a suitable color to prevent accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25.

- Use an EPA-approved dye or colorant that imparts an unnatural color to the seed.
- Any dye or colorant added to treated seed must be cleared for use under 40 CFR, Part 180.900, "Exemptions from the requirement of a tolerance". Federal regulations have established official tolerances or exemptions from tolerances for residues on food and forage crops that must not be exceeded when the product (dye or colorant) is used at specified rates.

Prepare no more mixture of **Acceleron® D-109 Fungicide Seed Treatment** than is needed for immediate application to seed. Refer to the **Tank Mixing Information** section of this label. Apply **Acceleron® D-109 Fungicide Seed Treatment** at the rates listed in **Acceleron® D-109 Fungicide Seed Treatment Crop-specific Use Directions** as a water-based mixture using standard slurry or mist-type seed treatment application equipment. The exact amount of water needed to provide the mixture or slurry rate for uniform and complete coverage of the seed surface is difficult to predict because weather conditions, seed type and surface, and equipment used all have a bearing on coverage. Thorough seed coverage and proper application equipment calibration are essential for good disease control. Consult a seed treatment specialist regarding slurry rates for the crop seed to be treated with **Acceleron® D-109 Fungicide Seed Treatment** and for calibration and operation procedures of the seed treatment equipment.

Tank Mixing Information

Tank Mix Partners

When using **Acceleron® D-109 Fungicide Seed Treatment** in a tank mixture with other seed treatment products, follow all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations on the tank mix partner label. **DO NOT** exceed label dosage or application rate. The most restrictive label precautions, limitations and restrictions must be followed. **Acceleron® D-109 Fungicide Seed Treatment** must not be mixed with any product that prohibits such mixing.

Compatibility Test for Tank Mix Components

When tank mixing **Acceleron® D-109 Fungicide Seed Treatment**, always test compatibility before use by performing a jar test as follows:

1. Add all intended seed treatment products and the appropriate amount of water in a clear glass jar.
2. Mix well and allow to sit for one hour.
3. Remix and observe for physical incompatibility.

Also test with the seed to ensure the combination of tank mix products does not have detrimental effects on the seed.

Mixing Order

For containers 5 gallons or less, shake well before use. For containers greater than 5 gallons, recirculate before use.

1. **Water** – Add 3/4 of the required volume of water to the mix tank and turn on agitation.
2. **Agitation** – Mechanical agitation is required for proper mixing of **Acceleron® D-109 Fungicide Seed Treatment**.
3. **Dye or colorant**
4. **Wettable powders**
5. **Soluble powders**
6. **Flowable formulations** (such as **Acceleron® D-109 Fungicide Seed Treatment**)
7. **Emulsifiable concentrates**
8. **Water** – Remaining 1/4 of required water volume is added last and can be used as rinsate for the chemical container or to adjust the total slurry to compensate for water displacement by powdered products.

Maintain agitation until the entire slurry mixture has been used.

Ensure seed is dry before bagging.

Use Restrictions

- **DO NOT** use treated seed for food, feed, or oil processing.
- **DO NOT** store **Acceleron® D-109 Fungicide Seed Treatment** near feed or foodstuffs.
- **DO NOT** contaminate feed or foodstuffs.
- **DO NOT** contaminate domestic or irrigation waters.
- **DO NOT** plant crops within 30 days after planting **Acceleron® D-109 Fungicide Seed Treatment**-treated seed unless permitted according to the seed labeling note.
- **DO NOT** use the same auger for treating seed that will handle commercial or feed grains.
- **DO NOT** store leftover treated seed where there is a chance of it becoming mixed with untreated seed. Store treated seed in cool, dry conditions.

SEED LABELING NOTE

Seed treated with this product that is then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag.

"This seed has been treated with Acceleron® D-109 Fungicide Seed Treatment containing pyraclostrobin. DO NOT use treated seed for food, feed or oil purposes. Store treated seed away from food and feedstuffs. DO NOT allow children, pets or livestock to have access to treated seed. Wear long pants, long-sleeved shirt, shoes and socks, and chemical resistant gloves made of any waterproof material when opening this bag or handling (e.g., loading, pouring) treated seed or seed pieces.

Treated seed exposed on soil surfaces may be hazardous to wildlife. Plant treated seed into the soil to the recommended minimum depth or greater to minimize exposure. Cover or collect treated seed spilled during loading and planting, in particular at row ends and field corners.

Dispose of all excess treated seed by burying seed away from bodies of water. DO NOT enter or allow worker entry into areas with treated seeds during the restricted entry interval of 12 hours, with the exception that workers may enter if they will have no contact with the soil/media subsurface.

Environmental Hazards: This product is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters or rinsate.

Surface Water Advisory: This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of pyraclostrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Groundwater Advisory: Pyraclostrobin is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Dispose of seed packaging or containers in accordance with local requirements. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

In the event of crop failure or harvest of a crop grown from this seed, the field may be replanted immediately with alfalfa, barley, brassica vegetables, bulb vegetables, citrus, corn, cotton, cucurbits, dried shelled peas and beans, edible podded legume vegetables, flax, fruiting vegetables, grass grown for seed, leafy vegetables, leaves of root and tuber vegetables, mint, oat, peanut, pecan, rapeseed (includes canola and crambe), root vegetables, rye, sorghum, soybean, succulent shelled peas and beans, sugar beet, sugar cane, sunflower, tuberous and corn vegetables (including potato), wheat and triticale, or any other crop for which a tolerance for pyraclostrobin exists. For all other crops, the minimum plant back interval is 30 days from the date this seed was planted. A cover crop without a tolerance for pyraclostrobin may be planted sooner than the 30 day interval for purposes of erosion control or soil improvement; however, the crop may not be grazed or harvested for food or feed."

Acceleron® D-109 Fungicide Seed Treatment Crop-specific Use Directions

Crop	Target Disease	Product Use Rate ml/100 kg seed (grams ai/100 kg seed)	Product Use Rate ft ozs/100 lbs seed (lbs ai/100 lbs seed)
Cotton ¹	Control of: <ul style="list-style-type: none"> Seed and seedling disease caused by: <ul style="list-style-type: none"> - <i>Rhizoctonia solani</i> Suppression of: <ul style="list-style-type: none"> Seed and seedling disease caused by: <ul style="list-style-type: none"> - <i>Fusarium</i> spp. - <i>Pythium</i> spp. 	98 to 196 (19.6 to 39.1)	1.5 to 3.0 (0.02 to 0.04)
Soybean ² (<i>Glycine</i> spp.)	Control of: <ul style="list-style-type: none"> Seed and seedling disease caused by: <ul style="list-style-type: none"> - <i>Rhizoctonia solani</i> Suppression of: <ul style="list-style-type: none"> Seed and seedling disease caused by: <ul style="list-style-type: none"> - <i>Fusarium</i> spp. - <i>Phomopsis</i> spp. - <i>Phytophthora</i> spp. - <i>Pythium</i> spp. 	26 to 98 (5.2 to 19.6)	0.4 to 1.5 (0.005 to 0.02)

¹Cotton - 4,500 Seeds per pound

²Soybean - 2,800 Seeds per pound

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

Acceleron® is a registered trademark of Bayer Group.
2020-04-306-0206

©2023 Bayer Group

ACCELERON[®] D-109

Fungicide Seed Treatment

For commercial and on-farm use
Seed treatment for disease control and plant health
in cotton and soybean.

PYRACLOSTROBIN GROUP 11 FUNGICIDE

Active Ingredient*:

pyraclostrobin: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]methoxy-,methyl ester).....18.4%

Other Ingredients:81.6%

Total:100.0%

*Equivalent to 1.67 pounds of pyraclostrobin per gallon formulated as a flowable concentrate for seed treatment.

EPA Reg. No. 7969-266-264

EPA Est. No. 51036-GA-001

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) See full label for complete First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty, and state-specific crop and/or use site restrictions. For MEDICAL AND TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577. For PRODUCT USE Information Call 1-866-998BAYER (1-866-992-2937). **FIRST AID:** If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. **DO NOT** give anything to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes. Call a poison control center or doctor for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. For MEDICAL AND TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577. **Precautionary Statements: Hazards to Humans and Domestic Animals: CAUTION.** Harmful if swallowed. Avoid contact with skin or clothing. **Environmental Hazards:** This product is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading. **STORAGE AND DISPOSAL:** DO NOT contaminate water, food, or feed by storage or disposal. **Pesticide Storage:** Store in original container only. Keep container closed when not in use. **DO NOT** store near food or feed. **Pesticide Disposal:** To avoid pesticide waste, use all material in this container by application, according to label directions. If pesticide waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry). **Container Handling:** Nonrefillable Container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities. **Triple rinse containers small enough to shake (capacity < 5 gallons) as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Acceleron[®] is a registered trademark of Bayer Group. ©2023 Bayer Group

Article No.: 81142952
2020-04-306-0206

Product of Germany
MON99999984A 220926A 03/23

Net Contents: 1.25 GAL

**PRODUCED FOR: Bayer CropScience LP • 800 N. Lindbergh Blvd.
St. Louis, Missouri 63167**