

FLUOPYRAM	GROUP	7	FUNGICIDE
BACILLUS FIRMUS STRAIN I-1582	GROUP	UNB	NEMATOCIDE

ACCELERON[®]

N-314

Seed Treatment

Seed treatment with 2 modes of action for use on corn and cotton for the protection against damage caused by early season plant pathogenic nematodes and for the promotion of plant growth.

ACTIVE INGREDIENTS:

fluopyram: N-[2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl]-2-(trifluoromethyl)benzamide: 29.5%

Bacillus firmus strain I-1582: 11.5%

OTHER INGREDIENTS: 59.0%

TOTAL: 100.0%

Contains 500 g ai/liter or 4.17 lbs ai/gallon

- 3 lbs of fluopyram per gallon (360 g fluopyram per liter)

- 1.17 lbs of Bacillus firmus strain I-1582 per gallon (140 g ai/liter) and a minimum of 4.0×10^9 cfu/ml of product

EPA Reg. No. 7969-460-264

EPA Est. 51036-GA-001

FLOWABLE SUSPENSION

KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete First Aid, Precautionary Statements, Directions for Use, Conditions of Sale and Warranty,
 and state-specific crop and/ or use site restrictions.

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours a Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

Net Contents: 30 Gallons

81167257 NVA 2023-04-0615-0045

MON99999988B 230627B 07/23

**PRODUCED FOR: Bayer CropScience LP • 800 N. Lindbergh Blvd.
 St. Louis, MO 63167 • 1-866-992-2937**

FIRST AID

If swallowed:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER

In case of emergency, call the toll-free Bayer CropScience
Emergency Response telephone number: 1-800-334-7577.
Have a product container or label with you when calling a poison
control center or doctor, or going for treatment.

Note to Physician: Treat symptomatically.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Workers involved with treating the seed (e.g. connecting and disconnecting hoses and transfer pumps, mixing, equipment calibration, etc.); baggers and bag sewers; cleaners/ repairers of seed treatment equipment and others exposed to the concentrate must wear:

- Long-sleeve shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made out of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride ≥ 14 mils or Viton ≥ 14 mils
- Wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air-purifying respirator with an HE filter . (Repeated exposures to high concentrations of microbial proteins can cause allergic sensitization.)

ENGINEERING CONTROLS

When engineering controls that meet the OSHA requirements defined in 29 CFR Part 1910.1000 are used the respirator requirement may be waived or modified as specified in 29 CFR Part 1910.1000.

USER SAFETY REQUIREMENTS

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features, such as ponds, streams, and springs will reduce the potential loading of fluopyram. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seed spilled during loading.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label and observe all label directions and precautions before use.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage, disposal or cleaning of equipment.

Pesticide Storage

Store in original container away from feed and food. Store in cool, dry area. DO NOT store in direct sunlight. DO NOT allow prolonged storage in temperatures that exceed 86° F (30° C) or in temperatures that fall below 32° F (0° C).

Pesticide Disposal

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling

Dilutable Seed Treatment Products in Nonrefillable Plastic Containers
Rigid nonrefillable containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)

Nonrefillable Container. DO NOT reuse or refill this container. After emptying product from container, either return container to Bayer CropScience, or rinse and either recycle or dispose of the container as follows:

Bottom discharge Intermediate Bulk Container (IBC) (e.g. Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container.

To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top discharge IBC, drums, kegs (e.g. Snyder 120 Next Gen, Bonar B120, drums, and kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IN CASE OF SPILL

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- Bayer CropScience 1-800-334-7577

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- Bayer CropScience 1-800-334-7577

Steps to take if material is released or spilled:

- Wear Personal Protective Equipment (PPE) and avoid exposure when managing a spill. (See **Precautionary Statements** section of this label for required PPE.)
- Dike and contain the spill with inert, absorbent material (e.g., sand, earth) and transfer liquid and solid diking material to separate containers for disposal. Small-scale spills of this product (that can be cleaned up with a typical spill kit) may be applied to labeled sites.
- Remove contaminated clothing, and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep spill out of all sewers and open bodies of water.

PRODUCT INFORMATION

ACCELERON® N-314 SEED TREATMENT is a systemic and biological seed treatment that when applied to the seed, aids in protecting the seed and seedling against damage caused by early season plant pathogenic nematodes and promotes plant growth.

In areas of high nematode infestation additional control measures may be warranted.

USE RESTRICTIONS

- For use only in commercial seed treatment equipment.
- Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.
- **DO NOT** use this product except as directed on this label or on other Bayer CropScience supplemental labeling for this product.
- **DO NOT** broadcast plant seeds.
- **DO NOT** apply more than 0.45 lb ai/A of flupyradimorph per year (including seed treatment, foliar, in-furrow, or banded applications).
- Refer to the specific use directions and restrictions in the **Detailed Use Directions** section.
- This product does not contain colorant. The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, Part 153.155.

FUNGICIDE RESISTANCE MANAGEMENT (FRAC)

For resistance management, ACCELERON N-314 SEED TREATMENT contains a Group 7 fungicide. Any fungal population may contain individuals naturally resistant to ACCELERON N-314 SEED TREATMENT and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of ACCELERON N-314 SEED TREATMENT or other Group 7 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your Bayer CropScience representative, pesticide distributor, or university extension specialist.

APPLICATION INSTRUCTIONS

Use ACCELERON N-314 SEED TREATMENT in commercial seed treatment facilities with commercial or professional seed treatment application equipment only.

- Use good quality, well-conditioned seed free of chaff.
- Thorough and uniform seed coverage is required for optimum performance.
- Apply at specified rates with either seed treatment equipment that has "direct inject" or dedicated pump capabilities or in a waterbased mixture using standard slurry seed treatment application equipment. Consult a seed treatment specialist about crop-specific slurry rates and seed treatment calibration and operation procedures. If applying in a water-based mixture mix only as much product as is needed for immediate application to seed.

TANK MIXING OTHER PRODUCTS

ACCELERON N-314 SEED TREATMENT can be used in combination with other registered Bayer CropScience seed treatment fungicides and insecticides. Pretest for compatibility with other seed treatment products.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Compatibility Test

Before mixing components, always perform a compatibility jar test.

1. In a clean quart jar containing 1/2 the intended amount of water for a final slurry application rate, add the labeled rate of ACCELERON N-314 SEED TREATMENT and follow with all intended tank mix components.
2. Add remaining balance of water. Total slurry volume depends on seed size and complete, uniform coverage of the seed, as well as the commercial seed treating application equipment used.
3. Mix well and allow to sit for one hour.
4. Evaluate - **DO NOT** use mixtures that curdle, precipitate, or gel.

MIXING INSTRUCTIONS

When applied with seed treatment application equipment designed to treat seed utilizing a treatment slurry mixture:

- Add approximately 1/2 to 2/3 of the required water to the treatment slurry tank followed by the accompanying seed treatment products and ACCELERON N-314 SEED TREATMENT.

- The order of product addition is dependent on the accompanying seed treatment products and application equipment used.
- Allow each slurry component to disperse completely prior to the next addition until a uniform suspension is obtained.
- After all slurry components have been added and dispersed completely, add the remaining 1/2 to 1/3 of required water for proper slurry volume and mix until uniform.
- **DO NOT** store mixed slurries for greater than 72 hours.
- Maintain adequate agitation until treatment slurry is applied.
- For additional instructions for use, slurry component mixing order, and other application suggestions contact your Bayer CropScience representative.

DETAILED USE DIRECTIONS

ALL CROPS

- For use only in commercial seed treatment equipment.
- Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.
- Apply with seed treatment equipment that has "direct inject" or dedicated pump capabilities or standard slurry-type.
- All seed treated with this product must be colored with an EPA-approved dye or colorant that imparts an unnatural color to the seed to help prevent the inadvertent use of treated seed for food, feed, or oil purposes.
- Treated seed must be labeled in accordance with the requirements of the Federal Seed Act.
- **DO NOT** broadcast plant seeds.
- **DO NOT** apply more than 0.45 lb ai/A of fluopyram per year (including seed treatment, foliar, in-furrow, or banded applications).

CORN**ACCELERON N-314 SEED TREATMENT RATE**

Soilborne Nematodes	# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	ml/80,000 seeds ¹ (fl ozs/80,000 seeds)	mg ai/seed ¹	g/A ² (lb/A)
Ring (<i>Criconebella</i> spp.)					
Root lesion (<i>Pratylenchus</i> spp.)				0.035 to 0.35 [delivers 0.025 to 0.25 mg fluopyram/ seed]	1.41 to 14.09 (0.0031 to 0.031) [delivers 0.0022 to 0.022 lb fluopyram/A]
Southern root-knot (<i>Meloidogyne incognita</i>)	3,706 (1,681)	25.94 to 259.4 (0.4 to 3.98)	5.6 to 56.0 (0.19 to 1.89)		
Sting (<i>Belonolaimus</i> spp.)					
Stubby-root (<i>Paratrichodorus</i> spp.)					

¹ Rate is calculated based on number of seeds/kg average of corn (3,706).

² Based on seeding rate of 40,250 seeds/acre [(40,250 seeds/A) x (mg ai/seed) x (g/1,000 mg)]

COTTON**ACCELERON N-314 SEED TREATMENT RATE**

Soilborne Nematodes	# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	ml/100,000 seeds ¹ (fl ozs/100,000 seeds)	mg ai/seed ¹	g/A ² (lb/A)
Lance (<i>Hoplolaimus</i> spp.)					
Reniform (<i>Rotylenchulus reniformis</i>)	9,921 (4,500)	476.19 to 972.23 (7.3 to 14.91)	48 to 98 (1.62 to 3.31)	0.24 to 0.49 [delivers 0.175 to 0.35 mg fluopyram/ seed]	20.4 to 41.6 (0.0449 to 0.0917) [delivers 0.033 to 0.066 lb fluopyram/A]
Southern root-knot (<i>Meloidogyne incognita</i>)					

¹ Rate is calculated based on number of seeds/kg average of cotton (9,921).

² Based on seeding rate of 85,000 seeds/acre [(85,000 seeds/A) x (mg ai/seed) x (g/1,000 mg)]

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with **ACCELERON N-314 SEED TREATMENT** must be labeled with the following statements:

- This seed has been treated with **ACCELERON N-314 SEED TREATMENT**, which contains **29.5% of fluopyram** and **11.5% Bacillus firmus strain I-1582**.
- **DO NOT** use treated seed for food, feed, or oil purposes.

User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

- Store treated seeds away from food and feedstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- Wear long-sleeved shirt, long pants and chemical- resistant gloves made out of: barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride \geq 14 mils or Viton \geq 14 mils when handling treated seed.
- Excess treated seed may be used for ethanol production only if (1) byproducts are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading or planting.
- Dispose of all excess treated seed. Leftover treated seed may be buried away from water sources in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment washwaters.
- Treated seed must be adequately covered with soil at planting.
- Dispose of seed packaging or containers in accordance with local requirements. **DO NOT** use empty seed bags for any other purpose.
- To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's specifications. Surplus seed or empty seed containers must be stored or disposed according to local federal regulations.
- Treat at manufacturer's specified rate.

Restrictions:

- **DO NOT** broadcast plant seeds.
- **DO NOT** apply more than 0.45 lb ai/A of fluopyram per year (including seed treatment, foliar, in-furrow, or banded applications).

Rotational Crops Plant-back Restrictions:

- The following crops can be replanted immediately following the last application of fluopyram: artichoke (globe); Brassica (cole) leafy vegetables (group 5); bulb vegetables (subgroup 3-07A and 3-07B); carrot; cereals (group 15) [which include: barley, buckwheat, field corn, popcorn, sweet corn, millet (pearl and proso), oats, rice; rye, sorghum, teosinte, triticale, and wheat]; citrus (group 10-10); cotton (subgroup 20C); cucurbits (group 9); dill seed; fruiting vegetables (group 8-10); ginseng; grapes and small vines (except fuzzy kiwifruit) (subgroup 13-07F); herb (subgroup 19A); hop; leafy vegetables (except watercress) (group 4); legume vegetables (except soybean, cowpea and dry peas); peanut; pome fruit (group 11-10); potato and other root, tuberous and corm vegetables (except sugar beet) (subgroup 1B and 1C); rapeseed (subgroup 20A); small berries (canberries and bushberries) (subgroup 13-07A and 13-07B); soybean; stone fruits (group 12-12); strawberry and other low-growing berries, except cranberry (subgroup 13-07G); sugar beet; sunflower (subgroup 20B); tobacco; tree nut (group 14-12).
- Alfalfa can be planted after 14 days and sugarcane can be planted after 14 days in Region 3 - Florida and boot of Alabama.
- **DO NOT** rotate to crops other than those listed above.

CONDITIONS OF SALE AND WARRANTY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

FLUOPYRAM	GROUP	7	FUNGICIDE
BACILLUS FIRMUS STRAIN I-1582	GROUP	UNB	NEMATOCIDE

ACCELERON[®]

N-314

Seed Treatment

Seed treatment with 2 modes of action for use on corn and cotton for the protection against damage caused by early season plant pathogenic nematodes and for the promotion of plant growth.

ACTIVE INGREDIENTS:

fluopyram: N-[2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl]-2-(trifluoromethyl)benzamide: 29.5%

Bacillus firmus strain I-1582: 11.5%

OTHER INGREDIENTS: 59.0%

TOTAL: 100.0%

Contains 500 g ai/liter or 4.17 lbs ai/gallon

- 3 lbs of fluopyram per gallon (360 g fluopyram per liter)

- 1.17 lbs of Bacillus firmus strain I-1582 per gallon (140 g ai/liter) and a minimum of 4.0 x 10⁹ cfu/ml of product

EPA Reg. No. 7969-460-264

EPA Est. 51036-GA-001

FLOWABLE SUSPENSION

KEEP OUT OF REACH OF CHILDREN

CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete First Aid, Precautionary Statements, Directions for Use, Conditions of Sale and Warranty, and state-specific crop and/ or use site restrictions.

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours a Day 1-800-334-7577 For PRODUCT USE information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID: If swallowed: Immediately call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give anything by mouth to an unconscious person. If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. **HOTLINE NUMBER:** In case of emergency, call the toll-free Bayer CropScience Emergency Response telephone number: 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment. Note to Physician: Treat symptomatically. **PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION.** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. **ENVIRONMENTAL HAZARDS:** For terrestrial uses: DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwater or rinsate. This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fluopyram. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff. This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seed spilled during loading. **STORAGE AND DISPOSAL: DO NOT** contaminate water, food or feed by storage, disposal or cleaning of equipment. **Pesticide Storage:** Store in original container away from feed and food. Store in cool, dry area. DO NOT store in direct sunlight. DO NOT allow prolonged storage in temperatures that exceed 86° F (30° C) or in temperatures that fall below 32° F (0° C). **Pesticide Disposal:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry). **Container Handling:** Dilutable Seed Treatment Products in Nonrefillable Plastic Containers. Rigid nonrefillable containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs). Nonrefillable Container. DO NOT reuse or refill this container. After emptying product from container, either return container to Bayer CropScience, or rinse and either recycle or dispose of the container as follows: **Bottom discharge Intermediate Bulk Container (IBC) (e.g. Schuetz Caged IBC or Snyder Square Stackable).** Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40, PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve. **Top discharge IBC, drums, kegs (e.g. Snyder 120 Next Gen, Bonar B120, drums, and kegs).** Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times. Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Acceleron[®] is a registered trademark of Bayer Group. ©2023 Bayer Group. All rights reserved.

Produced For: Bayer CropScience LP - 800 N. Lindbergh Blvd. - St. Louis, MO 63167

Net Contents: 30 Gallons

81167257 NVA 2023-04-0615-0045

MON99999988B 230627B 07/23