ACTION®

Guaranteed Analysis

0-0-0

Calcium 5%

Derived from Calcium Chloride

CAUTION Keep out of Reach of Children May Cause Eye Irritation

USER SAFETY RECOMMENDATIONS

Avoid contact with skin, eyes or clothing. Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove wet clothing immediately. Wash skin thoroughly and put on clean clothing.

FIRST AID

If in eyes: Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. If on skin: Take off contaminated clothing and rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. Have product container or label with you when calling a poison control center, or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

STORAGE AND DISPOSAL


Store this product out of direct sunlight, in a cool place above 40°F. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Dispose of empty container in a sanitary landfill or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

NET WEIGHT: 9.2 lbs/gal or 1.10 kg/L

NET CONTENTS:

☐ 2.5 gallons/23.0 lbs ☐ 10 liter/11.0 Kg ☐ 5.0 gallons/46.0 lbs ☐ 20 liter/22.0 Kg

COMPATIBILITY: ACTION is tank-mix compatible with most pesticides and fertilizers. Do not mix **ACTION** with products containing phosphorus or sulfur or other chemicals known to react with calcium. Add **ACTION** to the diluted solution last. Always conduct a compatibility test for new or untried tank-mix combinations.



GENERAL INFORMATION

ACTION contains immediately available forms of the essential plant nutrient calcium. Apply **ACTION** to the seed, soil or crop foliage for uptake by the plant. **ACTION** promotes maximum gene expression, cell division and plant growth maximizing root production and plant vigor.

DIRECTIONS FOR USE

It is a violation of federal law to use this product inconsistent with its labeling. **ACTION** can be applied as a seed treatment, in-furrow, in the transplant water, as a side-dress or foliar application, and through flood, drip, overhead, pivot and micro irrigation systems. Apply the recommended amount in a volume sufficient to provide uniform coverage.

RECOMMENDED USES: Apply as a seed treatment at 4 to 8 ounces per 100 pounds of seed for large-seeded crops. Do not use treated seed for food, feed or oil purposes. Label treated seed in accordance with the requirements of the Federal Seed Act and applicable state and local laws. Use an approved dye to distinguish treated seed and prevent inadvertent use for food, feed or oil purposes. For small-seeded crops, apply 4 to 8 ounces in furrow per planted acre.

Apply to the soil or foliage starting with the transplant water or when seeded crops have 2 to 4 true leaves and continue through harvest every 7 days using 4 to 8 ounces per planted acre or apply every 14 days using 16 ounces per planted acre. Use **ACTION** at the recommended rate on the following crops:

Field Crops: alfalfa, bean crops, canola, clover, corn, cotton, forage grasses, peanut, potato, rape (canola), rice, sorghum, soybean, sugar beet, sugarcane, sunflower, tobacco, and wheat/barley/oat.

Vegetable Crops: artichoke, asparagus, beet, carrot, cruciferous, fennel, herbs, leafy vegetables, melon, onion/garlic, cucurbits, eggplant, pepper, strawberry, sweet potato and tomato.

Fruit and Nut Crops: almond, apple, apricot, avocado, berry crops, cherry, citrus, date palm, fig, filbert, grapes, hazelnut, hops, nectarines, oil palm, olive, persimmon, peach, pear, pecan, pistachio, plums/prune, pomegranate, subtropical fruit and walnut.

Transplants: apply ACTION as a root dip or drench at 4 to 16 ounces per 100 gallons of water immediately before or after planting.