

This information is for promotional purposes only. Space considerations may require information to be omitted.
Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

ASULAM GROUP 18 HERBICIDE

ADDAX™

36.2 SL



**Apply Addax™ 36.2 SL for postemergence control of listed weeds in
Christmas Tree Plantings Sugarcane, Turf, Ornamentals
and in Non-Cropland Areas**

Not intended for use in residential areas. Only for commercial or agricultural use.

ACTIVE INGREDIENT:

Sodium salt of asulam (methyl sulfanylcarbamate)* 36.2%

OTHER INGREDIENTS:..... 63.8%

TOTAL:..... 100.0%

*Equivalent to 33.1% asulam or not less than 3.34 lbs. sodium asulam per gallon.

EPA Reg. No. 91234-157

KEEP OUT OF REACH OF CHILDREN CAUTION

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)**

See below for additional Precautionary Statements.

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none">▪ Take off contaminated clothing.▪ Rinse skin immediately with plenty of water for 15-20 minutes.▪ Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes.▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.▪ Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-884-685-9173 for emergency medical treatment information.	

**For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, chemical-resistant gloves (such as Nitrile, Butyl, Neoprene, and/or Barrier Laminate), and shoes plus socks. Follow manufacturer's instructions for cleaning /maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cab that meets the definition listed in the Worker Protection Standard for agricultural pesticides [40 CFR 170.305].

USER SAFETY RECOMMENDATIONS

Users should leave the treated area, remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination. Surface water contamination may occur in areas with poorly draining soils and little or no buffers or in areas where drainage systems flow directly to surface water.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not clean equipment or dispose of equipment washwater in a manner that will contaminate resources. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

RESTRICTIONS

Pilots must use an enclosed cab that meets the definition listed in the Worker Protection Standard for agricultural pesticides [40 CFR 170.305].

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is coveralls, chemical resistant gloves, and shoes plus socks.

WEED RESISTANCE MANAGEMENT

For resistance management, Addax 36.2 SL is a Group 18 herbicide. Any weed population may contain or develop plants naturally resistant to Addax 36.2 SL and other Group 18 herbicides. Weed species with acquired resistance to Group 18 herbicides may eventually dominate the weed population if Group 18 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Addax 36.2 SL or other Group 18 herbicides. Users should scout before and after application.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To delay herbicide resistance:

- Avoid the consecutive use of Addax 36.2 SL or other target site of action Group 18 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern (an herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides)
- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.



- Scout fields after application to verify that the treatment was effective.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

Report any incidence of non-performance of this product against a particular weed species to your Atticus, LLC retailer or representative. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemicals means to remove escapes, as practical, with the goal of preventing further seed production.

SPRAY DRIFT

Aerial Applications:

- Do not release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

Ground Boom Applications:

- Apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground crop canopy.
- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

Adjust nozzles – Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and move laterally in a concentrated cloud (under low wind conditions) indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.



PRODUCT INFORMATION

Do not apply Addax 36.2 SL through irrigation systems of any type.

CHRISTMAS TREE PLANTINGS

Apply Addax 36.2 SL as a postemergence application in the following species of Christmas Tree Plantings:

- Douglas Fir
- Grand Fir
- Noble Fir
- Scotch Pine

Apply this product as a spray mixture combined with water. Use a minimum of 20 gallons of solution per acre via ground application.

Weed	Rate of Addax 36.2 SL	Directions
Western Bracken (<i>Pteridium aquilinum var. pubescens</i>)	1 gallon per acre (3.34 lbs. active ingredient)	Treat with Addax 36.2 SL application solution following bud break and after firming/hardening of new tree growth. Prior to application, target species must be in full frond.

Restrictions:

- Do not graze or feed foliage from treated areas to livestock.
- Do not use a wetting agent with Addax 36.2 SL.
- Do not apply this product via aerial application.
- Apply Addax 36.2 SL a maximum of once per year.
- Do not apply more than 1 gallon of product per acre per year (3.34 lbs. active ingredient per acre per year).

SUGARCANE

Apply this product as a spray mixture combined with water to sugarcane grown from stubble or plant cane. Only apply Addax 36.2 SL when weeds are growing actively.

The application of fertilizer and/or some cultivation/cultural practices that disturb the target species root system may lead to sub optimal control of target species. Do not employ practices that will disturb the root system of target species for a minimum of 7 days before application of this product or for a minimum of 7 days after treatment with this product.

IMPORTANT: In Louisiana there have been reports of varying tolerances among crops to applications of Addax 36.2 SL. For further information, refer to the local County Agent or University Extension Specialist.

In order to enhance weed control in sub-optimal environmental conditions, use one of the following adjuvants:

Adjuvant*	Directions
Crop Oil Concentrate	Must contain 15-20% non-ionic surfactant and 80-85% paraffin based petroleum oil. Apply at the rate of 4 qts./100 gals. (1% V/V) of application mixture.
Non-Ionic Surfactant	Must contain 80% active ingredient minimum. Apply at the rate of 1-2 qts./100 gals. (0.25/0.5% V/V) of application mixture.

*Adjuvants must be cleared for application in growing crops.

Ground Application

Depending on local application practice, apply Asulam 36.5 SL in 15-100 gallons of water per acre.

Aerial Application: Except in Hawaii, apply Addax 36.2 SL in 3-5 gallons of water per acre. In Hawaii, apply Addax 36.2 SL in 5-10 gallons of water per acre.

In order to calculate the correct rate for banded application, use the following formula:

(Band width(in inches) ÷ Row Width (in inches))	x	Broadcast rate (see tables below)	=	Band Rate per acre
-------------------------------------------------	---	-----------------------------------	---	--------------------

Rates for broadcast application of Asulam 3.6F for a maximum of one application per year:

Weed	Rate of Addax 36.2 SL	Directions
Alexandergrass (<i>Brachiaria plantaginea</i>)	6-8 Pints per acre (2.50-3.34 lbs. active ingredient)	Use the higher rate where the grass is greater than 8 inches in height. Use the lower rate where grass is 6-8 inches in height or less.
Barnyardgrass (<i>Echinochloa crusgalli</i>)	6-8 Pints per acre (2.50-3.34 lbs. active ingredient)	Use the higher rate where the grass is greater than 8 inches in height. Use the lower rate where grass is 6-8 inches in height or less.
Broadleaf Panicum (<i>Panicum adspersum</i>)	6-8 Pints per acre (2.50-3.34 lbs. active ingredient)	Use the higher rate where the grass is greater than 8 inches in height. Use the lower rate where grass is 6-8 inches in height or less.
Crabgrass (<i>Digitaria spp.</i>)	6-8 Pints per acre (2.50-3.34 lbs. active ingredient)	Use the higher rate where the grass has reached the early seed head formation stage. Use the lower rate where grass is yet to reach the seed head formation stage.
Foxtail (<i>Setaria spp.</i>)	6-8 Pints per acre (2.50-3.34 lbs. active ingredient)	Use the higher rate where the grass is greater than 8 inches in height. Use the lower rate where grass is 6-8 inches in height or less.
Goosegrass (<i>Eleusine indica</i>)	6-8 Pints per acre (2.50-3.34 lbs. active ingredient)	Use the higher rate where the grass is greater than 8 inches in height. Use the lower rate where grass is 6-8 inches in height or less.
Itchgrass or Raoulgrass (<i>Rottboellia exaltata</i>)	8 Pints per acre (3.34 lbs. active ingredient)	Application mixture must be combined with a surfactant. Apply this product when grass is a maximum of 8 inches in height.
Johnsongrass (<i>Sorghum halepense</i>)	8 Pints per acre (3.34 lbs. active ingredient)	Apply Addax 36.2 SL when grass is growing actively and is 12-18 inches in height. At the time of application air temperature must be a minimum of 60°F.
Paragrass or Californiagrass (<i>Brachiaria mutica</i> or <i>Panicum purpurascens</i>)	8 Pints per acre (3.34 lbs. active ingredient)	Apply this product when grass is a maximum of 6-8 inches in height.

Two applications of Addax 36.2 SL in one season may be necessary where there are initially heavy infestations of target species, where rhizome Johnsongrass is present or when treating target species that germinate at different times during the growing season.



Rates for broadcast application of Asulam 3.6F for two applications per season:

Weed	Application 1: Rate of Addax 36.2 SL	Application 2: Rate of Addax 36.2 SL	Directions
Crabgrass (<i>Digitaria spp.</i>)	6 to 8 pints per acre (2.50-3.34 lbs. active ingredient)	6 to 8 pints per acre (2.50-3.34 lbs. active ingredient)	For the 1 st and 2 nd application, apply to grass before seed head formation.
Itchgrass or Raoulgrass (<i>Rottboellia exaltata</i>)	8 pints per acre (3.34 lbs. active ingredient)	8 pints per acre (3.34 lbs. active ingredient)	Application mixture must be combined with a surfactant. For the 1 st and 2 nd application grass must be a maximum of 8 inches in height.
Johnsongrass (<i>Sorghum halepense</i>)	8 pints per acre (3.34 lbs. active ingredient)	8 pints per acre (3.34 lbs. active ingredient)	For the 1 st and 2 nd application grass must be 12-18 inches in height.

Spot Treatment

Apply Addax 36.2 SL as a 5% v/v application spray (i.e. 1 gallon of Asulam 3.62 SL per 20 gallons of water).

RESTRICTIONS

- When applying Asulam 3.6 SL as a spot treatment, do not apply more than 8 pints of product per acre per year (3.34 lbs. active ingredient per acre per year).
- Apply Addax 36.2 SL a maximum of twice per year.
- Cover crops may be planted if they are ploughed under and not used for grazing.
- Mainland USA (except Louisiana): do not harvest crops for a minimum of 140 days following application of this product.
- Louisiana: do not harvest crops for a minimum of 100 days following application of this product.
- Hawaii: do not harvest crops for a minimum of 400 days following application of this product.
- Do not graze or feed sugarcane fodder to livestock.

NON-CROPLAND

Apply Addax 36.2 SL for control of listed species as a postemergence application in non-cropland sites, for example, railroad rights-of-way and yards, utility rights-of-way and yards, highway and roadside rights-of-way, pipeline rights-of-ways, storage areas and industrial plant sites, warehouse lots, lumberyards, boundary fences and fence rows.

The application solution of Addax 36.2 SL may be combined with an approved non-ionic surfactant at a rate of 0.25% by volume.

Prepare this product as a single water mix treatment at a rate of 20-100 gallons of application solution (dependent on local practice) per acre. Apply to the target area as a ground application to control species listed in the table below.

Weed	Rate of Addax 36.2 SL	Directions
Crabgrass (<i>Digitaria spp.</i>)	1 gallon per acre (3.34 lbs. active ingredient)	Treat with Addax 36.2 SL when grass is yet to reach the seed head formation stage.
Johnsongrass (<i>Sorghum halepense</i>)	1 gallon per acre (3.34 lbs. active ingredient)	Treat with Addax 36.2 SL when grass is a minimum of 18 inches in height. When applying as a spot treatment in Hawaii, apply this product in 100 gallons of solution. Do not exceed 50 gallons total of application solution per acre.
Paragrass or Californiagrass (<i>Brachiaria mutica</i> or <i>Panicum purpurascens</i>)	1 gallon per acre (3.34 lbs. active ingredient)	Treat with Addax 36.2 SL when grass is yet to reach the seed head formation stage. When applying as a spot treatment in Hawaii, apply this product in 100 gallons of solution. Do not exceed 50 gallons total of application solution per acre.
Western Bracken (<i>Pteridium aquilinum var. pubescens</i>)	7-8 pints per acre (2.92-3.34 lbs. active ingredient)	Treat with Addax 36.2 SL when ferns are in full frond.

Restrictions:

- Do not apply this product via aerial application
- Apply Addax 36.2 SL a maximum of once per Year.
- Do not apply more than 1 gallon of product per acre per year (3.34 lbs. active ingredient per acre per year).

TURF (Sod Farms Only)

Apply Addax 36.2 SL in Tifway 419 Bermudagrass and St. Augustinegrass and turf as a postemergence treatment in order to control listed weeds (see table below).

Apply Asulam 36.2L as a spray solution in 20-50 gallons of water per acre.

Weed	Rate of Addax 36.2 SL	Directions
Bullgrass (<i>Paspalum supinum</i>), Crabgrass (<i>Digitaria sp.</i>), Goosegrass (<i>Eleusine indica</i>)	5 pints per acre (2.09 lbs. active ingredient)	Apply Addax 36.2 SL in St. Augustinegrass to control listed weed species.
Sandbur (<i>Cenchrus sp.</i>)	5 pints per acre (2.09 lbs. active ingredient)	Apply Addax 36.2 SL in Tifway 419 Bermudagrass to control listed weed species.

Restrictions:

- Apply Addax 36.2 SL a maximum of once per year.
- Do not combine a surfactant with the application solution of Addax 36.2 SL.
- Do not treat freshly mown grass or to turf that is under stress.
- Do not apply more than 5 pints of product per acre per year (2.08 lbs. active ingredient per acre per year).



ORNAMENTALS

Apply this product as a single application. Apply as a postemergence broadcast treatment in a minimum of 20 gallons of water per acre. Use Addax 36.2 SL to treat specified target species in the following ornamental species:

Junipers: Juniperus Andorra, Juniperus chinensis, Juniperus conferta, Juniperus horizontalis, Juniperus litoralis, Juniperus Sabina

Yews: Podocarpus macrophyllus, Taxus cuspidate, Taxus media

Restrictions:

- Do not combine a surfactant with the application solution of Addax 36.2 SL.
- Apply Addax 36.2 SL a maximum of once per year.
- Do not apply more than 1 gallon of product per acre per year (3.34 lbs. active ingredient per acre per year).

Weed	Rate of Addax 36.2 SL	Directions
Barnyardgrass (<i>Echinochloa crusgali</i>) Crabgrass (<i>Digitaria sp.</i>) Fall Panicum (<i>Panicum dichotomiflorum</i>) Foxtails (<i>Setaria sp.</i>) Goosegrass (<i>Eleusine indica</i>) Horseweed (marestail) <i>Conyza Canadensis</i>	1 gallon per acre (3.34 lbs. active ingredient)	Apply to listed species between the early seedling stage and the early seed head formation stage.

The use of Addax 36.2 SL may be affected by local conditions. Refer to Experiment Station and/or State Agricultural weed specialists for information on possible lower dosages and for guidance on local weed problems.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store at temperatures above 32°F. Do not allow product to freeze.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable containers (five gallons or less). Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a rinse tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable containers (greater than five gallons). Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse or pressure rinse as follows. **Triple rinse:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Pressure rinse:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Return to point of sale or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.



**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Atticus, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Atticus, LLC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Atticus, LLC or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Atticus, LLC's election, one of the following: (1) Refund of purchase price paid by buyer or user for product bought, or (2) Replacement of amount of product used.

To the extent consistent with applicable law, Atticus, LLC shall not be liable for losses or damages resulting from handling or use of this product unless Atticus, LLC is promptly notified of such loss or damage in writing. In no case, to the extent consistent with applicable law, shall Atticus, LLC be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Atticus, LLC or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

F20200131a

SPECIMEN