

# Adrenalyn™ ETQ

Turf Fungicide

Tebuconazole	GROUP 3	FUNGICIDE
Azoxystrobin	GROUP 11	FUNGICIDE



ACTIVE INGREDIENTS:	% BY WT
Azoxystrobin: Methyl (E)-2-[[6-(2-cyanophenoxy-4-pyrimidinyl)]alpha-methoxymethylene] benzeneacetate	18.2%
Tebuconazole: (+)-alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl-1H-1,2,4-triazole-1-ethanol	27.3%
<b>OTHER INGREDIENTS:</b>	54.5%
<b>TOTAL:</b>	100.0%

Adrenalyn ETQ Turf Fungicide is a suspension concentrate fungicide containing 2.55 lbs. Tebuconazole and 1.69 lbs. Azoxystrobin per gallon.

## KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Emergency Phone Numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 222-1222 Poison Control Center
-------------------------	---

See additional Precautionary Statements and Directions for Use inside the booklet.

NET CONTENTS: 1 Gallon

Manufactured for:  
**SIPCAM AGRO USA, INC.**  
2525 Meridian Parkway  
Durham, NC 27713

# 1G

EPA Reg. No. 60063-62

EPA Est. No.: 60063-GA-1 (Lot number begins with VL)

EPA Est. No.: 70815-GA-1 (Lot number begins with CB)

EPA Est. No.: 86555-MO-1 (Lot number begins with AF)

EPA 20200730 (08/20)



**SIPCAM AGRO**  
USA, INC.

READ THE  
ENTIRE LABEL  
BEFORE USING  
THIS PRODUCT

FUNGICIDE

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING.** May be fatal if swallowed. Harmful if absorbed through skin or if inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Human flagging is prohibited.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

#### All handlers must wear:

- Long-sleeved shirt and long pants
- Water-proof gloves
- Shoes plus socks
- In addition, mixers/loaders/applicators using mechanically pressurized handwands, except when applying to nursery ornamentals and landscaping, must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Respirator fit testing, medical qualification, and training

Using a program that conforms to OSHA's requirements (see 29 CFR Part), employers must verify that any handler who uses a respirator is:

- Fit-tested and fit-checked,
- Trained, and
- Examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn. A qualified medical practitioner is a physician or other licensed health care professional who will evaluate the ability of a worker to wear a respirator. The initial evaluation consists of a questionnaire that asks about medical conditions (such as a heart condition) that would be problematic for respirator use. If concerns are identified, then additional evaluations, such as a physical exam, might be necessary. The initial evaluation must be done before respirator use begins. Handlers must be reexamined by a qualified medical practitioner if their health status or respirator style or use-conditions change.

Upon request by local / state / federal / tribal enforcement personnel, employers must provide documentation demonstrating how they have complied with these requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirement may be reduced or modified as specified in the WPS.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish, and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean highwater mark. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

### Ground Water Advisory

Azoxystrobin can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Tebuconazole is known to leach through soil into ground water under certain conditions as a result of label use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

Notify state and/or Federal authorities and Sipcam Agro USA, Inc. immediately if you observe any adverse environmental effects due to use of this product.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Human flagging is prohibited.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR PLANT INJURY.**

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area treated must be vacated by unprotected persons.

**DO NOT** enter or allow others to enter until sprays have dried.

**DO NOT** treat areas while unprotected humans or domestic animals are present in the treatment area. Because certain States may require more restrictive entry intervals, consult your State Department of Agriculture for further information.

### PRODUCT INFORMATION

This product has both systemic and curative properties and is effective against a wide range of organisms, providing preventative control of many significant plant diseases. Apply as a foliar spray or in tank mix combinations with other turf and ornamental products.

### RESTRICTIONS

- OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES:
  - Apply only during alternate years in sites adjacent to aquatic areas listed above.
  - **DO NOT** apply by ground or air within 100 feet of aquatic areas listed above.
  - **DO NOT** cultivate within 10 feet of an aquatic area to allow for growth of a vegetative filter strip.
- **DO NOT** spray this product where spray drift may reach apple trees. **DO NOT** use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

## **PRECAUTIONS**

- This product is extremely phytotoxic to certain apple varieties. AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees and apple fruit.

### **MANDATORY SPRAY DRIFT MANAGEMENT**

#### **Groundboom Applications:**

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

### **SPRAY DRIFT ADVISORIES**

**THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.**

## **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

## **BOOM HEIGHT**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

## **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

## **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

## **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## **RESISTANCE MANAGEMENT**

For resistance management, please note that this product contains both a Group 3 (tebuconazole) and a Group 11 (azoxystrobin) fungicide. Any fungal population may contain individuals naturally resistant to this product and other Group 3 or Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3 or Group 11 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your local Sipcarn Agro representative. You can also contact your pesticide distributor or university extension specialist to report resistance.

Use of a multiple-site contact fungicide (such as chlorothalonil containing products) as an alternating or tank-mix partner can help alleviate selection pressure on pathogens and delay the onset of resistance. **DO NOT** tank mix this product with any fungicide to which resistance has already been developed.

## DISEASE CONTROL IN GOLF COURSE TURF

### Use Restrictions

- For use only on golf course turf.
- **DO NOT** apply more than 6 applications per year. For use on gray snow mold and pink snow mold, **DO NOT** apply more than 3 applications per year.
- **DO NOT** exceed 4.5 fl. oz. product per 1,000 sq. ft. (or 200 fl. oz. product per acre) (3.98 lbs. a.i. tebuconazole per A) per year.
- **DO NOT** use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, play fields, etc.
- Not for residential use.
- Not for use on turf being grown for sale or commercial use as sod.
- **DO NOT** use clippings for animal feed.
- Apply by ground (includes any hand-held and mechanically pressurized handwands) only.
- **DO NOT** apply by air or through any type of Ultra Low Volume (ULV) spray system (<5 gallons per A).
- In the State of New York, **DO NOT** exceed 2.25 fl. oz. of this product per 1,000 sq. ft. (1.96 lbs. a.i. tebuconazole per A) per year.

### Spray Preparation

Begin application with a clean water spray tank (remove scale, pesticide residues, and other foreign matter, and flush the system with clean water).

### Mixing Instructions:

1. Fill the tank with ½ or ¾ the desired amount of water.
2. Start mechanical or hydraulic agitation.
3. Slowly add the specified amount of this product.
4. Add the remaining volume of water.

Maintain continuous agitation to keep the material in suspension. For best results, use spray mixture the same day it is prepared.

### Compatibility

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

To determine the physical compatibility of this product with other products, the following procedure should be followed: Pour the specified proportions of the products into a suitable container of water, mix thoroughly and allow to stand for at least five (5) minutes. If the combination remains uniformly mixed or can be re-mixed readily, the mixture is considered physically compatible.

Biological compatibility, i.e. effectiveness for disease control or lack of phytotoxicity, must be determined on a case-by-case basis. **DO NOT** mix this product with any other pesticide, fertilizer, adjuvant or other materials except water unless you know that such mixtures will be effective and non-injurious under your conditions of use. When an adjuvant is to be used, use a Council of Producers and Distributors of Agrotechnology (CPDA) certified adjuvant.

This product has demonstrated some phytotoxic effects when mixed with products that are formulated as EC's. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

### Prevention and Control of Golf Course Diseases

Use this product on both cool season and warm season turf grasses including, but not limited to, Bentgrasses, Bermudagrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, Zoysia grasses and Paspalum grasses or their mixtures.

Use this product for the prevention and control of the diseases listed in the table below. Begin applications when conditions are favorable for disease development and repeat applications as long as these conditions persist. Apply preventative treatments at 14- to 21-day intervals as indicated. When treating golf greens, always treat aprons and approaches. Use properly calibrated equipment to spray uniformly over the area to be treated. Use the highest rate and shortest interval listed in the table below for severe disease pressure.

### Turf Application Instructions

Apply the specified amount of this product in sufficient water for thorough coverage. Use a volume of 65 to 135 gallons per acre (1.5 to 3.0 gallons per 1,000 sq. ft.). Apply using properly calibrated low volume, hand-held, mechanical or motorized ground broadcast equipment. Maintain constant agitation during application. Make all applications after mowing and allow foliage to dry thoroughly before irrigation.

**Note:** Prior to pouring, slowly invert container several times to assure uniform mixture.

### GOLF COURSE TURF APPLICATION RATES

APPLICATION METHOD	fl.oz. Product / 1,000 ft <sup>2</sup>	Max. Number of Applications	Max. Single Application (fl.oz./1,000 ft <sup>2</sup> )	Max. Amount / Year*	
				fl.oz. Product / 1,000 ft <sup>2</sup>	fl.oz. Product / acre
Ground (All diseases except Snow Mold)	0.5 -1.0	6**	1.0	4.59**	199.78**
Ground (Snow Mold)	1.45	3	1.45	4.35	189.49
Mechanically Pressurized Hand-Held	0.275	6	0.275	1.65	71.87

\*In the State of New York, **DO NOT** exceed 2.25 fl. oz. of this product per 1,000 sq. ft. (1.96 lbs. a.i. tebuconazole per A) per year.

\*\* **DO NOT** apply more than 3.98 lbs.Ai / acre of tebuconazole per year (maximum of 4.59 fl.oz. product / 1,000 ft<sup>2</sup> or 199.78 fl.oz. product / acre)

### Equivalence Chart for Golf Course Turf Application Rates

fl.oz. Product / 1,000 ft <sup>2</sup>	lbs. Ai / 1,000 ft <sup>2</sup>		fl.oz. Product / acre	lbs. Ai / acre	
	Azoxystrobin	Tebuconazole		Azoxystrobin	Tebuconazole
0.275	0.46	0.70	11.98	0.16	0.24
0.5	0.85	1.28	21.78	0.29	0.43
1.0	1.70	2.55	43.56	0.58	0.87
1.45	2.45	3.70	63.16	0.83	1.26

## GOLF COURSE TURF APPLICATIONS AND INSTRUCTIONS

See **GOLF COURSE TURF APPLICATION RATE** table above for appropriate rates for the application method to be used.

TARGET DISEASES	APPLICATION DIRECTIONS
Anthracnose (Basal** and Foliar) <i>(Colletotrichum cereale)</i>	Use preventatively. Begin applications when disease conditions exist and prior to disease symptom development. Repeat application after 14 to 21 days. A second application may be made after 14 days for greens. ** Use the higher listed rate and shorter intervals for control of Basal Anthracnose.
Bentgrass Dead Spot <i>(Ophiosphaerella agrostis)</i>	Begin applications prior to disease development. Repeat application after 14 days.
Fairy Ring <i>(Lycoperdon spp., Agrocybe pediades, and Bovistia plumbea)</i>	Apply as disease symptoms develop. Make applications in 2 to 4 gallons of water per 1,000 sq. ft. Symptoms may take 2 to 3 weeks to disappear following application. Repeat applications may be required after 28 days in certain cases.
Brown Patch and Zoysia Patch, Large Patch, Leaf and Sheath Spot <i>(Rhizoctonia solani, R. spp., R. zeae)</i>	Brown Patch: Apply when disease conditions exist. Repeat application after 14 to 21 days. Zoysia Patch: Make 1 or 2 applications approximately one month prior to zoysia grass dormancy. Repeat application after 14 to 21 days. Large Patch: For control of Large Patch on all warm season turfgrasses, make 1 or 2 applications in the fall prior to infection or when conditions are favorable for infection. Repeat applications after 14 to 21 days. Leaf and Sheath Sport: Apply when disease conditions exist such as sequential periods of temperatures at or above 90°F. Curative control may require several applications. Target spray at the crown of turfgrass. Repeat application after 14 to 21 days.
Cool Weather Brown Patch, Yellow Patch <i>(Rhizoctonia cerealis)</i>	Make 1 or 2 applications in the fall when disease conditions exist. Repeat applications every 14 to 21 days.
Copper Spot <i>(Sclerotinia homoeocarpa)</i>	Apply preventatively when disease conditions exist. Repeat application after 14 to 21 days.
Dollar Spot <i>(Sclerotinia homoeocarpa)</i>	Apply preventatively when disease conditions exist. Repeat application after 14 to 21 days.
Gray Leaf Spot <i>(Pyricularia grisea)</i>	Apply in a preventative disease control program. Apply before disease is present and alternate with other fungicides that control gray leaf spot. Repeat application after 14 to 21 days.
Gray Snow Mold	Apply once in late fall prior to snow cover. For snow mold, do not make more than three applications at 1.45 fl.oz. / 1000 sq.ft..
Pink Snow Mold <i>(Microdochium nivale)</i> Fusarium Patch (prior to snow cover) (Intermittent snow cover and rainfall)	Apply preventatively prior to snow cover. For snow mold, do not make more than three applications at 1.45 fl.oz. / 1000 sq.ft.. Begin applications when conditions are favorable for disease infection and prior to disease symptom development. Repeat application after 14 to 21 days.
Leaf Rust Stem Rust Stripe Rust <i>(Puccinia spp.)</i>	Begin applications when conditions are favorable for disease infection and prior to disease symptom development. Repeat application after 14 to 21 days.

*(continued)*

**GOLF COURSE TURF APPLICATIONS AND INSTRUCTIONS (continued)**

TARGET DISEASES	APPLICATION DIRECTIONS
Leaf Spot ( <i>Bipolaris sorokiniana</i> )	Apply when disease conditions exist. Repeat application after 14 to 21 days.
Melting Out ( <i>Dechslera poae</i> )	Apply when disease conditions exist. Repeat application after 14 to 21 days.
Necrotic Ring Spot ( <i>Ophiosphaerella korrae</i> )	Apply when disease conditions exist. Repeat application after 14 to 21 days.
Pink Patch ( <i>Limonomyces roseipellis</i> )	Apply when disease conditions exist. Repeat application after 14 to 21 days.
Powdery Mildew ( <i>Erysiphe graminis</i> )	Begin applications when conditions are favorable for disease infection, prior to disease symptom development. Repeat application after 14 to 21 days.
Pythium Blight Pythium Root Rot ( <i>Pythium aphanidermatum</i> , <i>Pythium</i> spp.)	Use preventatively. Begin applications when disease conditions exist and prior to disease symptom development. During periods of prolonged favorable conditions, treat on the 10-day application interval. For use on newly seeded as well as established turf. Repeat application after 14 to 21 days.
Red Thread ( <i>Laetisaria fuciformis</i> )	Apply when disease conditions exist. Repeat application after 14 to 21 days.
Southern Blight ( <i>Sclerotium rolfsii</i> )	Apply when disease conditions exist. Repeat application after 14 to 21 days.
Spring Dead Spot ( <i>Ophiosphaerella korrae</i> ) or ( <i>Ophiosphaerella narmari</i> ) or ( <i>Ophiosphaerella herpotricha</i> )	Apply 1 or 2 applications approximately one month prior to bermudagrass dormancy. Irrigate with ¼" to ½" of water directly after application. Repeat application after 14 to 21 days.
Summer Patch ( <i>Magnaporthe poae</i> )	Apply when disease conditions exist. Repeat application after 14 to 21 days.
Take-all Patch ( <i>Gaeumannomyces graminis</i> var. <i>avenae</i> )	Make two applications at 28-day intervals in the spring and two applications at 28-day intervals in the fall.



# STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a dry, temperature-controlled, secure place.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

## CONTAINER HANDLING:

**Containers < 5 Gallons: Nonrefillable container.** Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Containers  $\geq$  5 Gallons: Nonrefillable container.** Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Bulk Containers: Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When the container is empty, replace the cap and seal all openings that have been opened during use and return to the point of purchase or to a designated location named at the time of purchase of this product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged or leaking, call CHEMTREC at (800) 424-9300. If not returned to the point of purchase or to a designated location, clean empty container as instructed above and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

If the container is damaged and leaking or material has been spilled, follow these procedures:

- Cover spill with absorbent material.
- Sweep into disposal container.
- Wash area with detergent and water and follow with clean water rinse.
- Do not allow to contaminate water supplies.
- Dispose of according to instructions.

**THIS CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

## **WARRANTY AND LIMITATION OF DAMAGES**

**Conditions of Sale:** To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. Sipcam Agro USA, Inc. disclaims all other warranties, express or implied. To the extent consistent with applicable law, Sipcam Agro USA, Inc. shall not be liable for consequential, special, or indirect damages resulting from the use or handling of this product, and Sipcam Agro USA, Inc.'s sole liability and buyer's and user's exclusive remedy shall be limited to the refund of the purchase price. Buyer and user acknowledge and assume all risks and liability resulting from handling, storage and use of this product. Sipcam Agro USA, Inc. does not authorize any agent or representative to make any other warranty, guarantee or representation concerning this product.

# Adrenalyn™ ETQ

Turf Fungicide

Tebuconazole	GROUP 3	FUNGICIDE
Azoxystrobin	GROUP 11	FUNGICIDE



ACTIVE INGREDIENTS:	% BY WT
Azoxystrobin: Methyl (E)-2-[[6-(2-cyanophenoxy-4-pyrimidinyl)alpha-methoxymethylene] benzeneacetate	18.2%
Tebuconazole: (+)-alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl-1H-1,2,4-triazole-1-ethanol	27.3%
<b>OTHER INGREDIENTS:</b>	54.5%
<b>TOTAL:</b>	100.0%

Adrenalyn ETQ Turf Fungicide is a suspension concentrate fungicide containing 2.55 lbs. Tebuconazole and 1.69 lbs. Azoxystrobin per gallon.

## KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Emergency Phone Numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 222-1222 Poison Control Center
-------------------------	---

See additional Precautionary Statements and Directions for Use inside the booklet.

NET CONTENTS: 1 Gallon

Manufactured for:  
**SIPCAM AGRO USA, INC.**  
2525 Meridian Parkway  
Durham, NC 27713

# 1G

EPA Reg. No. 60063-62

EPA Est. No.: 60063-GA-1 (Lot number begins with VL)

EPA Est. No.: 70815-GA-1 (Lot number begins with CB)

EPA Est. No.: 86555-MO-1 (Lot number begins with AF)

EPA 20200730 (08/20)



READ THE  
ENTIRE LABEL  
BEFORE USING  
THIS PRODUCT