

AgriPhage™-CMM

BACTERICIDE

Bactericide for use on tomatoes
Biological control for bacterial canker on tomatoes

FOR ORGANIC PRODUCTION

ACTIVE INGREDIENTS:

Bacteriophage active against *Clavibacter michiganensis*
subsp. *michiganensis** 0.0005%

OTHER INGREDIENTS: 99.9995%

Total: 100.0000%

*Contains at least 3.8 X 10¹² plaque-forming units (PFU) per gallon of product
(Contains at least 9.46 x 10¹¹ plaque-forming units (PFU) per quart of product,
Contains at least 1.0 x 10¹² plaque-forming units (PFU) per liter of product).

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See attached booklet for First Aid, additional Precautionary Statements and Directions for Use.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store this product at 39.2°F (4°C). Direct sunlight and/or extreme heat will cause degradation of the product and decrease product efficacy.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

EPA Reg. No. 67986-6-70051

EPA Est. No. 67986-UT-001

Manufactured by:



9075 South Sandy Parkway
Sandy, Utah 84070
801-746-3600
www.omnilytics.com

Not for sale or use after:

Manufactured for
Certis USA L.L.C.
9145 Guilford Road, Suite 175
Columbia, MD 21046



ESL20191202
rev20200411

This is a Specimen Label. It may not reflect the most-recent approved label for use in your state. Always refer to the label on the product packaging for approved use instructions. Please contact your Certis sales representative for more information.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Hotline Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For non-emergency information concerning this product, call the National Pesticide Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8:00 a.m. to 12:00 p.m. Pacific Time (NPIC Web site: www.npic.orst.edu). For emergencies, call your poison control center at 1-800-222-1222.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Waterproof gloves
- Shoes plus socks

Mixers/loaders and applicators must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air-purifying respirator with an HE filter. Repeated exposures to high concentrations of microbial proteins can cause allergic sensitization.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

User should:

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For Terrestrial Uses: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

CHEMIGATION: Do not apply this product through any irrigation system.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow workers entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Do not enter or allow workers to enter the treated greenhouse or enclosed space until the ventilation requirements in 40 CFR 170.405(b)(3) have been met and the REI of 4 hours has expired. Until then, only handlers wearing the appropriate personal protective equipment can enter the greenhouse or enclosed space.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water) is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

Product Information

AgriPhage-CMM is a bactericide that is comprised of a formulation of bacteriophage to be used as a preventive and curative product for the control of bacterial canker (caused by the bacterium *Clavibacter michiganensis* subsp. *michiganensis*) on tomatoes. Apply AgriPhage-CMM as a preventive to protect growing leaf tissue, as a curative when the first disease symptoms are visible, or when conditions are conducive to heavy disease pressure. In the greenhouse, apply AgriPhage-CMM as a foliar spray or by fogging. In the field, apply AgriPhage-CMM with conventional ground spray equipment immediately after planting and as the plants mature. See the Mixing and Application Instructions section of this label for additional details.

Mixing and Application Instructions

- MIX WELL PRIOR TO USE -

COMPATIBILITY: Do not combine AgriPhage-CMM in the spray tank with denaturing agents or copper salts. Apply AgriPhage-CMM in approved tank mixes or in an alternating spray program, applying denaturing agents or copper salts 4 days after AgriPhage-CMM application, as copper salts and denaturing agents compromise bacteriophage survival. AgriPhage-CMM is compatible with several commonly used fungicides, liquid fertilizers, herbicides, and insecticides, but has not been fully evaluated with all of these. To ensure compatibility of tank-mix combinations, please consult with an OmniLytics representative prior to tank mixing. AgriPhage-CMM cannot be mixed with another product with a prohibition against mixing. Use of the tank mix must be in accordance with the more restrictive label limitations and precautions.

Greenhouse Seedling Treatment

RATE	PRE-HARVEST INTERVAL (PHI)	APPLICATION NOTES
1 pint of AgriPhage-CMM per 12-25 gallons of water per 9600 square feet of greenhouse space	AgriPhage-CMM may be applied up to and including the day of harvest.	Apply daily treatments to foliage by spraying or fogging. Begin applications to seedlings, (at the 4-leaf stage), immediately after planting, as a preventive to protect the growing leaf tissue or as a curative when the first disease symptoms are visible. Use sufficient spray solution to ensure complete coverage. Avoid excessive amounts of spray solution that result in runoff.

Greenhouse Treatment – For Tomatoes Grown Hydroponically

RATE	PRE-HARVEST INTERVAL (PHI)	APPLICATION NOTES
½ to 1 quart of AgriPhage-CMM per 12-25 gallons of water per 9600 square feet of greenhouse space	AgriPhage-CMM may be applied up to and including the day of harvest.	Apply to foliage by spraying or fogging. Begin applications for foliar applications as a preventive to protect the growing leaf tissue or as a curative when the first disease symptoms are visible. Continue throughout the growing season. Begin at the lower application rate of ½ quart of AgriPhage-CMM per 9600 square feet of greenhouse space when plants are small. As plants increase in size and foliage, increase application rate to 1 quart of AgriPhage-CMM per 9600 square feet of greenhouse space to ensure adequate coverage. Repeat application 1-3 times per week. Use sufficient spray solution to ensure complete coverage. Avoid excessive amounts of spray solution that result in runoff. Under severe disease conditions, use the higher application rate of 1 quart of AgriPhage-CMM per 9600 square feet of greenhouse space.

(continued)

Field Treatment – Ground Application

RATE	PRE-HARVEST INTERVAL (PHI)	APPLICATION NOTES
1 to 2 pints of AgriPhage-CMM per 50-100 gallons of water per acre	AgriPhage-CMM may be applied up to and including the day of harvest.	Begin applications as a preventive to protect the growing leaf tissue or as a curative when the first disease symptoms are visible. Continue throughout the growing season. Begin at the lower application rate of 1 pint of AgriPhage-CMM per acre at planting (below the second tie). As plants increase in size and foliage (at or above the second tie), increase application rate to 2 pints of AgriPhage-CMM per acre to ensure adequate coverage. Repeat application 1-3 times per week. Use sufficient spray solution to ensure complete coverage. Avoid excessive amounts of spray solution that result in runoff. Under severe disease conditions and periods of heavy rain, apply immediately after rainfall, and use the higher application rate of 2 pints of AgriPhage-CMM per acre.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store this product at 39.2°F (4°C). Direct sunlight and/or extreme heat will cause degradation of the product and decrease product efficacy.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty Statement

OmniLytics, Inc. warrants that this material conforms to the description on the label and is reasonably fit for the purposes referred to in the Directions for Use. To the extent consistent with applicable law, OmniLytics, Inc. makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty. To the extent consistent with applicable law, in no case or circumstance shall OmniLytics, Inc. or seller be liable for consequential, special or indirect damages resulting from the use or handling of this product including but not limited to, loss of profits, business reputations, or customers; labor cost; or any other expenses incurred in planting, cultivating or harvesting.

Specimen Label



ESL20191202
rev20200411

AgriPhage™-CMM

BACTERICIDE

Bactericide for use on tomatoes
Biological control for bacterial canker on tomatoes

Batch Code:

FOR ORGANIC PRODUCTION

ACTIVE INGREDIENTS:

Bacteriophage active against *Clavibacter michiganensis*
subsp. *michiganensis** 0.0005%

OTHER INGREDIENTS: 99.9995%

Total: 100.0000%

*Contains at least 3.8 X 10¹² plaque-forming units (PFU) per gallon of product
(Contains at least 9.46 x 10¹¹ plaque-forming units (PFU) per quart of product,
Contains at least 1.0 x 10¹² plaque-forming units (PFU) per liter of product).

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See attached booklet for First Aid, additional Precautionary Statements
and Directions for Use.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing.
Wash thoroughly with soap and water after handling and before eating, drinking,
chewing gum, using tobacco, or using the toilet.

Net Contents: 1 qt.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store this product at 39.2°F (4°C). Direct sunlight and/or
extreme heat will cause degradation of the product and decrease product
efficacy.

Pesticide Disposal: Wastes resulting from the use of this product must be dis-
posed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.
Triple rinse container (or equivalent) promptly after emptying. Triple rinse as
follows: Empty the remaining contents into application equipment or a mix
tank and drain for 10 seconds after the flow begins to drip. Fill the container
¼ full with water and recap. Shake for 10 seconds. Pour rinsate into applica-
tion equipment or a mix tank, or store rinsate for later use or disposal. Drain
for 10 seconds after the flow begins to drip. Repeat this procedure two more
times. Then, offer for recycling if available or reconditioning if appropriate, or
puncture and dispose of in a sanitary landfill or by incineration, or if allowed by
state and local authorities, by burning. If burned, stay out of smoke.

EPA Reg. No. 67986-6-70051

EPA Est. No. 67986-UT-001

Manufactured by:



9075 South Sandy Parkway
Sandy, Utah 84070
801-746-3600
www.omnilytics.com

Not for sale or use after:

Manufactured for
Certis USA L.L.C.
9145 Guilford Road, Suite 175
Columbia, MD 21046



ESL20191202
rev20200411

PEEL BACK LEAFLET HERE

Item: 157011