



# ALL-SEASON BRUSH NO-MORE

Controls multiflora rose, brambles, cedars, locusts, poison oak, poison ivy, honeysuckle, thistles, kochia, kudzu, and many other trees, vines and broadleaf weeds.

#### **ACTIVE INGREDIENTS:**

| 2,4-D, 2-ethylhexyl ester    | 9.74%  |
|------------------------------|--------|
| 2,4-DP-p, 2-ethylhexyl ester | 4.78%  |
| Dicamba                      | 1.65%  |
| INERT INGREDIENTS:           | 83.83% |
| TOTAL 1                      | 00.00% |

#### This product contains:

0.49 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.46%. 0.24 lb. (+)-R-2-(2,4-dichlorophenoxy) propionic acid equivalent per gallon or 3.23%.

0.12 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.65%

Contains petroleum distillates.

Isomer Specific by AOAC Method

## WARNING



READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

#### PRECAUTIONARY STATEMENTS

#### **Hazards to Humans and Domestic Animals**

**WARNING:** Causes substantial but temporary eye injury. Causes skin irritation. Harmful if swallowed. Harmful if absorbed through the skin. Do not get in eyes, on skin, or on clothing.

#### Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are barrier laminate, nitrile rubber, neoprene rubber, or Viton. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear:

- coveralls over short-sleeved shirt and short pants,
- chemical-resistant footwear and socks,
- chemical-resistant gloves (category E),
- goggles or face shield,
- chemical-resistant headgear for overhead exposure, and
- chemical-resistant apron for mixing, loading, or cleaning spills or equipment or otherwise exposed to the concentrate.

#### **User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **User Safety Recommendations**

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product.
   Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

| First Aid                  |   |
|----------------------------|---|
| If in eyes:                | <ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>         |
| If on skin<br>or clothing: | <ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>   |
| If swallowed:              | <ul> <li>Immediately call a poison control center or doctor.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.</li> </ul> |
| If inhaled:                | Move person to fresh air.     If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.     Call a poison control center or doctor for treatment advice.   |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

**Note to Physician:** May pose an aspiration pneumonia hazard. Contains petroleum distillates.

#### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

#### **Physical or Chemical Hazards**

Combustible. Do not use or store near heat or open flame.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### 1. Precautions and Restrictions

- Do not enter or allow people (or pets) to enter the treated area until sprays have dried. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not apply this product through any type of irrigation system.
- Do not apply when temperatures exceed 85°F and humidity is high.

#### 2. Noncropland Sites

Gordon's® All-Season Brush-No-More® is recommended to control undesirable woody plants established in noncropland. It is effective for buckbrush, poison ivy, multiflora rose, and sumac established in the uncultivated areas presented on next page:

### Uncultivated agricultural areas and uncultivated nonagricultural areas:

#### A. Recommended Noncropland Sites.

- Barrier strips
- Farmyards
- Fencerows or fence lines
- Firebreaks
- Highway rights-of-way (principal, interstate, county, private, and unpaved roads): Roadsides, road shoulders, road embankments, dividers, and medians.
- Industrial sites: Lumberyards, tank farms, fuel or equipment storage areas.
- Municipal, state, and federal lands: Airports and military installations
- · Railroad rights-of-way
- Recreation areas: Fairgrounds, golf courses, parks, and areas adjacent to athletic fields.
- Utility rights-of-way: Telephone, pipeline, electrical powerlines, and communication transmission lines

#### B. Prohibitions for Noncropland Sites.

- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (noncropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply to wetlands (swamps, bogs, potholes, or marshes).
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals.
- Do not apply to agricultural drainage water or on agricultural ditchbanks.
- Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

#### 3. Application Timing for Mixed Brush

Spraying with ground equipment can be effective throughout the growing season from full leaf to leaf drop for mixed brush. Sprays should be applied during warm weather when brush are young and actively growing. All leaves, stems, and shoots should be thoroughly wetted to the ground. Do not cut brush until the herbicide has translocated throughout the plant causing root death.

Cut stump treatments of Gordon's® All-Season Brush-No-More® are appropriate during the dormant period before bud growth or any signs of active growth of the mixed brush.

#### **BRUSH CONTROLLED**

American chestnut Cherry Pine Cottonwood Ash Poplar Aspen Dogwood Sassafras Barberry Shortleaf pine Elm Birch Gooseberry Spruce Blackberry Honey locust Sumac Black cherry Honeysuckle Sycamore Black locust Kudzú Trumpetcreeper Brambles Maple Wild plum Buckbrush Multiflora rose Willow Cedar Oak

#### 4. Spray Drift Management

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

#### **Droplet Size**

Use only Medium or coarser spray nozzles according to ASAE (S 572) definition of standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

#### Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.

#### **Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable

atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

#### Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

#### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed. 2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

#### Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 feet above the target site. Aerial applications of this product are prohibited.

#### 5. Foliar Applications:

#### High volume foliar applications:

Apply a 2.0 to 8.0% vol/vol spray solution as a full cover spray with high volume equipment. Use the lower spray concentrations in the range for susceptible species and use the higher spray concentrations within the range for hard-to-control species, for mature plants during the late summer or under adverse environmental conditions (e.g. drought).

Spray woody plants or mixed brush uniformly and thoroughly by wetting all leaves, stems, bark and root collars. The total volume of spray solution required for adequate coverage of solid stands of mixed brush can range from 2 to 8 gallons of spray solution per treated 1000 sq. ft. The spray preparation chart for applications on a spray-to-wet basis is shown below in Table 1.

The maximum seasonal application rate for trees, brush and woody plant control is 8.0 gallons of product (4.0 lb. 2,4-D ae) per acre per application per site.

## For Backpack Sprayers, Knapsack Sprayers, and Hand-pressurized Pump Sprayers

**Table 1.** Instructions for preparing 1 to 3 gallons of spray solution at 2.0 to 8.0% spray concentration with water for high volume foliar applications.

| - 1 |                     |   |                  |            |            |
|-----|---------------------|---|------------------|------------|------------|
|     | Gallons<br>Of Water | Amount Of Product Needed for<br>Spray Concentration of: |                  |            |            |
|     | Oi water            | 2.0%  | 2.7 %            | 4.0 %      | 8.0 %      |
|     | 1                   | 5<br>tablespoons  | 7<br>tablespoons | 5 fl. oz.  | 10 fl. oz. |
|     | 2                   | 5 fl. oz.   | 7 fl. oz.        | 10 fl. oz  | 20 fl. oz. |
|     | 3                   | 8 fl. oz.   | 10 fl. oz.       | 15 fl. oz. | 30 fl. oz. |
|     |                     |   |                  |            |            |

Equal measures: 1 fl. oz. = 2 Tablespoons (Tbs.) = 6 Teaspoons (tsp.)

| Target<br>species | Application schedule                      | Maximum<br>application<br>rate, gallons<br>of product<br>per acre | Maximum<br>application rate,<br>pounds of 2,4-D<br>acid equivalent<br>per acre per<br>application | Maximum<br>number of<br>applications<br>per year | Minimum<br>days<br>between<br>applications |
|-------------------|---|---|---|--|--|
| Woody<br>plants   | Broadcast<br>and high<br>volume<br>foliar | 8.0 gal./A  | 4.0 lb./A   | 1  | NA   |

#### 6. Individual Plant Treatments:

**Cut Surface - Stump Treatment -** This method is most effective and economical on stumps with diameters larger than 3 to 4 inches. This treatment can be applied throughout the year except when snow, ice, or water prevents thorough spray coverage.

For Backpack Sprayers, Knapsack Sprayers, and Hand Pump Sprayers - Mix 20.0 fluid ounces of Gordon's® Cleanout® Brush and Stump Control with 1.0 gallon of oil (diesel oil, No. 1 or No. 2 fuel oil,

kerosene, mineral oil, or other oil blends formulated for basal applications). Penetrants appropriate for oil soluble herbicides may improve control. Refer to Table 2 for spray preparation. Spray thoroughly the cut surfaces, bark, and exposed roots. Treat entire circumference of the tree

Limitations for cut surface (stump) treatments - Use only one cut surface application per year. The maximum seasonal application rate for trees, brush and woody plant control is 8.0 gallons of product (4.0 lb. 2,4-D ae) per acre per application per site.

**Table 2.** Quick mix instructions for preparing 1 to 5 gallons of spray solutions with oil for cut surface applications.

| Spray Solution,<br>Gallons | Amounts of Gordon's All-Season<br>Brush-No-More required,<br>Fluid Ounces |  |
|----------------------------|---|--|
| 1 gal.                     | 20 fluid ounces (1.25 pints)  |  |
| 2 gal.                     | 40 fluid ounces (2.5 pints)   |  |
| 3 gal.                     | 60 fluid ounces (3.75 pints)  |  |
| 5 gal.                     | 100 fluid ounces (6.25 pints)   |  |
|                            |   |  |

Equal Measures: 16 fluid ounces = 1 pint = 1/2 quart = 2 cups

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal **PESTICIDE STORAGE:** Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse (or pressure rinse) container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

#### OR

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et.al. v. EPA, C01-0132C, (W.O. WA). For further information, please refer to EPA Web Site: http://www.epa.gov/espp.

#### LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

BRUSH-NO-MORE®, CLEANOUT® and GORDON'S® are registered trademarks of PBI/GORDON CORPORATION.

783/2-2010 AP102909 EPA REG. NO. 2217-775-33955



An Employee-Owned Company

MANUFACTURED BY PBI/GORDON CORPORATION 1217 WEST 12TH STREET KANSAS CITY, MISSOURI 64101 www.pbigordon.com