

SPECIMEN

Armezon[®]**Herbicide**

For postemergence weed control in all types of field corn, popcorn, sweet corn, sugarcane, and between crop applications

Active Ingredient:

topramezone: [3-(4,5-dihydro-isoxazolyl)-2-methyl-4-(methylsulfonyl)phenyl](5-hydroxy-1-methyl-1*H*-pyrazol-4-yl)methanone 29.7%

Other Ingredients: 70.3%

Total: 100.0%

1 gallon contains 2.8 pounds of topramezone free acid.

EPA Reg. No. 7969-262

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents:

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after first 5 minutes; then continue rinsing. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything to an unconscious person.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).</p>	

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber (includes natural rubber blends and laminates) ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. **DO NOT** apply this product through any type of irrigation system.

Product must be used in a manner which will prevent back-siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures, or rinsate.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at time of herbicide application.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Observe all precautions and limitations in this label and the labels of products used in combination with **Armezon® herbicide**. The use of **Armezon** not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions, and **Conditions of Sale and Warranty** are to be followed.

BASF Corporation does not recommend or authorize the use of this product in manufacturing, processing, or preparing custom blends with other products for application in crops.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber (includes natural rubber blends and laminates) \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils
- Shoes plus socks

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store product in original container only. Store product in a cool, dry place. **DO NOT** store this product under wet conditions. If this product has been stored where freezing temperatures have occurred, agitate or mix contents of container well before use. Avoid cross-contamination with other pesticides.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity \leq 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity $>$ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

(continued)

STORAGE AND DISPOSAL *(continued)*

Container Handling *(continued)*

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

Spills

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

Steps to take if this material is released into the environment or spilled:

- Wear **Personal Protective Equipment (PPE)** and avoid exposure when managing a spill. (See **Precautionary Statements** section of this label for required PPE.)
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing, and wash affected skin areas with soap and water. Wash clothing before reuse.
- Keep spill out of all sewers and open bodies of water.

Product Information

Armezon® herbicide is a suspension concentrate (SC) herbicide providing systemic postemergence control or growth suppression of emerged broadleaf and grass weeds in field corn (grown for grain, silage, or seed), popcorn (grown for ear, kernel, or seed), sweet corn (grown for ear, kernel, or seed), sugarcane, and between crop applications. This product may be used on conventional and herbicide-resistant/tolerant corn hybrids. BASF has not tested all inbred lines for tolerance to **Armezon**. Before using **Armezon**, refer to seed company instructions for use on inbred lines of field corn, popcorn, and sweet corn.

When applied postemergence as directed, **Armezon** will control or suppress the broadleaf weeds listed in **Table 1** and the grass weeds listed in **Table 2**.

To increase weed control spectrum for use in corn, tank mix **Armezon** with 0.25 lb to 1.5 lbs active ingredient of atrazine herbicide per acre. Use the lower rates of atrazine for added burndown of emerged weeds and the higher labeled rates for added weed residual control.

Armezon applications must include spray additives. See **Additives** and **Mixing Order** for details.

Table 1. Broadleaf Weeds Controlled Postemergence Broadcast
(including ALS-resistant¹, glyphosate-resistant, and triazine-resistant biotypes)

Common Name	Scientific Name	Armezon [®] herbicide Application Rate	
		0.5 fl oz/A	0.75 to 1.0 fl oz/A
		Maximum Weed Size ² (inches)	
Amaranth, Palmer	<i>Amaranthus palmeri</i>	4	6
Amaranth, Powell	<i>Amaranthus powellii</i>	4	6
Burcucumber	<i>Sicyos angulatus</i>	4	6
Canola, volunteer	<i>Brassica</i> spp.	4	6
Carpetweed	<i>Mollugo verticillata</i>	4	6
Chickweed, common	<i>Stellaria media</i>	2	4
Cocklebur, common	<i>Xanthium strumarium</i>	5	8
Dandelion	<i>Taraxacum officinale</i>	–	6*
Galinsoga, hairy	<i>Galinsoga ciliata</i>	4	6
Henbit	<i>Lamium amplexicaule</i>	3	4
Horseweed (Marestail)	<i>Conyza canadensis</i>	4	6
Jimsonweed	<i>Datura stramonium</i>	4	6
Kochia	<i>Kochia scoparia</i>	4	6
Lambsquarters, common	<i>Chenopodium album</i>	4	6
Lettuce, prickly	<i>Latuca serriola</i>	2	4
Mallow, common	<i>Malva neglecta</i>	2	3
Mallow, Venice	<i>Hibiscus trionum</i>	2*	3*
Morningglory	<i>Ipomoea</i> spp.	4*	6*
Mustard	<i>Brassica</i> spp.	4	6
Nightshade, black	<i>Solanum nigrum</i>	4	6
Nightshade, Eastern black	<i>Solanum ptycanthum</i>	4	6
Nightshade, hairy	<i>Solanum sarrachoides</i>	4	6
Pigweed, prostrate	<i>Amaranthus blitoides</i>	4	6
Pigweed, redroot	<i>Amaranthus retroflexus</i>	4	6
Pigweed, smooth	<i>Amaranthus hybridus</i>	4	6
Pigweed, tumble	<i>Amaranthus album</i>	2	4
Pusley, Florida	<i>Richardia scabra</i>	2	3
Ragweed, common	<i>Ambrosia artemisiifolia</i>	4	6
Ragweed, giant	<i>Ambrosia trifida</i>	5	8
Shepherd's-purse	<i>Capsella bursa-pastoris</i>	2	4
Sida, prickly	<i>Sida spinosa</i>	2	3
Smartweed, ladysthumb	<i>Polygonum persicaria</i>	2	3
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	2	3
Sunflower, volunteer	<i>Helianthus</i> spp.	5	8
Sunflower, wild (common)	<i>Helianthus annuus</i>	5	8
Thistle, Canada	<i>Cirsium arvense</i>	4*	6*
Thistle, Russian	<i>Salsola iberica</i>	2	4
Velvetleaf	<i>Abutilon theophrasti</i>	4	8
Waterhemp, common	<i>Amaranthus rudis</i>	4	6
Waterhemp, tall	<i>Amaranthus tuberculatus</i>	4	6

¹ ALS (acetolactate synthase)-resistant weeds include those weeds resistant to imidazolinone, sulfonamide, and/or sulfonylurea herbicides.

² For best performance, spray before weeds exceed the maximum stem height or vine length listed in this table.

* Partial control or suppression

Table 2. Grass Weeds Controlled or Suppressed² Postemergence Broadcast
(including ALS-resistant¹, glyphosate-resistant, and triazine-resistant biotypes)

Common Name	Scientific Name	Maximum Weed Leaf Stage ³	Maximum Weed Size ³ (inches)
Barnyardgrass	<i>Echinochloa crus-galli</i>	4	4
Crabgrass, large	<i>Digitaria sanguinalis</i>	4	3
Crabgrass, smooth	<i>Digitaria ischaemum</i>	4	3
Cupgrass, woolly	<i>Eriochloa villosa</i>	3*	3*
Foxtail, giant	<i>Setaria faberi</i>	4	4
Foxtail, green	<i>Setaria viridis</i>	3*	3*
Foxtail, yellow	<i>Setaria lutescens</i>	3*	3*
Goosegrass	<i>Eleusine indica</i>	4	3
Johnsongrass, seedling	<i>Sorghum halepense</i>	3*	4*
Millet, wild proso	<i>Panicum miliaceum</i>	3	3
Panicum, fall	<i>Panicum dichotomiflorum</i>	3*	3*
Shattercane	<i>Sorghum bicolor</i>	3*	4*
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>	3*	3*

¹ ALS-resistant weeds include those weeds resistant to imidazolinone and/or sulfonylurea herbicides.

² Growth suppression at 0.5 fl oz/A

³ For best performance, spray before grass exceeds the maximum leaf stage and/or height listed in this table.

* Growth suppression at 0.75 fl oz/A; control at 1.0 fl oz/A

Mode of Action

Armezon® herbicide is absorbed by leaves, roots, and shoots and translocated to the growing points of sensitive weeds to control emerged weeds. **Armezon** controls weeds by inhibiting carotenoid biosynthesis (HPPD-inhibitor **Group 27**). Temperatures and moisture conditions for active plant growth are important for optimum **Armezon** activity. **Armezon** application to weeds during periods of stress conditions, such as cold temperatures and/or drought, may result in reduced performance.

Herbicide Resistance Management

Resistance to **Armezon** or cross-resistance to other HPPD-inhibitor herbicides is known to exist. Repeated applications of a single mode of action in a weed management plan increase the probability of selecting for naturally occurring biotypes* with less susceptibility to herbicides using that mode of action. Therefore, tank mix **Armezon** with a herbicide having a different mode of action and/or use in a rotation with herbicides having a different mode of action. Other HPPD-inhibitor herbicides include **Balance® Flexx herbicide** (isoxaflutole), **Callisto® herbicide** (mesotrione), and **Laudis® herbicide** (tembotrione).

* A weed biotype is a naturally occurring individual within a given species that has a slightly different but distinct genetic makeup from other plants.

Crop Tolerance

Apply **Armezon** during favorable growing conditions for optimum crop tolerance and weed control. Crops under environmental stress are more likely to show injury from any herbicide application. Rarely, plants under these

conditions treated with **Armezon** may show transient bleaching of the portion of the leaves intercepting the spray application. These symptoms are temporary and occur infrequently; crop growth is not affected.

Cultivation

Avoid disturbing (e.g. cultivation) treated areas for at least 7 days following an application of **Armezon** to allow best herbicide uptake, translocation, and weed control.

Insecticide Information

Armezon may be used sequentially or in combination with soil or foliar applied insecticides registered for use in corn.

Application Instructions

Armezon is effective for post emergence control of annual weeds in conservation or conventional tillage production systems.

DO NOT apply **Armezon** within 30 feet of the downwind edge of native plant communities.

The applicator is responsible for any loss or damage that results from spraying **Armezon** in a manner other than directed in this label. In addition, applicator must follow all applicable state and local regulations and ordinances for spraying.

Application Timing

- Apply **Armezon** as a postemergence treatment when weeds are actively growing.
- For optimal weed control, apply **Armezon** before weeds exceed labeled height.
- Apply **Armezon** a minimum of one hour before rainfall or overhead irrigation.

Ground Application Methods and Equipment

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. Use higher water volumes treating larger weeds and/or high-density weed infestation. Weeds must be thoroughly covered with spray. Dense leaf canopies shelter small weeds and can prevent adequate spray coverage. **Armezon® herbicide** applications should be made with drop nozzles if the crop canopy prevents adequate weed coverage.

Spray Drift

DO NOT apply when weather conditions may cause drift to adjacent crops and vegetation; injury may result if this occurs. To avoid spray drift from treated areas, **DO NOT** make applications when wind speed exceeds 10 mph or during periods of temperature inversions.

Use of larger droplet sizes will reduce spray drift. Agriculturally approved drift-reducing additives may also be used.

Aerial Application Methods and Equipment

Uniformly apply with properly calibrated aerial equipment in 2 or more gallons of water per acre. Adequate spray volume must be used to provide accurate and uniform distribution of spray particles over the treated area and to avoid drift of spray particles to nontarget areas.

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following special aerial use directions and precautions:

- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than 3/4 the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- **DO NOT** spray when wind velocity is greater than 5 mph. Coarse sprays (larger droplets) are less likely to drift.

Additives

Postemergence applications of **Armezon** require the addition of an adjuvant and nitrogen fertilizer for optimum weed control.

Agriculturally approved drift-reducing additives may be used in applications with **Armezon**.

When an adjuvant is to be used with this product, BASF recommends the use of Chemical Producers and Distributors Association certified adjuvant.

Adjuvants

Unless specific tank mix directions are given in **Crop-specific Information**, always use a methylated seed oil (MSO) or a petroleum-based or vegetable seed-based oil concentrate (COC) with **Armezon**. For best performance

across a wide range of environmental conditions, including when weeds are under moisture and/or temperature stress, use an MSO adjuvant. Apply these oilbased adjuvant concentrates at 1.0 to 1.5 gallons per 100 gallons of water (1.0% to 1.5% volume/volume [v/v]). Use the higher rate when applying during periods of hot, dry weather.

AND

Nitrogen Fertilizer

Recommended nitrogen-based fertilizers include urea ammonium nitrate (UAN; 28% to 34%) or ammonium phosphate (10-34-0) at 1.25 to 2.5 gallons per 100 gallons of water (1.25% to 2.5% v/v). Instead of a liquid fertilizer, spray grade ammonium sulfate (AMS) at a minimum rate of 8.5 to 17 pounds per 100 gallons of water may be used. Use the higher rate when applying during periods of hot, dry weather.

Mixing Instructions

For product containers 5 gallons or less, shake well before use. For product containers more than 5 gallons, recirculate before use.

DO NOT use liquid fertilizer as a carrier for postemergence applications. Use only water as a carrier.

1. **Water** - Fill the spray tank 1/2 to 3/4 full with clean water and start agitation.
2. **Armezon** - Add required amount to spray tank while agitating.
3. After the **Armezon** has visibly dispersed, add spray additives and fill the remainder of the tank with water.

Maintain agitation throughout mixing and application until spraying is completed. Limit the amount of spray mixture prepared to that needed for immediate use.

Tank Mixing Order

When tank mixing **Armezon** with recommended herbicides, add the other herbicides and other components in the following order. Maintain agitation throughout mixing and application until spraying is completed. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.

1. **Water** - Fill tank 1/2 to 3/4 full with clean water and start agitation.
2. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
3. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. **Water-soluble additives** (including dry and liquid fertilizers such as AMS or UAN)
5. **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)

- 6. **Water-soluble products**
- 7. **Emulsifiable concentrates** (including MSO adjuvants)
- 8. **Remaining quantity of water**

Cleaning Spray Equipment

To avoid injury to sensitive crops, drain and clean application equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions. Triple rinse equipment before and after applying this product.

Tank Mixing Information

Armezon® herbicide may be applied sequentially or tank mixed with other herbicides as part of a complete weed control program. Tank mix recommendations are for use only in states where the sequential or tank mix product and application site are registered. Refer to **Crop-specific Information** for more details and for specific tank mix restrictions. Read and follow the applicable restrictions and precautions and **Directions For Use** on all products included in any tank mix. **The most restrictive labeling applies to tank mixes.**

Rotational Crop Restrictions

The following rotational crops may be planted after applying **Armezon** at the specified rate. Planting earlier than the specified interval may result in crop injury. Avoid overapplication by minimizing overlap of spray swaths and by switching off spray boom when turning (end rows).

For rotational crops following the use of sequential applications of **Armezon**, the rotational interval begins after the last **Armezon** application.

Rotational Crop	Rotational Interval (months)		
	Armezon Application Rate (fl oz/A)		
	0.5	0.75	1.0 to 2.0
Corn (all)	0	0	0
Sugarcane	0	0	0
Cereal grains	3	3	3
Grass grown for seed	3	3	3
Rice	3	3	3
Alfalfa	9	9	9
Cotton	9	9	9
Grain sorghum	9	9	9
Peanut	9	9	9
Potato	9	9	9
Soybean	9	9	9
Sunflower	9	9	9
Canola	9	9	18

(continued)

Rotational Crop (continued)	Rotational Interval (months)		
	Armezon Application Rate (fl oz/A)		
	0.5	0.75	1.0 to 2.0
Lima bean, succulent	9	9	18
Dry beans¹	9	9*	18**
Pea	9	9	18**
Snap/Garden bean	9	9*, †	18
Sugar beet	9*	9*	18**
All Other Crops	18	18	18

¹ For cranberry beans in Idaho, Utah, and the area east of the Cascade Mountains in Oregon and Washington, follow the guidelines for snap/garden bean.

* 18 months for the following states: Colorado, Michigan, Minnesota, Montana, Nebraska (west of Highway 83), North Dakota, South Dakota, Wisconsin, and Wyoming

** 9 months for Idaho, Oregon, and Washington

† 18 months for Idaho, Utah, and the area east of the Cascade Mountains in Oregon and Washington.

Crop-specific Information

Corn (field corn, popcorn, seed corn, sweet corn)

Apply **Armezon** postemergence on all corn types including conventional, **Clearfield®**, **Roundup Ready®**, and **LibertyLink®** hybrids. In addition, **Armezon** may be applied on inbred lines used in field corn, popcorn and sweet corn seed production. Refer to seed company instructions before use on inbred lines.

Armezon may be used in tank mixes or sequential applications with other herbicides registered for use in corn. If **Armezon** is tank mixed with other herbicides, follow label restrictions for the most restrictive tank mix products. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Corn Restrictions

- **DO NOT** apply more than 1.0 fl oz/A **Armezon** (0.022 lb topramezone per acre) per year in corn.
- In the event of a crop loss because of weather, any corn type can be replanted following an application of **Armezon**. If **Armezon** was tank mixed with other herbicides, the label restrictions for these herbicides must also be followed.
- **DO NOT** apply **Armezon** within 45 days of corn harvest (fresh market sweet corn, silage, fodder, or grain) or after the V8 stage of corn growth, whichever comes first.

- **DO NOT** graze or feed treated corn forage, silage, fodder, or grain for at least 45 days after an application of **Armezon® herbicide**.

Tank Mixes

Armezon may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **G-Max Lite™ herbicide**
- **Guardman Max® herbicide**
- **Outlook® herbicide**
- **Prowl® H2O herbicide**
- **Status® herbicide**
- atrazine
- glyphosate (e.g. **Roundup® herbicide**)

Armezon tank mixes or sequential applications with products containing mesotrione are not recommended.

Sequential Herbicide Combinations and Uses

In addition to control of many emerged broadleaf weeds, **Armezon** controls or suppresses growth of several emerged grass weed species. To target a broader spectrum of grasses, use **Armezon** as a sequential postemergence treatment following a preemergence grass herbicide such as **Guardman Max**, **Outlook**, or **Prowl H2O**. **Armezon** may also be used in sequential programs with registered burndown herbicides.

Armezon may be applied in sequence with products containing isoxaflutole (e.g. **Balance® Flexx herbicide**) if the isoxaflutole rate used is less than or equal to 0.0625 lb active ingredient per acre (equal to 4 fl ozs/A of **Balance Flexx**).

Between Crop Application (Fallow)

Armezon may be used as a foliar application to control emerged broadleaf and grass weeds at any time of the year during the fallow period after crop harvest and before the following crop is planted. For rotational crops following the use of a between crop application of **Armezon**, the rotational interval begins after the last **Armezon** application; see **Rotational Crop Restrictions** section for intervals.

Application Rate and Timing

Apply **Armezon** as a broadcast spray at 0.5 fl oz/A to 2.0 fl ozs/A. Best product performance is obtained when weeds are small and actively growing. Thorough coverage of existing weeds is essential, and higher spray volume may be needed for best performance. Sequential applications may be made with a minimum of 14 days between applications. **DO NOT** apply more than the maximum cumulative amount of 2.0 fl ozs/A of **Armezon** per year.

Sugarcane

Armezon can be applied to plant cane or sugarcane grown from stubble (ratoon). **Armezon** may be used in tank mixes or sequential applications with other herbicides registered for use in sugarcane such as atrazine, metribuzin, or **Prowl H2O**. If **Armezon** is tank mixed with other herbicides, follow label restrictions for the most restrictive tank mix product. Application of **Armezon** may cause transient discoloration, chlorosis, or yellowing of sugarcane.

Armezon may be applied between growing seasons as either an early preplant in plant cane or post harvest in ratoon cane prior to cane initiating regrowth. Apply 0.5 to 2.0 fl ozs/A of **Armezon** with a minimum of 14 days between sequential applications. **DO NOT** apply more than 4.0 fl ozs/A of **Armezon** per year.

Special Weeds Controlled/Suppressed

In addition to the weeds controlled early postemergence as described in **Table 1** and **Table 2**, **Armezon** controls or suppresses the following weeds.

- **Common Bermudagrass** (*Cynodon dactylon*) - Apply early in the season at onset of Bermudagrass greenup or emergence of new leaves. Apply 1.0 to 2.0 fl ozs/A of **Armezon** per application. Up to four sequential applications at 2 to 3 week intervals may be necessary for best control. **DO NOT** apply more than 4.0 fl ozs/A of **Armezon** (0.0875 lb topramezone per acre) per year. Apply using MSO or COC spray adjuvant plus nitrogen fertilizer such as AMS or UAN. See **Adjuvants** and **Nitrogen Fertilizer** sections for details.
- **Fall panicum** (*Panicum dichotomiflorum*) - For rescue suppression of large fall panicum more than 12-inches tall or other annual grasses listed in **Table 1** and **Table 2**, apply 2.0 fl ozs/A **Armezon** and use a minimum of 20 gallons per acre spray volume for proper spray coverage. Apply using MSO or COC spray adjuvant plus nitrogen fertilizer such as AMS or UAN. See **Adjuvants** and **Nitrogen Fertilizer** sections for details.

Sugarcane Restrictions

- **DO NOT** apply more than 2.0 fl ozs/A of **Armezon** (0.0438 lb topramezone/A) per application in sugarcane.
- **DO NOT** apply more than 4.0 fl ozs/A of **Armezon** (0.0875 lb topramezone/A) per year in sugarcane.
- **DO NOT** exceed a seasonal total of 2.0 fl ozs/A of **Armezon** during the final year of sugarcane production, prior to rotation to another crop.
- **DO NOT** apply **Armezon** within 100 days of sugarcane harvest.
- **DO NOT** graze or feed treated sugarcane for at least 100 days following an application of **Armezon**.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Armezon, Clearfield, Guardsman Max, Outlook, Prowl, and Status are registered trademarks of BASF.

G-Max Lite is a trademark of BASF.

Balance, Laudis, and LibertyLink are registered trademarks of Bayer.

Callisto is a registered trademark of a Syngenta Group Company.

Roundup and Roundup Ready are registered trademarks of Monsanto Technology LLC.

U.S. Patent No. 8,017,556

© 2022 BASF Corporation
All rights reserved.

007969-00262.20220823.NVA 2022-04-0216-0159
Supersedes: NVA 2015-04-216-0157

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


We create chemistry