



- When applying to vegetation, such as bushes and trees, spray to the point of run-off.
- To kill ants, apply directly to trails, around doors, windows and other places where ants may be found.

**For Outdoor Treatment of Carpenter Ants**

- To kill carpenter ants, apply as a perimeter treatment as described above.
- Locate and treat carpenter ant nests for best results. Also apply to trails around doors and windows and other places where carpenter ants have been observed or are expected to forage.
- **DO NOT directly treat firewood with this product. To protect firewood from carpenter ants (and termites), spray the soil beneath where the firewood will be stacked. 32 oz. will treat an area of up to 640 sq.ft. Not intended to replace structural treatments. Not recommended as the sole protection against termites. Applications of Bug-No-More Large Property Insect Control are not a substitute for mechanical alterations, soil or foundation treatment but merely as a supplement. For active infestations, seek professional inspection and treatment.**

**For Outdoor Treatment of Bees and Wasps**

- To kill bees and wasps, thoroughly spray 1 fl.oz. per gallon of water onto nest entrance and any surrounding areas where these pests are found.
- Apply late in the evening when insects are at rest.

**For Outdoor Treatment of Blacklegged and Brown Dog Ticks**

- To kill blacklegged ticks and brown dog ticks and to prevent infestation of buildings, apply 15.0 fl.oz. per gallon of water per 1000 sq.ft. to soil and vegetation around the exterior perimeter of the building as described in the instructions for use in perimeter treatment.
- Spray to the point of runoff, applying spray thoroughly and uniformly on treated areas.
- Also treat areas of taller vegetation (2 – 3 ft.) around the exterior of the property and soil and vegetation which border wooded areas of the property where blacklegged and brown dog ticks may be harboring.

**For Outdoor Treatment of Spiders, Scorpions, Mole Crickets, Crickets and Grasshoppers**

- To kill spiders, scorpions, spiders, mole crickets, crickets and grasshoppers and to prevent infestation of buildings, apply 15 fl.oz. per gallon of water per 1,000 sq.ft. to the soil and vegetation around the exterior perimeter of the building as described in the instructions for use in perimeter treatment.
- Spray to the point of runoff, applying spray thoroughly and uniformly on treated areas.

**ORNAMENTALS, INCLUDING FLOWERS, ROSES, SHRUBS and TREES**

**For treatment of listed pests in annual and perennial flowers, roses, bedding plants, ground covers, shrubs and deciduous and evergreen trees**

- Spray 15 fl.oz. per gallon of water per 1000 sq.ft. up to the point of run-off making sure to cover all sides of leaves and tree trunks.
- For ground covers, bedding plants, foliage, annuals and perennials, apply 15 fl.oz. per gallon of water as a uniform spray. 32 fl.oz. will treat 2133 sq.ft. Repeat treatment as necessary to achieve control using higher application rates as pest pressure & foliage area increases.

Plant Safety Note: Certain plants may be sensitive to the final spray solution. A limited area should be treated and observed for one week prior to application to the entire planting.

**HOME GARDENS:**

**To kill insects on fruit, nut and vegetable plants**

- 1). Use 5 fl.oz per 1 gallon of water to cover 1000 sq.ft.
- 2). Spray leaves to point of dripping.
- 3). Reapply if insects return, but should not be more than once every seven days and according to harvest directions.
- 4). Follow chart below for list of plants, pests and harvesting directions.

Plants	Pests	Use Before Harvest
<b>VEGETABLES</b>		
<b>Leafy Vegetables</b> Amaranth (leafy amaranth, Chinese spinach, tampala); Arugula; Cardoon; Cilantro; Corn salad; Garden cress; Upland cress; Dandelion; Dock; Endive; Florence Fennel; Head Lettuce; Leaf Lettuce; Orach; Parsley; Garden Purslane; Winter Purslane; Radicchio (red chicory); Rhubarb; Spinach; Swiss chard	Corn Earworm/Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetles  Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Tobacco Budworm, Armyworms, Loopers, Lygus Bugs, Onion Thrips, Stink Bugs, Wireworm	<b>Spray up to 1 day before harvesting.</b>
<b>Fruiting Vegetables including:</b> Eggplant; groundcherry (Physalis spp.); pepino (Melon pear); pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); tomatillo; tomato	Armyworm, Southern  Armyworm, True Armyworm, Yellow-striped Celery Leaf Tier, Colorado Potato Beetle, Corn Borer, European, Corn Borer, Southwestern  Corn Earworm, Cucumber Beetle, Cutworm, Flea Beetle, Garden Webworm, Green Stink Bug, Hornworms, Leafminers (adults), Leafhoppers, Meadow Spittlebug, Pepper Maggot (adults), Pepper Weevil, Plant Bugs, Tobacco Budworm, Tomato Fruitworm, Tomato Pinworm, Aphids, Armyworm, Beet Armyworm, Fall Cabbage Looper, Grasshoppers, Lygus Bugs, Brown Stink Bug, Tomato Psyllid, Thrips, Whiteflies.	<b>Spray up to 1 day before harvesting.</b>
<b>Brassica Vegetables including:</b> Broccoli; Chinese Broccoli (gai lan, white flowering broccoli); Brussels Sprouts; Cauliflower; Cavalo broccolo; Kohlrabi; Cabbage; Chinese Cabbage (napa); Leafy Brassica Greens including: Broccoli Raab (rapini); Chinese cabbage (bok choy); Chinese Mustard Cabbage (gai choy); Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens	Corn Earworm, Cucumber Beetles, Cutworm, Diamondback Moth, Flea Beetles, Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Southern Cabbageworm, Tobacco Budworm Alfalfa Looper, Armyworms, Cabbage Looper, Cabbage Webworm, Ground Beetles, Leafminers (adults), Lygus Bugs, Onion Thrips, Stinkbugs, Wireworm (adults), Aphids, and Whiteflies	<b>Spray up to 1 day before harvesting.</b>
<b>Root &amp; Tuber Vegetables</b> Arracacha; Arrowroot; Artichoke; Garden Beet; Edible Burdock; Edible Canna; Carrot; Turnip-rooted chervil; Chicory; Chufa; Dasheen (Taro); Ginger; Ginseng; Horseradish; Leren; Turnip-Rooted Parsley; Parsnip; Potato; Oriental Radish (Daikon); Rutabaga; Salsify (Oyster Plant); Black Salsify; Spanish Salsify; Skirret; Sweet Potato; Tanier (Cocoyam); Turmeric; Turnip; Yam Bean; Yan (True)	Cutworms, Cabbage Looper  Cucumber Beetle, European Corn Borer, Fleabeetles, Leafhoppers, Southern Corn Rootworm (adult), Vegetable Weevil, Whitefringed Beetle (adult), Armyworm, Beet Armyworm, Yellowstriped Cabbage Maggot, Colorado Potato Beetle, Grasshoppers, Imported Cabbageworm, Potato Leafhopper/Potato Psyllid, Potato Tuberworm, Tarnished Plant Bug	<b>Spray up to 1 day before harvesting.</b>

(cont. on next page)



PESTS KILLED	
Caterpillars, Butterflies and Moths	Tent forming caterpillars, cocoon forming caterpillars, chrysalis forming caterpillars. Leaf feeding caterpillars (including leaves of flowering plants, bushes and other ornamentals, vegetables and trees), fruit eating caterpillars, legume feeding caterpillars, vegetable feeding caterpillars, stem and trunk boring caterpillars, and leafrolling caterpillars. Caterpillars of the Yucca Moth family; Clothes moth family; bagworm moth family, Gelechiid moth family; Ermine Moth family; Clear Winged moth family; Tortricid Moth family (millers) such as Codling moth; Swallowtails; Whites, Sulphurs and Orange Tips family; Gossamer-Winged Butterflies Family; Metalmark family; Snout Butterflies family; Brush-Footed Butterfly family; Hackberry and Goatweed Butterfly family; Satyrs, Nymphs and Arctics family; Milkweed Butterfly family; Smoky Moth family; Pyralid Moths family such as Sod Webworm Moth; Plume Moth family; Measuringworm Moth family; Tiger Moth family; Ctenuchid Moth family; Tussock moth family; Owlet Moth family such as Eight-spotted Forester
Common Bristletails	Silverfish Family
Springtails	Podurid Springtail family such as Snow Pea, Seashore Springtail; Entomobryid Springtail family such as Ainsley's Springtail; Globular Springtail family
Cockroaches	Blattid Cockroach Family such as Oriental Cockroach (Asiatic Cockroach, Shad Cockroach), and American Cockroach (Waterbug); Blattellid Cockroaches family such as German Cockroach (Croton Bug)
Termites	Rotting-wood termite family such as Pacific Coast Termite; Subterranean termite family such as subterranean termites. Not intended to replace structural treatments. For termites – not recommended as sole protection against termites. Such applications are not a substitute for mechanical alterations, soil or foundation treatment but merely as a supplement. For active infestations, get a professional inspection.
Earwigs	Long-Horned Earwig family such as Ring-legged Earwig, and Riparian Earwig; Common Earwig family such as European Earwig
Grasshoppers & Crickets	Pygmy Grasshopper family; Long-horned grasshoppers and Katydid family; Camel Crickets family; True Crickets family such as House Cricket
True Bugs	Plant Bug family; Assassin bug family; Lace Bug family; Seed Bug family; Stilt Bug family; Leaf-footed Bug family; Scentless Plant Bug family such as Eastern Boxelder Bug and Western Boxelder Bug; Stink Bug family
Cicadas and Kin	Cicada family such as Periodical Cicadas, Dogday Harvestfly, and Grand Western Cicada
Treehoppers	Treehopper family, such as Thorn-mimic Treehopper, Oak Treehopper, Buffalo Treehopper, and Locust Treehopper
Leafhoppers	Leafhopper family such as Grape Leafhopper, Scarlet-and-green Leafhopper, Sharpshooter
Thrips	Banded Thrips family such as Banded-wing Thrips; Common Thrips family
Beetles	Tiger Beetle Family; Ground Beetle family such as Bombardier Beetle; Predacious Diving Beetle family; Whirligig Beetle family; Hister Beetle family; Rove Beetle family; Carrion Beetle family; Stag Beetle; Bessbugs family; Scarab Beetle family; Click Beetle family; Soldier Beetle family, Net-winged Beetle family; Spider Beetle family; Branch and Twig Borer family; Checkered Beetle family; Flat Bark Beetle family; Ladybug Beetles; Darkling Beetles; Fire-colored Beetles family; Blister Beetles family; Long-horned Beetle family; Seed Borer family; Leaf Beetle family; Primitive Weevil family; Snout Beetles and Weevil family; Bark and Ambrosia Beetle family
Bees, California Carpenter Bees, Ants, Wasps, Sawflies	Cimicid Sawflies family; Common Sawfly family; Horntail family; Stem Sawflies family; Braconids family; Ichneumons family; Torymid Wasp family; Chalcid family; Gall Wasp family; Pelecinid family; Tiphid Wasp family; Velvet-Ants; Scoliid Wasp family; Ant family; Vespid Wasp family; Spider Wasps; Sphecid Wasp family; Bee families
Flies	Crane fly family; Phantom Crane Flies; Net-winged Midges; Black Flies; March Flies family; Fungus Gnats family; Snipe Fly family; Stiletto Fly family; Flower-loving Fly family; Mydas Fly family; Robber Fly family; Bee Flies; Dance Flies; Long-Legged Flies; Hover Fly family; Thick Headed Fly family; Pyrgotid Fly family; Fruit Fly family; Seaweed Fly family; Marsh Fly family; Shore fly family; Pomace fly family; Anthomyiid Fly family; Muscid Fly family; Louse Fly family, Blow Fly family; Flesh Fly family; Tachinid Fly family
Spiders	Folding trapdoor spider family; Tarantula family; Trapdoor Spider family; Spitting spider family; Violin Spider family; Comb-footed Spider family; Sheet-web Weaver family; Orb Weaver family; Large-jawed Orb Weaver family; Funnel Web Weaver family; Nursery Web Spider family; Wolf Spider family; Lynx Spider family; Sac Spider family; Wandering spider family; Giant Crab spider family; Selenopid Crab Spider family; Crab Spider family; Philodromid family Daddy-long-legs family
Ticks	Blacklegged tick, Brown dog tick, Deer tick
Psyllid, Whiteflies, Aphids, Mealybugs, Scale Insects	Psyllid family; Whitefly family; Aphid family; Woolly and Gall-Making Aphid family; Giant Scale Insects family; Armored Scale Insects family; Mealybug family; Cochineal Bug family
Millipedes	Millipedes

IMPORTANT – Directions for Storage and Disposal	
<b>Pesticide Storage:</b>	Protect from freezing. Store this product in its original container in a cool, dry (preferably locked) area that is inaccessible to children and animals. After partial use, replace original lid and close tightly.
<b>Container Disposal:</b>	Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling, if available. If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY

DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

724/8-2019 AP081414  
EPA REG. NO. 279-3472-2217



**An Employee-Owned Company**  
**MANUFACTURED BY**  
**PBI/GORDON CORPORATION**  
 P.O. BOX 860350  
 SHAWNEE, KANSAS 66286  
 www.GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.