

Kills On Contact Controls Listed Pests For Up To 4 Weeks

ACTIVE INGREDIENT:

Permethrin [*(3-phenoxyphenyl)methyl (±) cis-trans-3-(2, 2-dichloroethenyl)-2, 2-dimethylcyclopropanecarboxylate].... 0.25%

*cis-trans isomer ratio: min. 35%(±) cis, max. 65% (±) trans.

KEEP OUT OF REACH OF CHILDREN CAUTION



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid inhalation of dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Avoid contamination of food and feed

FIRST AID				
Have the product container or label when calling a poison control center or doctor, or going for treatment.				
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.			
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.			
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.			

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

For outdoor use around the home only.

Gordon's Bug-No-More® Lawn Insect Granules contains permethrin, a long lasting insecticide which provides contact and residual control for up to 4 weeks of troublesome outdoor insect pests. Controls grubs and other listed root feeding

Three Easy Steps to Use this Product

Determine size of the job, measure the area to be treated.

Use spreader setting chart to determine appropriate setting for your

spreader.

Step #3: Treat as directed in rate chart. Thorough coverage is important. Water

in granules where indicated.

Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

Note: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary, in order to assure correct rate of application.

SPREADER SETTINGS

Spreader Make and Model	Grass or Dichondra Lawns 2 pounds Per 1,000 Sq. Ft.	St. Augustine Lawn 3 pounds Per 1,000 Sq. Ft.	White Grubs, Billbugs 2½ pounds Per 1,000 Sq. Ft.
Ames	12	_	_
Bigfoot	F	G	F-G
Central Imperial Make No. 1662-G	6½	71/4	7
Central Suburbia Lawn Spreader No. 1622-O	6½	7½	7
Cyclone Model B1 Rotary	41/2	5	43/4
Cyclone Pro	3.25	3.5	3.4
Cyclone SB3000	6	_	_
Earthway	12	_	_
Earthway 2400 Rotary	14	15	14
Earthway Model 73008	12	_	_
Greenfield Model 105	6½	71/4	6¾
Lely Broadcast	2.3	2.5	2.4
Lesco Rotary	10(E)	11(F)	10(E)
Ortho Drop Spreader Model 3000	41/2	5½	5
Ortho Drop Broadcast Spreader Model 4000	41/2	7	6
Precision Red Devil	1	_	_
Prizelawn CB Rotary	3.8	4.2	4.0
Scotts Accurgreen AG-3	2	_	_
Scotts EasyGreen EG-3	26	_	_
Scotts EvenGreen EV-1	6	_	_
Scotts Model 35	4	5	4½
Scotts PrecisionGreen PF-4	5	_	_
Scotts R8A	F	G	G
Scotts SpeedyGreen SG-3	5	_	_
Sears Craftsman Model 671.19198	6	7	6½
Spyker 24-22 Rotary	3.6	3.9	3.75

DIRECTIONS FOR APPLYING GORDON'S BUG-NO-MORE® LAWN INSECT GRANULES

Treatment Areas	Target Pest	Remarks	Use Rate
Band Treatment Around House Foundation	Ants, Fleas, Brown dog ticks, Millipedes, Clover mites, Roaches, Spiders, Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs	These pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses. Apply a 5-foot band of granules uniformly. Water lightly after application	3.2 oz. for each 100 sq. ft. area (5' by 20')
Overall Lawn Application for Grass and Dichondra Lawns	Ants, Armyworms, Brown dog ticks, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs	Apply with a fertilizer spreader. See spreader setting chart above. Apply when grass is dry. Water lawn lightly after application.	For application with other types of equipment, apply at the rate of 2 lbs. per 1000 sq. ft., (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10'). For heavy lawns such as St. Augustinegrass, the rate should be increased to 3 lbs. per 1,000 sq. ft. (or 4.8 oz. per 100 sq. ft.)
	Grubs, Japanese beetle grubs, Ateanius spretulus (dung beetles)	Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with ½ - ½ inches of water after application. Treat again in late summer or early fall if a second generation occurs.	Apply 2.5 lbs. per 1000 sq. ft. of lawn.
Spot Treatment	Ants	Water lightly after application.	Apply 1½ teaspoons over and around each ant hill.
	Fire Ants	Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. Note: For best results, apply in cool weather, 65° - 80°F, or in early morning or late evening hours. Treat new mounds as they appear.	Apply 1 cup over and 2 ft. around each fire ant mound.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container in a dry place. Store in locked area out of reach of children.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes can not be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Offer for recycling, if available. If recycling is not available, then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Read the entire Directions for Use and Warranty before using this Product. By using this product, user or buyer accepts the following warranty.

The manufacturer is committed to providing high quality products. To the extent consistent with applicable law, the manufacturer makes no warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. To the extent consistent with applicable law, the manufacturer or the Seller shall not be liable for indirect, special, incidental or consequential damages. To the extent consistent with applicable law, the exclusive remedy of the user or buyer shall not exceed the purchase

604/11-2014 AP090110 EPA REG. NO. 53883-39-2217



An Employee-Owned Company

MANUFACTURED FOR **PBI/GORDON CORPORATION** 1217 WEST 12TH STREET KANSAS CITY, MISSOURI 64101 www.pbigordon.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.