

CIDETRAK® CM

A MATING DISRUPTANT FORMULATION FOR:

CODLING MOTH (*Cydia pomonella*)

AND HICKORY SHUCKWORM (*Cydia caryana*)

ACTIVE INGREDIENT:

(8E,10E)-8,10-dodecadien-1-ol2.00 %

OTHER INGREDIENTS:98.00 %

TOTAL:100.00%

NET CONTENTS:

Dispensers per package: 400 units

One dispenser contains: 120 mg, active ingredient

EPA Registration No. 51934-10

EPA Establishment No. 51934-OK-002

MANUFACTURED BY: TRÉCÉ INCORPORATED, 7569 Hwy 28 W · Adair, Oklahoma 74330

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wear chemical resistant gloves and protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID STATEMENT

IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice.
HOT LINE NUMBER:	<ul style="list-style-type: none"> Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-866-785-1313 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water when disposing of product, wash water, or when cleaning equipment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING

CIDETRAK® CM disrupts the mating communication between adult male and female moths, reduces fertile egg laying, suppresses subsequent larval infestation, and reduces the need for applying insecticides. For maximum effectiveness, growers should remove and destroy infested fruit from trees and the ground. Growers should maintain accurate records of CM adult population monitoring and larval damage within their orchards. Consult your local extension specialist, certified crop advisor, or Trece representative for assistance in designing the best CIDETRAK® CM program for your orchard.

Crop	Apple, Pear, Walnut, Pecan, Quince, and other pome fruits
Targeted Pest	Codling Moth (<i>Cydia pomonella</i>), Hickory Shuckworm (<i>Cydia caryana</i>).
Rate	Minimum 320 dispensers per acre (800 dispensers per hectare) or 38 gm a.i. per acre per application. Maximum of 800 dispensers per acre (2,000 dispensers per hectare) or 96 gm a.i. per acre per application. Apply 400 dispensers per acre (1000 dispensers per hectare) or 48 gm a.i. per acre per application to moderately infested orchards. Apply 10-20% of the total per acre rate to all orchard borders, to orchards that are moderately heavily infested with CM, or adjacent to an untreated orchard. Do not exceed 150 grams of active ingredient per acre per year.
Application	Attach dispensers securely to lateral branches in upper one-third of tree canopy. Attach the dispensers from the ground using a pole applicator or from a moving trailer.
Timing	Apply prior to moth emergence in early spring. Monitor moth activity using PHEROCON® IIB or PHEROCON VI traps and lures. Begin monitoring in early spring and continue throughout the season to assess treatment effectiveness. Depending on temperature, CIDETRAK® CM dispensers will release pheromone for 120-140 days. In areas with long field season (i.e. more than 120 days) a second application may be necessary, based on monitoring results, prior to subsequent CM flights. NOTE: CIDETRAK CM will not prevent crop damage from immigration of mated female moths into treated orchards. Nearby (500 yards) untreated Codling Moth hosts, such as apple, walnut, pear, quince, crabapple, plum and others, can be a source of these mated females. Treatment with pheromone is recommended only when all host crops within or near treated blocks have been treated with CIDETRAK® CM.
Treatment Tips	CIDETRAK® CM suppresses mating of Codling Moth and Hickory Shuckworm. Immigration of mated female moths from adjacent, infested orchards can reduce the level of control. Area-wide treatment of the entire host plant block is the most effective strategy. To supplement CIDETRAK CM, use insecticides to control high populations. Monitor all pest populations to determine timely use of insecticides. To manage immigration and high population pressures, consider: <ol style="list-style-type: none"> 1. Treatment of external sources of infestation with CIDETRAK CM 2. Treatment of external sources of infestation with an insecticide 3. Treatment of pheromone treated orchard with insecticide Consult your local extension specialist, certified crop advisor, or Trece representative for local field condition management strategies.

STORAGE AND DISPOSAL

Do not contaminate food or feed by storage or disposal

PESTICIDE STORAGE: Store unopened original package in a dry location at a temperature below 40° F.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Dispose of empty containers and foil envelopes in trash. Offer for recycling.

WARRANTY AND LIMITATION OF DAMAGES

To the extent consistent with applicable law, all statements concerning the use of this product apply only when used as directed as stated on this product label. Trece Inc. warrants that the product conforms to this label and under normal conditions of use according to this label, is reasonably fit for the stated label purpose. The manufacturer makes no express or implied warranties of merchantability or fitness for a purpose concerning this product, or its use, extending beyond the label. Under no circumstance will Trece Inc. be liable for damage (special, consequential or incidental) that result from the handling, storage, or use of this product which is not in compliance with the label. Read all label directions carefully.

Buyer's exclusive remedy, should this product be defective, is replacement of the product, or if this is impractical, a purchase price refund.



Mailing Address: P.O. Box 129 · Adair, Oklahoma 74330
Shipping Address: 7569 Hwy 28 W · Adair, Oklahoma 74330
Phone: (918) 785-3061 · Fax: (918) 785-3063

PHEROCON® and CIDETRAK® are registered Trademarks of TRECE Incorporated, Adair, OK USA
©2016 TRECE, Adair, OK USA · TRE0879