

CIDETRAK[®] CMDA COMBO[™] PP

A MATING DISRUPTION FORMULATION FOR:

CODLING MOTH (*Cydia pomonella*)

AND HICKORY SHUCKWORM (*Cydia caryana*)

ACTIVE INGREDIENTS:	Percent by Wt.
8,10-Dodecadien-1-ol, (8E, 10E).....	1.80%
2,4-decadienoic acid, ethyl ester, (E, Z)	1.20%
OTHER INGREDIENTS:	97.00%
TOTAL:	100.00%

EPA Registration No. 51934-13 EPA Establishment No. 51934-OK-002

NET CONTENTS:

Dispensers per package: 400

One dispenser contains: 90 mg 8,10-Dodecadien-1-ol, (8E, 10E)

60 mg 2,4-decadienoic acid, ethyl ester, (E, Z)

MANUFACTURED BY: TRÉCÉ INCORPORATED, 7569 Hwy 28 W · Adair, Oklahoma 74330

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

FIRST AID STATEMENT

IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.
HOT LINE NUMBER:	<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p> <p>For emergency information on product, use, etc., call the National Pesticides Information Center at 1-800-858-7378 6:30 AM to 4:30 PM Pacific Time (PT), seven days a week. During other times, call the poison control center at 1-800-222-1222. For additional information, you may also contact Trece Inc. at 1-866-785-1313.</p>

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING

CIDETRAK[®] CMDA COMBO[™] PP disrupts the mating communication between adult male and female moths, reduces fertile egg laying, suppresses subsequent larval infestation, and reduces the need for applying insecticides.

For maximum effectiveness, growers should remove and destroy infested fruit from trees and the ground. Growers should maintain accurate records of CM adult population monitoring and larval damage within their orchards. Consult your local extension specialist, certified crop advisor, or Trece representative for assistance in designing the best CIDETRAK[®] CMDA COMBO[™] PP program for your orchard.

Crop	Apple, Pear, Walnut, Pecan, Quince, and other pome fruits
Targeted Pest	Codling Moth (<i>Cydia pomonella</i>), Hickory Shuckworm (<i>Cydia caryana</i>).
Rate	CIDETRAK[®] CMDA COMBO[™] PP: Minimum of 200 dispensers per acre (500 dispensers per hectare) or 18 gm CM a.i. and 12 gm DA a.i. per acre per application. Maximum of 400 dispensers per acre (1,000 dispensers per hectare) or 36 gm a.i. and 24 gm DA a.i. per acre per application. Apply 320 dispensers per acre (800 dispensers per hectare) for moderate populations. Apply the maximum dispenser rate to all orchard borders, to orchards that are heavily infested with CM, or adjacent to an untreated orchard. Do not exceed 150 gm combined a.i. per acre per year.
Application	Pome Fruit - Attach dispensers securely to lateral branches at 5-10% of the distance between the trunk and branch terminal in the upper 1/5th of the canopy. Attach the dispensers from the ground using a pole applicator or from a moving trailer.
Timing	Apply prior to moth emergence in early spring. Monitor moth activity using PHEROCON [®] IIB or PHEROCON VI traps and PHEROCON CMDA COMBO + AA lures. Begin monitoring in early spring and continue throughout the season to assess treatment effectiveness. Note - CIDETRAK[®] CMDA COMBO[™] PP will <u>not prevent crop damage</u> from immigration of mated female moths into treated orchards. Nearby (500 yards) untreated Codling Moth hosts, such as apple, walnut, pear, quince, crabapple, plum and others, can be a source of these mated females. Treatment with pheromone is recommended only when all host crops within or near treated blocks have been treated with CIDETRAK CMDA COMBO PP .
Treatment Tips	CIDETRAK[®] CMDA COMBO[™] PP suppresses mating of Codling Moth. Immigration of mated female moths from adjacent, infested orchards can reduce the level of control. Area-wide treatment of the entire host plant block is the most effective strategy. To supplement CIDETRAK CMDA COMBO PP , use insecticides to control high populations. Monitor all pest populations to determine timely use of insecticides. To manage immigration and high population pressures, consider: 1. Treatment of external sources of infestation with CIDETRAK CMDA COMBO PP 2. Treatment of external sources of infestation with an insecticide 3. Treatment of pheromone treated orchard with insecticide Consult your local extension specialist, certified crop advisor, or Trece representative for local field condition management strategies. Monitoring Tip - PHEROCON CMDA COMBO [™] + AA lures capture significantly more adults than PHEROCON CMDA COMBO [™] lures in DA treated orchards.
STORAGE AND DISPOSAL	
Do not contaminate food or feed by storage or disposal of unused or used dispensers.	
PESTICIDE STORAGE: Store unopened original package in a dry location at temperature below 40°F. To avoid contamination of food or feed items, only unopened product packages may be stored in cold storage facilities that are used for food storage.	
PESTICIDE DISPOSAL: Product wastes resulting from use of this product may be disposed of on use site or at an approved waste disposal facility.	
CONTAINER DISPOSAL: Offer for recycling, if available. Dispose of empty dispensers and foil envelopes in the trash.	

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed as stated on this product label. To the extent consistent with applicable law, Trece Inc. warrants that the product conforms to this label and under normal conditions of use according to this label, is reasonably fit for the stated label purpose. To the extent consistent with applicable law, the manufacturer makes no express or implied warranties of merchantability or fitness for a purpose concerning this product, or its use, extending beyond the label. Under no circumstance will Trece Inc. be liable for damage (special, consequential or incidental) that result from the handling, storage, or use of this product which is not in compliance with the label. Read all label directions carefully.

Buyer's exclusive remedy, should this product be defective, is replacement of the product, or if this is impracticable, a purchase price refund.



Mailing Address: P.O. Box 129 · Adair, Oklahoma 74330
Shipping Address: 7569 Hwy 28 W · Adair, Oklahoma 74330
Phone: (918) 785-3061 · Fax: (918) 785-3063

PHEROCON[®] and CIDETRAK[®] are registered Trademarks of TRECE Incorporated, Adair, OK USA
©2016 TRECE, Adair, OK USA · TRE0879