

# CLEANSE™ 2 EC

Herbicide

**ACTIVE INGREDIENT:**

\*Clethodim .....26.4%

**OTHER INGREDIENTS:** .....73.6%

**TOTAL:** .....100.0%

Contains Petroleum Distillates

\*(E)-2-[1-[[[3-chloro-2-propenyl]oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one  
Cleanse 2 EC Contains 2.0 lbs. clethodim per gal.

**BY WT.**

## KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for additional Precautionary Statements, First Aid, and complete Directions for Use.

EPA Reg. No. 83222-30

EPA Est. No. 37429-GA-001  
37429-GA-002  
42750-MO-001

Made in U.S.A.

**NET CONTENTS: 1 GALLON**



**DIRECT AG  
SOURCE™**

Manufactured For:  
Direct Ag Source, LLC  
30473 260th Street • Eldora, IA 50627



CLEANSE-N010515-REVA012915

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS & DOMESTIC ANIMALS**  
**WARNING**

Causes substantial but temporary eye irritation. Avoid contact with skin. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid breathing vapors or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

<b>FIRST AID</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give any liquid to the person.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 7:30 AM-3:30 PM PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information.	
<b>NOTE TO PHYSICIANS</b>	
Ingestion of this product or subsequent vomiting can result in aspiration of light hydrocarbon liquid, which can cause pneumonitis. If ingested, probable mucosal damage may contraindicate the use of gastric lavage.	

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- long-sleeved shirt and long pants,
- chemical-resistant gloves such as Barrier Laminate or Viton  $\geq$  14 mils,
- shoes plus socks, and
- protective eyewear.

**Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwater or rinsate.

The use of this product may pose a hazard to the federally designated endangered species of Solano Grass and Wild Rice. Use of this product is prohibited in the following areas where the species are known to exist:

Solano Grass: Solano County, California: the vernal lakes area bounded by the Union Pacific Railroad and Hastings Road to the north, Highway 113 to the east, Highway 12 to the south, and Travis Air Force Base to the west.

Wild Rice: Hays County, Texas.

### **PHYSICAL OR CHEMICAL HAZARDS**

Combustible. Do not use or store near heat or open flame.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL AND PAMPHLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls,
- Chemical-resistant gloves, such as Barrier Laminate or Viton  $\geq$  14 mils,
- Shoes plus socks,
- Protective eyewear.

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift.

Do not enter treated areas without protective clothing until sprays have dried.

## TANK MIXES

**Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.**

Read and follow the entire label of each product to be used in the tank mix with this product.

## CHEMIGATION

May be applied to onions and garlic by sprinkler irrigation systems. Do not apply by chemigation to any other crop, or to this crop using any other type of irrigation system.

## PRODUCT INFORMATION

**FOR USE ON:** Alfalfa, Artichoke (Globe), Asparagus, Bean and Pea (dry shelled)<sup>1</sup>, Bean and Pea (succulent shelled)<sup>2</sup>, Broccoli, Bushberry<sup>3</sup>, Cabbage, Caneberry<sup>4</sup>, Canola\*, Carrot, Cauliflower (and other Head and Stem Brassica Vegetables)<sup>5</sup>, Celery, Clover (grown in Idaho, Oregon and Washington only), Conifers, Cotton, Cranberry, Cucumber, Eggplant (and other Fruiting Vegetables)<sup>6</sup>, Fallow Land (and other non-producing agricultural areas), Flax\*, Field Corn<sup>7</sup>, Garden Beet, Garlic, Herbs<sup>7</sup>, Hops, Horseradish (and other Root Vegetables)<sup>8</sup>, Legume Vegetables (edible podded)<sup>9</sup>, Lettuce, Head and Leaf (and other leafy greens)<sup>10</sup>, Melons (including Cantaloupes and Watermelons)<sup>11</sup>, Mint, Mustard Greens (and other leafy brassica greens)<sup>12</sup>, Mustard Seed\*, Non-Bearing Food Crops, Non-Crop or Non-Planted Areas, Onions (dry bulb and green), Ornamentals, Peaches, Peanuts (including perennial), Peppers (bell and non-bell), Potato, Radish, Rhubarb (and other Leaf Petioles)<sup>13</sup>, Safflower, Sesame, Shallots (dry bulbs and green), Soybeans, Squash (including Pumpkins), Strawberry, Sugar Beet, Sunflower, Sweet Potato, Tomato and Yam (and other Tuberous and Corn Vegetables)<sup>14</sup>.

**\*Not for use in California**

- <sup>1</sup> Other Dry Shelled Bean and Pea crops approved for use with CLEANSE 2 EC include: Bean (*Lupinus* spp.), grain, sweet, white and white sweet; Bean (*Phaseolus* spp.), field, kidney, lima (dry), navy, pinto and tepary; Bean (*Vigna* spp.), adzuki bean, black-eyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, broad (dry), chickpea (garbanzo), guar, lablab bean and lentil; Pea (*Pisum* spp.), field and pigeon.
- <sup>2</sup> Other Succulent Bean and Pea crops approved for use with CLEANSE 2 EC includes: Bean (*Lupinus* spp.), grain, sweet, white and white sweet; Bean (*Phaseolus* spp.), field, kidney, lima (dry), navy, pinto and tepary; Bean (*Vigna* spp.), adzuki bean, black-eyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, broad (dry), chickpea (garbanzo), guar, lablab bean and lentil; Pea (*Pisum* spp.), field and pigeon.
- <sup>3</sup> Other bushberry crops approved for use with CLEANSE 2 EC include: Aronia Berry, Blueberry (highbush), Black Currant, Buffalo Currant, Chilean Guava, Cranberry (highbush), Elderberry, European Barberry, Gooseberry, Honeysuckle (edible), Huckleberry (edible), Jostaberry, Juneberry (Saskatoon berry), Native Currant, Salal, Sea Buckthorn, Red Currant; cultivars, varieties and/or hybrids of these.
- <sup>4</sup> Other Caneberry crops approved for use with CLEANSE 2 EC include: Blackberry, Loganberry, Raspberry (black and red), Wild Raspberry; cultivars, varieties and/or hybrids of these.
- <sup>5</sup> Other head and stem brassica vegetables approved for use with CLEANSE 2 EC include: Chinese broccoli, Brussels sprouts, Chinese (napa) cabbage, Chinese mustard, cavalo broccoli and kohlrabi.
- <sup>6</sup> Other Fruiting Vegetables (except tomato) approved for use with CLEANSE 2 EC include: eggplant, groundcherry, pepino, peppers (all) and tomatillo.
- <sup>7</sup> For burndown of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn.
- <sup>8</sup> Other Herb crops approved for use with CLEANSE 2 EC include: angelica, balm, basil, borage, burnet, camomile, catnip, chervil (dried), chive, Chinese chive, clary, coriander (leaf), costmary, cilantro (leaf), curry (leaf), dill (dillweed), horehound, hyssop, lavender, lovage (leaf), marigold, marjoram (*origanum* spp.), nasturtium, parsley (dried), pennyroyal, rosemary, rue, sage and savory, summer and winter.
- <sup>9</sup> Other root vegetables approved for use with CLEANSE 2 EC include: burdock, edible; celeriac; chervil, turnip-rooted; chicory; ginseng; parsley, turnip-rooted; parsnip; radish, oriental; rutabaga; salsify; salsify, black; salsify, Spanish; skirret and turnip.
- <sup>10</sup> Other Edible Podded Legume Vegetable crops approved for use with CLEANSE 2 EC include: Bean (*Phaseolus* spp.), runner, snap and wax; Bean (*Vigna* spp.), asparagus, Chinese longbean, moth, yardlong, jackbean; Pea (*Pisum* spp.), dwarf, edible-pod, snow, sugar snap, pigeon and sword bean.
- <sup>11</sup> Other Leafy Greens crops approved for use with CLEANSE 2 EC include: amaranth (Chinese spinach, leafy amaranth and tampala), arugula (rocket), chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden, yellow rock and winter), dandelion, dock (sorrel), endive (escarole), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio (red chicory), spinach (New Zealand and Vine (Indian and malabar)).
- <sup>12</sup> Other cucurbit crops approved for use with CLEANSE 2 EC include: Chayote (fruit), Chinese Wax Gourd, Citron Melon, Edible Gourd, Gherkin and Muskmelons (all) including Honeydew Melon.
- <sup>13</sup> Other leafy brassica greens approved for use with CLEANSE 2 EC include: broccoli raab, Chinese (bok choy) cabbage, collards, kale, mizuna, mustard greens, mustard spinach, rape greens and turnip greens.
- <sup>14</sup> Other leaf petiole crops approved for use with CLEANSE 2 EC include: cardoon, celtuce, Chinese celery, Florence fennel, and Swiss chard.

<sup>15</sup> Other tuber and corm vegetables approved for use with CLEANSE 2 EC include: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible burdock, edible canna, bitter and sweet cassava, chayote (root), chufa, dasheen (taro), ginger, leren, tanier, turmeric and bean yam.

CLEANSE 2 EC is not for use on vegetable crops being grown for seed production unless specific use directions are provided in this label.

CLEANSE 2 EC is a selective postemergence herbicide for control of annual and perennial grasses.

CLEANSE 2 EC does not control sedges or broadleaf weeds.

Repeated use of CLEANSE 2 EC (or similar postemergence grass herbicides with the same mode of action) may lead to the selection of naturally occurring biotypes that are resistant to these products in some grass species.

If poor performance occurs and cannot be attributed to adverse weather or application conditions, a resistant biotype may be present. This will most likely occur in fields where other control strategies such as crop rotation, mechanical removal, and other classes of herbicides are not used from year to year.

Do not allow CLEANSE 2 EC to come in contact with desirable grass crops such as corn, rice, sorghum, small grains, or turf, as these and other grass crops will be injured or killed. Minor leaf spotting may occur on treated plants under certain environmental conditions. New foliage is not affected.

#### **Control Symptoms**

Treated grass weeds show a reduction in vigor and growth. Early chlorosis/necrosis of younger plant tissue is followed by a progressive collapse of the remaining foliage. Symptoms will generally be observed in 7 to 14 days, depending on grass species treated and environmental conditions.

## **APPLICATION INFORMATION**

#### **Timing of Applications**

Apply CLEANSE 2 EC postemergence to actively growing grasses according to rate table directions in this label. Applications made to grass plants stressed by insufficient moisture or cold temperatures, or to grass plants exceeding specified growth stages, may result in unsatisfactory control.

In arid regions where irrigation is used to supplement limited rainfall, apply CLEANSE 2 EC as soon as possible, after an irrigation (within 7 days). In arid regions, a second application of CLEANSE 2 EC will generally provide more effective control of perennial grass weeds than a single application. Make second application to actively growing grass 2 to 3 weeks after emergence of new growth.

Cultivation of treated grasses 7 days prior to or within 7 days after application of CLEANSE 2 EC may reduce weed control. DO NOT APPLY CLEANSE 2 EC if rainfall is expected within one hour, since control may be reduced.

## ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE

CROP	ADJUVANT
Alfalfa, Cotton, Dry Bean & Pea (shelled), Edible Podded Legume Vegetables, Peanuts (including perennial), Potato, Soybeans, Succulent Bean & Pea (shelled), Sugar Beet and Sunflower	Always use a crop oil concentrate* at 1.0 qt./A by ground or 1 % v/v (but not less than 1 pt./A) in the finished spray volume by air. 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N or 32%N), or an equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added to CLEANSE 2 EC applications, in addition to the specified rate of crop oil concentrate. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
Artichoke (Globe), Asparagus, Bushberry, Caneberry, Canola, Carrot, Clover, Cranberry, Cucurbits, Flax, Fruiting Vegetables (except Tomato), Garden Beet, Garlic, Head & Stem Brassica Vegetables, Herbs, Hops, Leaf Petioles, Leafy Brassica Greens, Leafy Greens, Mint, Mustard Seed, Onions (Dry Bulb & Green), Peaches, Root Vegetables, Safflower, Sesame, Shallots (Dry Bulb & Green), Strawberry, Sweet Potato (Yam & other Tuberous and Corm Vegetables except Potato) and Tomato	Always use a crop oil concentrate at 1% v/v in the finished spray volume unless tank mix instructions indicate otherwise. Do not add liquid fertilizer for these crops.
Non-Bearing Food Crops, Ornamental Plants	Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v). Do not use crop oil concentrate since it may injure flowers and foliage.
Conifer Trees, Fallow Land (and other non-producing agricultural areas), and Non-Crop or Non-planted Areas	Always use a crop oil concentrate containing at least 15% emulsifier at 1% v/v (but not less than 1 pt./A) in the finished spray volume.

\*Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality, and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

### Ground Application

Use a minimum of 5 gals. and a maximum of 40 gals. of spray solution per acre. Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Under the following conditions a minimum of 10 gals. per acre is required: narrow row soybeans, broadleaf herbicide tank mixes, perennial grasses, volunteer corn, drought or stress conditions, heavy grass pressure or when grasses are at or near maximum height. Failure to use a minimum of 10 gals. per acre under these conditions can result in poor coverage and reduced grass control requiring repeat applications. Spray pressures must reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

Make applications to onions (dry bulbs and green), garlic, and shallots (dry bulbs and green) in a minimum of 20 gals. of spray solution per acre.

### **Air Application**

Use a minimum of 3 gals. of spray solution per acre. Increase spray volumes up to 10 gals. as grass or crop foliage becomes dense. For onions (dry bulbs and green), garlic, or shallots (dry bulbs and green): When applying by air, do not exceed 8 fl. oz./A in a single application. In California, make air applications to onions, garlic or shallots in a minimum of 20 gals. of spray solution per acre.

**NOTE: Crop injury may occur when CLEANSE 2 EC is applied to onions, garlic or shallots with aerial equipment.**

### **Spot Treatment**

When using hand sprayers or high volume sprayers utilizing hand guns, mix 1/4% to 1/2% (0.33 oz. to 0.65 oz. per gal.) CLEANSE 2 EC and treat to wet vegetation, while not allowing runoff of spray solution. For uses requiring crop oil concentrate, include crop oil concentrate at 1% (1.3 oz. per gal.) by volume. For uses requiring non-ionic surfactant, include non-ionic surfactant at 1/4% (0.33 oz. per gal.) by volume.

**NOTE: If CLEANSE 2 EC is applied as a spot treatment, do not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.**

## **CHEMIGATION - ONIONS (Dry Bulb and Green) AND GARLIC SPRINKLER IRRIGATION APPLICATION**

**Do not apply CLEANSE 2 EC by chemigation in the states of Idaho, Montana, Oregon and Washington.**

Apply CLEANSE 2 EC at the high rate specified for annual grasses (16 fl. oz. per acre) when the grass height is at the low end of the range (application to larger grasses may not provide adequate control). Add a crop oil concentrate containing at least 15% emulsifier at 1 quart per acre.

Apply CLEANSE 2 EC in 0.1 to 0.2 acre-inch of water either at the end of a regular irrigation set or as a separate application not associated with a regular irrigation using the least amount of water that provides proper distribution and coverage. Application of more than label specified quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness. Use a metering device to inject the CLEANSE 2 EC into the irrigation water at a constant flow. Constant agitation must be maintained in the chemical supply tank during the entire period of herbicide application. Inject the product with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period.

CLEANSE 2 EC is not to be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

### **Use Precautions**

1. Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.
3. If you have any questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.



5. A person knowledgeable of the chemigation system and responsible for its operation or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
6. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
12. Do not apply when wind speed favors drift beyond the area intended for treatment.

## **RESTRICTIONS AND LIMITATIONS**

- Do not apply if rain is expected within 1 hour of application as control may be unsatisfactory.
- Do not apply a postemergence broadleaf herbicide within one day following application of CLEANSE 2 EC or reduced grass control may result.
- CLEANSE 2 EC is not to be used on vegetable crops being grown for seed production unless specific use directions are provided.
- For canola, do not apply more than 6 fl. oz. of CLEANSE 2 EC (0.08 lb. ai) per acre per season. For clover, flax, mustard seed and radish crops, do not apply more than 16 fl. oz. of CLEANSE 2 EC (0.25 lb. ai) per acre per season. For all other crops, do not apply more than 32 fl. oz. of CLEANSE 2 EC (0.50 lb. ai) per acre per season. Application on Long Island, New York, is restricted to no more than 16 fl. oz. of CLEANSE 2 EC (0.25 lb. ai) per acre per season.
- Do not apply more than 8 fl. oz./A of CLEANSE 2 EC per application to the following crops: asparagus, brassica vegetables (head and stem), bean (succulent), carrot, cranberry, cucurbits, flax, fruiting vegetables (except tomato), garden beet, green onion, herbs, hops, leaf petioles, leafy brassica greens, leafy greens, legume vegetables (edible podded), non-bearing food crops, pea (shelled), pea (succulent), root vegetables, safflower, sesame and strawberry. Do not apply more than 6 fl. oz./A of CLEANSE 2 EC per application to canola or mustard seed. For all other crops, do not apply more than 16 fl. oz. of CLEANSE 2 EC (0.25 lb. ai) per acre per application. Exceeding these directions may result in unacceptable crop injury.
- Do not apply under conditions of stress. Applying CLEANSE 2 EC under conditions that do not promote active grass growth will reduce herbicide effectiveness. These conditions include drought, excessive water, extremes in temperature, low humidity and grasses either partially controlled or stunted from prior pesticide applications. Grasses under these kinds of stressful conditions will not absorb and translocate CLEANSE 2 EC effectively, and will be less susceptible to herbicide activity.
- Best perennial grass control can be obtained if rhizomes or stolons are cut up by preplant tillage practices (discing, plowing, etc.) to stimulate maximum emergence of grass shoots. Cultural practices, such as continuous no-tillage in which the perennial grass rhizomes or stolons are not cut up, result in a very staggered, non-uniform weed emergence. Due to this non-uniform weed emergence, apply no fewer than two CLEANSE 2 EC applications per season per year at the appropriate weed-growth stage rate under continuous no-till conditions.

- Grass crops such as corn, rice, sorghum, small grains, or turf, etc. are highly sensitive to CLEANSE 2 EC.
- While all the vegetable crops on this label have been tested and are tolerant to CLEANSE 2 EC, not all specialty varieties of these crops have been tested. It is advised that, before applying CLEANSE 2 EC to specialty varieties of vegetable crops on this label, crop tolerance be investigated first using a small section of the field. It is possible that injury symptoms can occur. Symptoms may appear as leaf speckling or stunting.
- Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.
- Tank mixes of CLEANSE 2 EC and broadleaf herbicides may result in reduced grass control. If grass regrowth occurs, an additional application of CLEANSE 2 EC may be necessary.

#### **AVOID SPRAY DRIFT.**

Do not allow spray from ground or aerial equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including:

1. Do not spray if wind speeds are or become excessive.
2. Do not spray if wind speed is 10 mph or greater. If sensitive crops or plants are downwind, extreme caution must be used under all conditions.
3. Do not spray if winds are gusty.
4. Use extreme caution when conditions are favorable for drift (high temperatures, drought, low relative humidity), especially when sensitive plants are located nearby.
5. Do not apply when a temperature inversion exists. If inversion conditions are suspected, consult with local weather services before making an application.
6. Further reductions in drift can be obtained by:
  - a) Using large droplet size sprays. Do not use nozzles that produce small droplets. Orient nozzles downward and slightly backward as needed to reduce drift for ground applications.
  - b) Orienting nozzles straight back with the windstream, using straight stream orifices for aerial applications. Use the lowest number of nozzles practical with the largest possible orifice size to obtain the minimum 3 GPA volume. Application height and boom length must be set according to manufacturer's instructions to minimize drift.
  - c) Increasing the volume of spray mixture (for example, a minimum of 10 GPA for ground applications) by using higher flow rate nozzles. Using lower pressure with the appropriate nozzle to obtain higher volumes will also reduce drift.
  - d) Applying as close to target plants as practical while maintaining a good spray pattern for adequate coverage.

**Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

## CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Alfalfa</b> including: Sainfoin Holy Clover Birdsfoot trefoil <sup>(3)</sup>	15 days before grazing, feeding or harvesting (cutting) for forage or hay	6-16 fl. oz. <sup>(4)</sup>	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Do not plant rotational crops until 30 days after application of CLEANSE 2 EC. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Artichoke (Globe)</b>	5 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval. Do not apply more than 32 fl. oz./A of CLEANSE 2 EC (0.5 lb. ai/A) per season.
<b>Asparagus</b>	1 day	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.
<b>Bean, Dry Shelled</b> including: Bean ( <i>Lupinus</i> spp.) Grain Sweet White White Sweet Bean ( <i>Phaseolus</i> spp.) Field Kidney Lima (dry) Navy Pinto Tepary	30 days	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Do not apply more than 16 fl. oz./A of CLEANSE 2 EC per application. For repeat applications, make on a minimum of a 14-day interval. Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn.

(continued)

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC (continued)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Bean, Dry Shelled</b> including: Bean ( <i>Vigna</i> spp.) Adzuki Bean Black-eyed Pea Catjang Cowpea Crowder Pea Moth Bean Mung Bean Rice Bean Southern Pea Urd Bean Broad (dry) Chickpea (garbanzo) Guar Lablab Bean Lentil	30 days	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Do not apply more than 16 fl. oz./A of CLEANSE 2 EC per application.  For repeat applications, make on a minimum of a 14-day interval.  Refer to appropriate Table for reduced rate directions for the control of small annual grasses.  The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn.
<b>Bean, Succulent Shelled</b> including: Bean ( <i>Phaseolus</i> spp.) Broad Bean (succulent) Lima Bean (green) Bean ( <i>Vigna</i> spp.) Black-eyed Pea Cowpea Southern Pea	21 days	6-8 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Refer to appropriate Table for reduced rate directions for the control of small annual grasses.  Do not apply more than one (1) application of CLEANSE 2 EC per acre per season.  The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn.
<b>Beet, Garden</b>	30 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application.  For repeat applications, make on a minimum of a 14-day interval.

(continued)

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC** *(continued)*

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Brassica Vegetables, Head &amp; Stem</b> including: Broccoli Brussels sprouts Cabbage Cauliflower	30 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.
<b>Bushberry,</b> including: Aronia berry, Blueberry, highbush Chilean guava Cranberry, highbush Currant, black Currant, buffalo Currant, native Currant, red Elderberry European barberry Gooseberry Honeysuckle, edible Huckleberry Jostaberry Juneberry Saskatoon berry Salal Sea buckthorn Cultivars, varieties and/or hybrids of these	14 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. Do not apply more than 32 fl. oz. of CLEANSE 2 EC (0.5 lb. ai/A) per season. For repeat applications, make a minimum of a 14-day interval. Apply at the base of the plant where grassy weeds are growing near the ground. Do not apply to low growing berries. Do not apply to Bushberry grown for root stock.

*(continued)*

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC** *(continued)*

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Caneberry</b> , including: Blackberry Loganberry Raspberry, black Raspberry, red Raspberry, wild Cultivars, varieties and/or hybrids of these	7 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. Do not apply more than 32 fl. oz./A of CLEANSE 2 EC (0.5 lb ai/A) per season. For repeat applications, make on a minimum of a 14-day interval. Apply at the base of the plant where grassy weeds are growing near the ground. Do not apply to low growing berries. Do not apply to Caneberry grown for root stock.
<b>Canola*</b>	70 days	4-6 fl. oz.	1.0% v/v in the finished spray volume	Do not apply after crop has begun bolting. Crop injury may occur when CLEANSE 2 EC is applied during the bloom period. Do not apply more than 6 fl. oz./A of CLEANSE 2 EC in a single application. Do not exceed 6 fl. oz./A of CLEANSE 2 EC in a season. *Not for use in California unless accompanied by a supplemental label.
<b>Carrot</b>	30 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.
<b>Clover</b>	15 days before grazing, feeding, or harvesting (cutting) for forage or hay	6-16 fl. oz.	1.0% v/v in the finished spray volume	For use on clover grown in the states of Idaho, Oregon and Washington only. Do not exceed 16 fl. oz./A of CLEANSE 2 EC in a season.

*(continued)*

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC** *(continued)*

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<p><b>Corn, Field</b> For burndown of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn.</p>	90 days	3 fl. oz.	Do not use COC or MSO with CLEANSE 2 EC in this use pattern.	<p>Do not make more than 1 application of CLEANSE 2 EC per season. Do not apply more than 3 fl. oz./A of CLEANSE 2 EC per season. To control the existing stand, replant no sooner than 6 days after application. See Directions for Use in Roundup Ready Field Corn (Burndown) table.</p>
<p><b>Cotton</b></p>	60 days	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	<p>Do not graze treated fields or feed treated forage or hay to livestock. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p>
<p><b>Cranberry</b></p>	30 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	<p>Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. Do not apply between the "hook" stage and full fruit set. For repeat applications, make on a minimum of a 14-day interval.</p>
<p><b>Cucurbits</b> including: Cantaloupes (all) Chayote (fruit) Chinese Wax Gourd Citron Melon Cucumber Gherkin Gourd, Edible Honeydew Melon Muskmelons (all) Pumpkin Squash (all) Watermelon</p>	14 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	<p>Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.</p>

*(continued)*

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC** *(continued)*

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Fallow Land Conifer Trees</b> (and other non-producing agricultural areas) <b>Non-Crop or Non-Planted Areas</b>	N/A	6-16 fl. oz.	1.0% v/v (but not less than 1 pt./A) in the finished spray volume using a crop oil concentrate containing at least 15% emulsifier	Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop.
<b>Flax*</b>	60 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Apply prior to bloom. Crop injury may occur when CLEANSE 2 EC is applied during the bloom period.  Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application.  Do not exceed 16 fl. oz./A of CLEANSE 2 EC in a season.  *Not for use in California unless accompanied by a supplemental label.
<b>Fruiting Vegetables</b> (except Tomato) including: Eggplant Groundcherry Pepino Peppers (all) Tomatillo	20 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application.  For repeat applications, make on a minimum of a 14-day interval.

*(continued)*



**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC (continued)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Herbs</b> including: Angelica Balm Basil Borage Burnet Camomile Catnip Chervil (dried) Chive Chive, Chinese Clary Coriander (leaf) Costmary Culantro (leaf) Curry (leaf) Dill (dillweed) Horehound Hyssop Lavender Lovage (leaf) Marigold Marjoram ( <i>Origanum</i> spp.) Nasturtium Parsley (dried) Pennyroyal Rosemary Rue Sage Savory, Summer and Winter	14 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	CLEANSE 2 EC has not been tested on all herbs and herb varieties. It is the responsibility of the user to test CLEANSE 2 EC on a small portion of the crop to be treated before treating the entire field.  Crop tolerance must be verified to CLEANSE 2 EC on a small area of the herb crop, at the desired CLEANSE 2 EC rate and with the same crop oil concentrate that will be used on the herb field. If no crop response is evident seven (7) days after treatment, CLEANSE 2 EC may be used on the entire field at the rate tested and with the same crop oil used in the tolerance test.  Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application.  For repeat applications, make on a minimum of a 14-day interval.
<b>Hops</b>	21 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application.  For repeat applications, make on a minimum of a 14-day interval.

*(continued)*

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC** *(continued)*

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Leaf Petioles</b> including: Cardoon Celery Celtuce Chinese Celery Fennel, Florence (finocchio) Rhubarb Swiss Chard	30 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.
<b>Leafy Brassica Greens</b> including: Broccoli Raab Cabbage, Chinese (bok choy) Collards Kale Mizuna Mustard Greens Mustard Spinach Rape Greens Turnip Greens	14 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.

*(continued)*

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC (continued)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Leafy Greens</b> including: Amaranth Chinese Spinach Leafy Amaranth Tampala Arugula (rocket) Chervil Chrysanthemum, Edible-leaved Garland Corn Salad Cress Garden Upland (yellow rock and winter) Dandelion Dock (sorrel) Endive (escarole) Lettuce, Head and Leaf Orach Parsley Purslane Garden Winter Radicchio (red chicory) Spinach New Zealand Vine (indian and malabar)	14 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.

*(continued)*

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC (continued)**

<b>Crops<sup>(1)</sup></b>	<b>Minimum Time From Application to Harvest (PHI)</b>	<b>Use Rates Per Acre</b>	<b>Crop Oil Concentrate Rates Per Acre<sup>(2)</sup></b>	<b>Special Use Instructions</b>
<b>Legume Vegetables, Edible Podded</b> including: Bean ( <i>Phaseolus</i> spp.) Runner Snap Wax Bean ( <i>Vigna</i> spp.) Asparagus Chinese Longbean Moth Yardlong Jackbean Pea ( <i>Pisum</i> spp.) Dwarf Edible-pod Snow Sugar Snap Pigeon Sword Bean	21 days	6-8 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. Do not apply more than one (1) application per acre per season. For peas, apply before bloom, but no later than 21 days before harvest. Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Mint</b>	21 days	6-16 fl. oz. <sup>(4)</sup>	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air	Do not apply more than 16 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.
<b>Mustard Seed*</b>	75 days	4-6 fl. oz.	1.0% v/v in the finished spray volume	Do not apply after crop has begun bolting. Crop injury may occur when CLEANSE 2 EC is applied during the bloom period. Do not apply more than 6 fl. oz./A of CLEANSE 2 EC in a single application. Do not exceed 16 fl. oz./A of CLEANSE 2 EC in a season. *Not for use in California unless accompanied by a supplemental label.

*(continued)*

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC (continued)**

<b>Crops<sup>(1)</sup></b>	<b>Minimum Time From Application to Harvest (PHI)</b>	<b>Use Rates Per Acre</b>	<b>Crop Oil Concentrate Rates Per Acre<sup>(2)</sup></b>	<b>Special Use Instructions</b>
<b>Onion</b> (Dry Bulb Only) <b>Garlic</b> <b>Shallot</b> (Dry Bulb Only)	45 days	6-16 fl. oz. <sup>(7), (8)</sup>	1.0% v/v in the finished spray volume	Minimum of 20 gals./A spray volume by ground in entire U.S. Minimum of 20 gals./A spray volume by air in California. <sup>(9)</sup> In states other than California, air applications to onions, garlic or shallots must be made in a minimum of 10 gals./A.
<b>Onions, Green</b> including: Green Eschalot Green Shallots Japanese Bunching Onions Leeks Scallions or Spring Onion	14 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.
<b>Ornamentals</b>	N/A	6-16 fl. oz.	Do not use crop oil concentrate since it may injure flowers and foliage. See Special Use Instructions.	Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).
<b>Non-Bearing Food Crops</b>	N/A	6-8 fl. oz. <sup>(8)</sup>	Do not use crop oil concentrate since it may injure flowers and foliage. See Special Use Instructions.	Sugar maples cannot be tapped for syrup within one year of CLEANSE 2 EC application.
<b>Pea, Dry Shelled</b> including: Pea ( <i>Pisum</i> spp.) Field Pigeon	30 days	6-8 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC per application. Do not apply more than one (1) application of CLEANSE 2 EC per acre per season. Apply before bloom but not later than 30 days prior to harvest. <sup>(10)</sup> Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn.

*(continued)*

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC (continued)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<p><b>Pea, Succulent Shelled</b> including: Pea (<i>Pisum</i> spp.) English Pea Garden Pea Green Pea Pigeon Pea</p>	21 days	6-8 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(3)</sup>	<p>Do not apply more than 8 fl. oz./A of CLEANSE 2 EC per application.</p> <p>Do not apply more than one (1) application per acre per season.</p> <p>Apply before bloom but not later than 21 days prior to harvest.<sup>(10)</sup></p> <p>Refer to appropriate Table for reduced rate directions for the control of small annual grasses.</p> <p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn.</p>
<b>Peach</b>	14 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	<p>Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application.</p> <p>Do not apply more than 32 fl. oz./A of CLEANSE 2 EC (0.5 lb. ai/A) per season.</p> <p>For repeat applications, make on a minimum of a 14-day interval.</p> <p>Do not apply to Peach grown for root stock.</p>
<b>Peanut (including Perennial)</b>	40 days	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(3)</sup>	<p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p>
<b>Potato</b>	30 days	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(3)</sup>	<p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p>

(continued)

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC (continued)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Radish</b>	15 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. Do not apply more than 16 fl. oz./A of CLEANSE 2 EC (0.25 lb. ai) per season. For repeat applications, make on a minimum of a 14-day interval.
<b>Root Vegetables</b> (except Radish), including: Chicory Ginseng Horseradish Turnip	30 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.
<b>Safflower</b>	70 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.
<b>Sesame</b>	14 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply during flowering. Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.
<b>Soybean</b>	60 days	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(3)</sup>	Do not graze treated fields or feed treated forage or hay to livestock. Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Strawberry</b>	4 days	6-8 fl. oz.	1.0% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A of CLEANSE 2 EC in a single application. For repeat applications, make on a minimum of a 14-day interval.

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLEANSE 2 EC (continued)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Sugar Beet</b>	40 days	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(3)</sup>	Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Sunflower</b>	70 days	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(3)</sup>	The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Sweet Potato, Yam and other Tuberous and Corm Vegetables (except Potato) including:</b> Artichoke Chinese Jerusalem Cassava Bitter Sweet Ginger	30 days	6-16 fl. oz.	1.0% v/v in the finished spray volume	The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Tomato</b>	20 days	6-16 fl. oz.	1.0% v/v in the finished spray volume	For repeat applications, make on a minimum of a 14-day interval.

N/A = Not Applicable

- <sup>(1)</sup> Do not use CLEANSE 2 EC on vegetable crops being grown for seed production unless specific use directions are provided.
- <sup>(2)</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the Addition of Adjuvant and Crop Oil Concentrate section for further information.
- <sup>(3)</sup> CLEANSE 2 EC may be applied to seedling or established alfalfa grown for seed, hay, silage, green chop or direct grazing.
- <sup>(4)</sup> For weed control in established alfalfa and mint, the minimum use rate is 10 fl. oz./A of CLEANSE 2 EC.
- <sup>(5)</sup> 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N or 32%N), or an equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added to CLEANSE 2 EC applications, in addition to the specified rate of crop oil concentrate.
- <sup>(6)</sup> Do not apply CLEANSE 2 EC plus 2,4-DB as a tank mix to alfalfa unless the 60 day feeding, grazing, and harvesting restriction on the 2,4-DB label can be observed.



- (7) For ground applications to garlic or shallots, do not exceed 8 fl. oz./A of CLEANSE 2 EC in a single application. For air applications to onion, garlic or shallots, do not exceed 8 fl. oz./A of CLEANSE 2 EC in a single application. For garlic and shallots, do not exceed 2 applications of CLEANSE 2 EC per season. In CA, for air applications to onions, do not exceed 2 applications per season.
- (8) If CLEANSE 2 EC is applied as a spot treatment to onions, garlic, shallots, or non-bearing food crops, care must be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.
- (9) In California, do not apply CLEANSE 2 EC to onions, garlic, or shallots until the crop has at least two full leaves. In California, 14-day spray intervals are necessary between the application of CLEANSE 2 EC and liquid nitrogen or other herbicide applications. Injury to crop may occur when shorter intervals are observed.
- (10) Applications of CLEANSE 2 EC to peas during the bloom period could result in severe crop injury, including loss of yield and delayed maturity.

## NON-BEARING FOOD CROPS

**DO NOT APPLY CLEANSE 2 EC TO NON-BEARING FRUIT OR NUT CROPS THAT ARE GROWN FOR ROOT STOCK.**

**Crop injury to non-bearing fruit and nut crops can occur if CLEANSE 2 EC is improperly applied. Do not apply CLEANSE 2 EC directly over the top of these plant types. Instead, direct spray at the base of the plant where grassy weeds are growing near the ground.**

Non-bearing fruit and nut crops are plants which will not bear fruit or nuts for at least one year following CLEANSE 2 EC application.

COMMON NAME	SCIENTIFIC NAME
Apples	<i>Malus</i> spp.
Berries	<i>Vaccinium</i> spp.
	<i>Rubus</i> spp.
Cherry, Sweet	<i>Prunus avium</i>
Citrus Fruits	<i>Citrus</i> spp.
Grapes	<i>Vitis</i> spp.
Olives	<i>Olea</i> spp.
Peach	<i>Prunus persica</i>
Pears	<i>Pyrus communis</i>
Prunes	<i>Prunus</i> spp.
Stone Fruits	<i>Prunus</i> spp.
Strawberries	<i>Fragaria</i> spp.
Tree Nuts	
Almond	<i>Prunus dulcis</i>
Filbert	<i>Corylus maxima</i>
Pecan	<i>Carya illinoensis</i>
Pistachio	<i>Pistacia vera</i>
Walnut	<i>Juglans</i> spp.

## CONIFER TREES

CLEANSE 2 EC can be used to control labeled grasses in Christmas tree farms, conifer nurseries, and conifer plantations (but not in forests).

COMMON NAME	SCIENTIFIC NAME
Arborvitae, American	<i>Thuja occidentalis</i>
Cedars	<i>Cedrus</i> spp.
Cypress	<i>Taxodium</i> spp.
Fir, Douglas	<i>Pseudotsuga menziesii</i>
Firs	<i>Abies</i> spp.
Hemlock, Canadian/Eastern	<i>Tsuga canadensis</i>
Hemlock, Western	<i>Tsuga heterophylla</i>
Pines	<i>Pinus</i> spp.
Spruces	<i>Picea</i> spp.
Yew	<i>Taxus</i> spp.

## NON-CROP OR NON-PLANTED AREAS

The following areas are considered non-crop or non-planted areas: rights-of-way including railroads, highways, roads, dividers, medians, pipelines, public utility lines, pumping stations, transformer stations and substations. Around airports, electric utilities, commercial buildings, manufacturing plants, storage yards, rail yards, fence lines, parkways, and post-harvest croplands. Also beneath greenhouse benches and around golf courses.

**ANNUAL GRASSES**  
(EXCEPT FOR IN ESTABLISHED ALFALFA AND MINT)

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.
- Do not apply more than 8 fl. oz./A of CLEANSE 2 EC per application to the following crops: asparagus, carrot, cranberry, cucurbits, flax, fruiting vegetables (except tomato), garden beets, green onion, head and stem brassica vegetables, herbs, hops, leaf petioles, leafy brassica greens, leafy greens, non-bearing food crops, root vegetables, safflower, sesame and strawberry. Do not apply more than 6 fl. oz./A of CLEANSE 2 EC per application to canola or mustard seed.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	RATE FL. OZ./ ACRE	HIGH RATE <sup>(1)</sup>
Barnyardgrass	<i>Echinochloa crus-galli</i>	2 to 8	6	8
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	2 to 6	6	8
Brome				
California	<i>Bromus carinatus</i>	2 to 6	6	8
Cheatgrass	<i>Bromus secalinus</i>	2 to 6	6	8
Downy	<i>Bromus tectorum</i>	2 to 6	6	8
Ripgut	<i>Bromus diandrus</i>	2 to 6	6	8
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	6	8
Crabgrass				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	6	8
Large	<i>Digitaria sanguinalis</i>	2 to 6**	6	8
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	6	8
Southern	<i>Digitaria ciliaris</i>	2 to 6**	6	8
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	6	8
Fall Panicum	<i>Panicum dichotomiflorum</i>	2 to 8	6	8
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	6	8
Foxtail				
Giant	<i>Setaria faberi</i>	2 to 12	6	8
Green	<i>Setaria viridis</i>	2 to 8	6	8
Yellow	<i>Setaria glauca</i>	2 to 8	6	8

(continued)

## ANNUAL GRASSES (continued)

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	RATE FL. OZ./ ACRE	HIGH RATE <sup>(1)</sup>
Goosegrass	<i>Eleusine indica</i>	2 to 6**	6	8
Itchgrass	<i>Rottboellia exaltata</i>	2 to 6	6	8
Junglerice	<i>Echinochloa colona</i>	2 to 6	6	8
Lovegrass (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	6	8
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	6	8
Red Rice	<i>Oryza sativa</i>	1 to 3	6	8
Ryegrass				
Hardy	<i>Lolium remotum</i>	2 to 6	6	8
Italian	<i>Lolium multiflorum</i>	2 to 6	6	8
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	6	8
Shattercane	<i>Sorghum bicolor</i>	6 to 18	6	8
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	6	8
Sprangletop				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	6	8
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	6	8
Mexican	<i>Leptochloa uninervia</i>	2 to 6	6	8
Red	<i>Leptochloa filiformis</i>	2 to 6	6	8
Texas Panicum	<i>Panicum texanum</i>	2 to 6	6	8
Volunteer Cereals <sup>(2)</sup>				
Barley	<i>Hordeum vulgare</i>	2 to 6	6	8
Oats	<i>Avena sativa</i>	2 to 6	6	8
Rye	<i>Secale cereale</i>	2 to 6	6	8
Wheat	<i>Triticum aestivum</i>	2 to 6	6	8
Volunteer Corn <sup>(3)</sup>	<i>Zea mays</i>	4 to 12	4	6
Volunteer Corn <sup>(3)</sup>	<i>Zea mays</i>	12 to 24	6	8
Volunteer Corn (S.R.) <sup>(4)</sup>	<i>Zea mays</i>	4 to 12	8 (suppression only)	
Volunteer Grain Sorghum	<i>Sorghum bicolor</i>	8 to 12	6	8

(continued)

**ANNUAL GRASSES** (continued)

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	RATE FL. OZ./ ACRE	HIGH RATE <sup>(1)</sup>
Wild Oats	<i>Avena fatua</i>	2 to 6	6	8
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	6	8
Witchgrass	<i>Panicum capillare</i>	2 to 8	6	8
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	6	8
* Generally occurs between 3-leaf stage and tillering. ** Length of lateral growth.				
<p>(1) Rates higher than 8 fl. oz./A of CLEANSE 2 EC may be applied in certain geographic areas, cropping situations, or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 8 to 16 fl. oz./A of CLEANSE 2 EC may be applied. Do not apply more than 8 fl. oz./A of CLEANSE 2 EC per application to the following crops: asparagus, carrot, cranberry, cucurbits, flax, fruiting vegetables (except tomato), garden beet, green onion, head and stem brassica vegetables, herbs, hops, leaf petioles, leafy brassica greens, leafy greens, non-bearing food crops, root vegetables, safflower, sesame and strawberry. Do not apply more than 6 fl. oz./A of CLEANSE 2 EC per application to canola or mustard seed.</p> <p>(2) When a cereal grain crop (such as wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum CLEANSE 2 EC use rate for control is 8 fl. oz./A.</p> <p>(3) Includes Roundup Ready®, Liberty Link® and IMI-CORN® volunteer corn.</p> <p>(4) Sethoxydim resistant volunteer corn.</p>				

## ANNUAL & PERENNIAL GRASS CONTROL IN ESTABLISHED ALFALFA AND MINT WITH CLEANSE 2 EC

GRASS SPECIES	WEED STAGE	RATE FL. OZ./ ACRE	HIGH RATE
Annual & Perennial Grasses Listed in Grass Table	See Table	10	16
<p><b>Mowing:</b> The best control of annual grasses can be achieved by applying CLEANSE 2 EC before grass weeds are mowed. Once a grass is mowed it becomes tougher to control, as much of the available leaf surface has been removed. In areas without a killing frost, some annuals can over-winter after having been mowed multiple times. These grasses form large crowns and may contain many viable buds. These grasses, even though they may be an annual grass, may require repeated applications of CLEANSE 2 EC for partial or complete control.</p> <p><b>Irrigated Alfalfa and Mint:</b> Irrigation practices can be very critical to the successful use of CLEANSE 2 EC in established alfalfa and mint and may be necessary to initiate active growth of the weeds prior to application. Generally, applications 2 to 4 days after an irrigation are most effective. Irrigation made shortly after application (2 days) can be effective, but more consistent grass control occurs when the irrigation is made before the application.</p> <p><b>Aerial Application:</b> Apply CLEANSE 2 EC in a minimum of 10 GPA in established alfalfa and mint when applying by air.</p> <p><b>Annual Grass Control:</b> Apply CLEANSE 2 EC at the grass sizes indicated in the Annual Grass Table and rates indicated above (10 to 16 fl. oz/A). If a grass has been cut, apply CLEANSE 2 EC after active growth has resumed and regrowth has reached the minimum height and before it reaches the maximum height indicated. Apply before the alfalfa/mint canopy covers the grasses and interferes with the spray coverage. Some annual grasses are spring- and summer-germinating plants, while others are fall-germinating plants, and the time they are actively growing and most susceptible to CLEANSE 2 EC may vary from region to region. Also, some annuals germinate over an extended period of time, and because control of small grasses is desired, applications after each weed flush may be required. As a general rule, spray spring and summer-germinating grasses as early in the season as possible, after initial green-up. Spray fall-germinating weeds in the fall soon after they begin growing but before any damage is done due to frost. Late fall applications may be less effective due to environmental conditions, such as frost, slower plant growth, or the onset of flowering.</p> <p><b>Perennial Grass Control:</b> CLEANSE 2 EC effectively controls perennial grasses such as bermudagrass, Johnsongrass, quackgrass, wirestem muhly, tall fescue, foxtail barley and orchardgrass. Due in part to lack of tillage, perennial grasses are more difficult to control in a perennial crop such as established alfalfa or mint. A program of repeated applications is usually necessary for best results. The best way to control perennial grasses is to do so in the year of stand establishment before rhizomes and stolons become large and difficult to kill.</p> <p>Use the high rate under heavy grass pressure and/or when grasses are at or near maximum height.</p> <p>Always add a crop oil concentrate at 1 qt./A by ground or 1.0% v/v (but not less than 1 pt./A) to the finished spray volume by air.</p>			

## ANNUAL BLUEGRASS CONTROL WITH CLEANSE 2 EC

GRASS SPECIES	WEED STAGE	RATE FL. OZ./ ACRE	HIGH RATE
Annual Bluegrass ( <i>Poa annua</i> )	to 4-leaf	6*	16
<p>*Use a minimum of 8 fl. oz./A of CLEANSE 2 EC to control annual bluegrass in seedling and established alfalfa and mint.</p> <ul style="list-style-type: none"> <li>• Apply under favorable soil moisture and humidity that exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).</li> <li>• Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.</li> <li>• Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.</li> <li>• Always add a crop oil concentrate at 1 qt./A by ground to the finished spray volume.</li> </ul>			

## ROUNDUP READY FIELD CORN (BURNDOWN)

APPLICATIONS RATES		
GRASS SPECIES	WEED SIZE (inches)	Rate when applied alone or with glyphosate
Field Corn	Up to 12	3 fl. oz./A
<p>For control of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn. Care must be taken to avoid in-field boom (spray) overlaps or excessive injury may occur. Replant no sooner than 6 days after application.</p> <p><b>Adjuvant specifications:</b> NIS at 0.25% v/v plus AMS at 2.5 to 4 lbs./A. Do not use a COC or MSO with CLEANSE 2 EC in this use pattern.</p>		

## DIRECTIONS FOR REDUCED RATE TO CONTROL

### SMALL ANNUAL GRASSES IN CANOLA, BEAN & PEA (DRY SHELLED) (INCLUDING SOYBEANS), EDIBLE PODED LEGUME VEGETABLES, FLAX, MUSTARD SEED, BEAN & PEA (SUCCULENT SHELLED) AND SUGAR BEET

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Regrowth by tillering may occur if application is made when plants are stressed by lack of moisture, excessive moisture, low temperatures and/or under very low humidity.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (inches)	RATE FL.OZ/ ACRE <sup>(1)</sup>
Barnyardgrass	<i>Echinochloa crus-galli</i>	1 to 4	4
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	1 to 4	5
<b>Crabgrass</b>			
Large	<i>Digitaria sanguinalis</i>	1 to 3*	4
Large	<i>Digitaria sanguinalis</i>	1 to 4*	5
Smooth	<i>Digitaria ischaemum</i>	1 to 3*	4
Smooth	<i>Digitaria ischaemum</i>	1 to 4*	5
Southern	<i>Digitaria ciliaris</i>	1 to 4*	5
Fall Panicum	<i>Panicum dichotomiflorum</i>	1 to 4	4
<b>Foxtail</b>			
Giant	<i>Setaria faberi</i>	1 to 4	4
Green	<i>Setaria viridis</i>	1 to 4	4
Millet	<i>Setaria italica</i>	1 to 4	5
Yellow	<i>Setaria glauca</i>	1 to 4	4
Seedling Johnsongrass	<i>Sorghum halepense</i>	1 to 6	5
Shattercane	<i>Sorghum bicolor</i>	4 to 10	4
Texas Panicum	<i>Panicum texanum</i>	1 to 4	5
<b>Volunteer Cereals</b>			
Barley	<i>Hordeum vulgare</i>	1 to 4	5
Oats	<i>Avena sativa</i>	1 to 4	5
Wheat	<i>Triticum aestivum</i>	1 to 4	5
Volunteer Corn**	<i>Zea mays</i>	4 to 12	4
Wild Oats	<i>Avena fatua</i>	1 to 4	5
Wild Proso Millet	<i>Panicum millaceum</i>	1 to 6	4
* Length of lateral growth			
** Not sethoxydim resistant Corn			
(1) Always add a crop oil concentrate at 1 qt./A by ground application to the finished spray volume.			



## PERENNIAL GRASSES

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.
- Do not apply more than 8 fl. oz./A of CLEANSE 2 EC per application to the following crops: asparagus, carrot, cranberry, cucurbits, flax, fruiting vegetables (except tomato), garden beet, green onion, head & stem brassica vegetables, herbs, hops, leaf petioles, leafy brassica greens, leafy greens, non-bearing food crops, root vegetables, safflower, sesame and strawberry. Do not apply more than 6 fl. oz./A of CLEANSE 2 EC per application to canola or mustard seed.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (inches)	RATE FL. OZ./ACRE	HIGH RATE
<b>Bermudagrass</b>	<i>Cynodon dactylon</i>			
First Application		3 (or up to 6" runners)	8	16
Repeat Application(s) (if regrowth occurs)		3 (or up to 6" runners)	8	16
<b>Fescue, Tall</b>	<i>Festuca arundinacea</i>			
First Application		4 to 8	8	16
Repeat Application(s) (if regrowth occurs)		4 to 8	8	16
<b>Foxtail Barley</b>	<i>Hordeum jubatum</i>			
First Application		2 to 6	8	16
Repeat Application(s) (if regrowth occurs)		2 to 6	8	16
<b>Orchardgrass</b>	<i>Dactylis glomerata</i>			
First Application		4 to 8	8	16
Repeat Application(s) (if regrowth occurs)		4 to 8	8	16
<b>Quackgrass*</b>	<i>Elytrigia repens</i>			
First Application		4 to 12	8	16
Repeat Application(s) (if regrowth occurs)		4 to 12	8	16
<b>Rhizome Johnsongrass</b>	<i>Sorghum halepense</i>			
First Application		12 to 24	8	16

(continued)

**PERENNIAL GRASSES** (continued)

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (inches)	RATE FL. OZ./ACRE	HIGH RATE
Repeat Application(s) (if regrowth occurs)		6 to 18	6	8
<b>Wirestem Muhly</b>	<i>Muhlenbergia frondosa</i>			
First Application		4 to 8	8	16
Repeat Application(s) (if regrowth occurs)		4 to 8	8	16
<b>Perennial Bluegrass*</b>				
<b>Roughstalk</b>	<i>Poa trivialis</i>			
<b>Kentucky</b>	<i>Poa pratensis</i>			
First Application		2 to 4	8	16
Repeat Application(s) (if regrowth occurs)		2 to 4	8	16
<b>Bentgrass*</b>	<i>Agrostis spp.</i>			
First Application		2 to 4	-	16
Repeat Application(s) (if regrowth occurs)		2 to 4	-	16

\*Control of quackgrass, perennial bluegrass and bentgrass with CLEANSE 2 EC may be enhanced by adding AMS at 2.5 to 4.0 lbs./A.

## TANK MIXES

The labels for each of the herbicides labeled for tank mixing with CLEANSE 2 EC are unique to the characteristics of those products and contain restrictions and limitations that may be more restrictive than the CLEANSE 2 EC label in certain considerations. Those concerns may include, but are not limited to:

1. Geographic restrictions - all products are not registered for use in all areas and rates may vary from one region of labeled use to another;
2. Crop rotation restrictions;
3. Applicator certification requirements;
4. Worker safety rules (e.g. protective clothing requirements, reentry time);
5. Soil type or soil characteristics;
6. Maximum dosage or number of applications allowed per season;
7. Rain free period required;
8. Application timing (e.g. pre-harvest interval);
9. Do not exceed the total season rates.

**THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.**

## **TANK MIX APPLICATION OF CLEANSE 2 EC AND BROADLEAF HERBICIDES FOR CONTROL OF GRASSES AND BROADLEAF WEEDS**

- Apply only to actively growing grass and broadleaf weeds at specified height or growth stage listed on each label.
- Apply when the first grass or broadleaf weed species in a mixed population reaches the specified height or growth stage for treatment.
- Apply under favorable soil moisture and humidity which exist a few days after rainfall or within seven days after irrigation.
- Always add the appropriate adjuvant to the spray mix at the rate specified for each specific tank mix combination.
- Tank mix applications may sometimes result in reduced grass control and possible increases in crop injury as compared to either product used alone. If regrowth occurs, or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC, as specified in the respective size and rate tables.
- Do not tank mix CLEANSE 2 EC when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.

### **MIXING INSTRUCTIONS**

Mixing and compatibility qualities must be verified by a jar test.

Failure to agitate the spray solution may result in improper mixing of the herbicides and unsatisfactory weed control.

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, add the correct amount of CLEANSE 2 EC. Agitation should create a rippling or rolling action on the water surface.
3. If tank mixing CLEANSE 2 EC with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates, and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
4. Add any required adjuvants (crop oil concentrate, non-ionic surfactant and/or nitrogen solution).
5. Fill spray tank to desired level with water. Continue agitation until all spray solution has been applied.

### **INFORMATION ON ANTAGONISM**

Tank mixes of CLEANSE 2 EC with postemergence broadleaf herbicides have shown some reduction or failure to control certain grass species which would have otherwise been controlled when CLEANSE 2 EC is applied alone. Activity of the postemergence broadleaf herbicide in the tank mix is not affected.

## ALFALFA

**Table 1. CLEANSE 2 EC TANK MIXES WITH BROADLEAF HERBICIDES FOR ALFALFA**

(Refer to the tables above for specific grasses and growth stages)

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
CLEANSE 2 EC + 2,4-DB <sup>(4)</sup>	10 to 16 fl. oz. + Refer to 2,4-DB label	10 to 16 fl. oz. + Refer to 2,4-DB label	1.0%	1.0%
CLEANSE 2 EC + PURSUIT DG <sup>(5)</sup> or PURSUIT <sup>(5)</sup>	10 to 16 fl. oz. + 1.08 to 2.16 oz. or 3 to 6 fl. oz.	-	1.0%	1.0%
CLEANSE 2 EC + BUCTRIL <sup>®</sup> 2L <sup>(6)</sup> or BUCTRIL GEL <sup>(6,7)</sup>	10 to 16 fl. oz. + 1.0 to 1.5 pts. or 0.5 to 0.75 pt.	-	0.5%	0.5%

- (1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC alone (without a tank mix herbicide), according to the appropriate size and rate directions.
- (2) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Do not tank mix in these situations.
- (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- (4) CLEANSE 2 EC plus 2,4-DB may increase the severity of crop injury when tank mixed. Alfalfa plants will generally outgrow this temporary crop injury within a few weeks.
- (5) Before using this tank mix, read and understand the PURSUIT or PURSUIT DG labels for geographical restrictions and restrictions regarding alfalfa growth stage and type. Failure to do so can result in crop injury to alfalfa. Do not feed, graze, or harvest alfalfa for 30 days following an application of PURSUIT to alfalfa.
- (6) In the states of Washington, Oregon, Idaho, Montana, Wyoming, Colorado, Utah, Nevada and the western halves of North Dakota, South Dakota, Nebraska, and Kansas: The CLEANSE 2 EC plus BUCTRIL or BUCTRIL GEL tank mix must be applied in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 2 trifoliolates. Unacceptable crop injury may occur to alfalfa seedlings less than the 2 trifoliolate leaf stage. CLEANSE 2 EC plus BUCTRIL or BUCTRIL GEL applications made when temperatures are expected to exceed 80°F at (and 3 days following) application can result in unacceptable crop injury. In the states not listed above, apply in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 4 trifoliolate leaves. When alfalfa stand is uneven and conditions favor leaf burn, unacceptable crop injury may occur to alfalfa in the 2 trifoliolate or smaller stage of growth. CLEANSE 2 EC plus BUCTRIL or BUCTRIL GEL applications made when temperatures are expected to exceed 70°F at (and 3 days following) application can result in unacceptable crop injury. Crop leaf burn can occur following CLEANSE 2 EC plus BUCTRIL or BUCTRIL GEL application. Warm, humid conditions may enhance leaf burn. New crop growth will not be affected.
- (7) Do not apply when alfalfa is under moisture, temperature, insect or disease stress, or has been stressed by other pesticide carryover or application.

## CANOLA\*

**Table 2. REDUCED RATE CLEANSE 2 EC TANK MIXES WITH BROADLEAF HERBICIDES FOR CANOLA**  
(Refer to the tables above for specific grasses and growth stages.)

PRODUCT	APPLICATION RATES/ACRE		AMMONIUM SULFATE	
	ANNUAL GRASSES <sup>(1)</sup>	PERENNIAL GRASSES	GROUND	AIR
CLEANSE 2 EC <sup>(2)</sup> + LIBERTY <sup>(3)</sup>	4 to 5 fl. oz. + 34 fl. oz.	-	3.0 lbs.	3.0 lbs.

(1) Annual grasses and sizes controlled with these tank mixtures are those that are identified in the **DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES** table.  
 (2) Do not apply CLEANSE 2 EC tank mix during or after bolting or flowering or crop injury may occur.  
 (3) For use only on LibertyLink® canola.  
 \*Not for use in California unless accompanied by a supplemental label.

## COTTON

**Table 3. CLEANSE 2 EC TANK MIXED WITH COBRA® AND MSMA APPLIED POST DIRECTED TO COTTON**

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup>	COMMENTS
	ANNUAL GRASSES	PERENNIAL GRASSES	V/V GROUND	
CLEANSE 2 EC <sup>(4)</sup> + COBRA + MSMA (4.0 lbs./gal.) or MSMA (6.6 lbs./gal.)	6 to 8 fl. oz.	8 to 16 fl. oz.	1.0%	Reduce broadcast rate in proportion to the band area actually treated.
	See COBRA® label for rates to control broadleaf weeds and height limitations for cotton. Refer to the CLEANSE 2 EC label for weed height and species controlled.			
	See MSMA label for rates to control broadleaf weeds and height limitations for cotton. Refer to the CLEANSE 2 EC label for weed height and species controlled.			

(1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Do not tank mix in these situations.  
 (2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.  
 (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.  
 (4) If at the time of application, grass height is so tall that post-directed applications cannot get good coverage over the top of the grassy weeds, then poor control may result and a second (non-post directed) application of CLEANSE 2 EC may be necessary.

## COTTON (continued)

**Table 4. CLEANSE 2 EC TANK MIXED WITH BUCTRIL® 4 EC HERBICIDE TO CONTROL EMERGED WEEDS IN BXN COTTON AS A BROADCAST APPLICATION**

PRODUCT <sup>(1)</sup>	APPLICATION RATE/ACRE <sup>(2)</sup>	CROP OIL CONCENTRATE PER ACRE <sup>(3)</sup>	COMMENTS <sup>(7)</sup>
	ANNUAL GRASSES		
CLEANSE 2 EC + BUCTRIL® 4 EC HERBICIDE <sup>(4,5,6)</sup>	8 to 16 fl. oz. See BUCTRIL® 4 EC HERBICIDE label for rates to control broadleaf weeds and height limitations for cotton.	1 qt.	See charts for grasses controlled.
<p>(1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage.</p> <p>(2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC at the specified rate with the appropriate amount of crop oil concentrate in a non-BUCTRIL® tank mix.</p> <p>(3) Always add a crop oil concentrate at 1 qt./A by ground in the finished spray solution.</p> <p>(4) Applications of BUCTRIL® 4 EC can be made only to cotton that has been genetically modified for crop tolerance to postemergence over-the-top applications of bromoxynil.</p> <p>(5) Do not apply the CLEANSE 2 EC plus BUCTRIL® tank mix within 75 days of harvest.</p> <p>(6) Do not exceed 2 applications of BUCTRIL® before cotton is 12 inches tall and one application after cotton is 12 inches tall.</p> <p>(7) Use a minimum of 10 gals. of spray solution per acre.</p>			

**Table 5. CLEANSE 2 EC TANK MIXED WITH GLYPHOSATE TO CONTROL EMERGED GRASSES IN COTTON AS A BROADCAST APPLICATION**

PRODUCT	APPLICATION RATE/ACRE <sup>(1)</sup>		ADJUVANT		COMMENTS
	ANNUAL GRASSES	PERENNIAL GRASSES	Glyphosate formulation <b>with</b> built in adjuvant	Glyphosate formulation <b>without</b> built in adjuvant	
CLEANSE 2 EC + GLYPHOSATE	6 to 8 fl. oz. See glyphosate label for rates to control broadleaf weeds and height limitations for cotton.	8 to 16 fl. oz.	Non-ionic surfactant @ 0.125 to 0.25% v/v plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals. of carrier	Crop oil concentrate @ 1 pt./A plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals. of carrier	See charts for grasses controlled. Use a minimum of 10 gals. of spray solution per acre.
<p>(1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC at the specified rate with the appropriate amount of crop oil concentrate.</p>					

## BEANS (DRY AND SUCCULENT SHELLLED)

**Table 6. CLEANSE 2 EC TANK MIXES WITH BROADLEAF HERBICIDES FOR BEANS (DRY AND SUCCULENT SHELLLED)**  
(Refer to the tables above for specific grasses and growth stages.)

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
CLEANSE 2 EC + BASAGRAN®	8 to 10 fl. oz. + 1.0 to 2.0 pts.	10 to 16 fl. oz. + 1 to 2 pts.	1%	1%

(1) Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Do not tank mix in these situations.  
 (2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC alone (without a tank mix herbicide), according to the appropriate size and rate specifications.  
 (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

## FLAX\*

**Table 7. REDUCED RATE CLEANSE 2 EC TANK MIXES WITH BROADLEAF HERBICIDES FOR FLAX**  
(Refer to the tables above for specific grasses and growth stages)

PRODUCT	APPLICATION RATES/ACRE		CROP OIL CONCENTRATE	
	ANNUAL GRASSES <sup>(1)</sup>	PERENNIAL GRASSES	GROUND	AIR
CLEANSE 2 EC + BRONATE ADVANCED™(2,3)	4 to 5 fl. oz. + 11.4 fl. oz.	-	1 pt.	1 pt.
CLEANSE 2 EC + BRONATE®(2,3)	4 to 5 fl. oz. + 0.9 pt.	-	1 pt.	1 pt.
CLEANSE 2 EC + BUCTRIL® (2,3)	4 to 5 fl. oz. + 1.0 pt.	-	1 pt.	1 pt.

*(continued)*

FLAX\* (continued)

PRODUCT	APPLICATION RATES/ACRE		CROP OIL CONCENTRATE	
	ANNUAL GRASSES <sup>(1)</sup>	PERENNIAL GRASSES	GROUND	AIR
CLEANSE 2 EC + RHONOX <sup>®(2,3)</sup>	4 to 5 fl. oz. + 0.25 to 0.5 pt.	-	1 pt.	1 pt.

(1) Annual grasses and sizes controlled with these tank mixtures are those that are identified in the **DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table**.

(2) Do not apply CLEANSE 2 EC tank mix during or after the bud stage or to ornamental flax or crop injury may occur.

(3) Do not apply tank mixes if temperatures are expected to exceed 85°F at (or 3 days following) application or crop injury may occur.

\*Not for use in California unless accompanied by a supplemental label.

**SOYBEAN**

**Table 8. CLEANSE 2 EC TANK MIXES<sup>(3)</sup> TO CONTROL ANNUAL GRASSES WHEN USED AS A BURNDOWN IN NO-TILL SOYBEAN**

PRODUCT	PRODUCT RATE/ ACRE <sup>(1)</sup>	GRASS HEIGHT (inches)	CROP OIL CONCENTRATE /ACRE <sup>(2)</sup>	28%N OR 32%N QTS./A OR 2.5 TO 4.0 LBS. AMS
CLEANSE 2 EC + 2,4-D ester <sup>*(3)</sup>	3 fl. oz.	Foxtail 1 to 3 Fall Panicum 1 to 3	1 qt.	1 to 2 qts. or 2.5 to 4.0 lbs. AMS
	4 fl. oz.	Foxtail 1 to 4 Fall Panicum 1 to 4	1 qt.	1 to 2 qts. or 2.5 to 4.0 lbs. AMS
	6 to 8 fl. oz. + 0.5 lb. ai	(See Grass Chart for grasses claimed.)	1 qt.	1 to 2 qts. or 2.5 to 4.0 lbs. AMS

\*2,4-D ester must not be used where drift sensitive crops may be grown.

(1) If regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC according to the appropriate size and rate directions.

(2) Always use a crop oil concentrate at the listed rate in the finished spray volume.

(3) The following products can be tank mixed with CLEANSE 2 EC plus 2,4-D ester: DUAL<sup>®</sup> MAGNUM<sup>®</sup>, PROWL<sup>®</sup>, SENCOR<sup>®</sup>, and SENCOR<sup>®</sup> plus DUAL<sup>®</sup> MAGNUM



**Table 9. CLEANSE 2 EC TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEANS**

(Refer to the tables above for specific grasses and growth stages.)

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>			
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
			GROUND	AIR
CLEANSE 2 EC + COBRA	6 to 8 fl. oz. + 12.5 fl. oz.	8 to 16 fl. oz. + 12.5 fl. oz.	0.5 to 1.0%	1.0%
CLEANSE 2 EC + BASAGRAN® 4 SL	8 to 10 fl. oz. + 1 to 2 pts.	10 to 16 fl. oz. + 1 to 2 pts.	1.0%	1.0%
CLEANSE 2 EC + Glyphosate (For use on Roundup Ready soybeans only)	6 to 8 fl. oz. + 0.75 to 3.0 lb. ai	8 to 16 fl. oz. + 0.75 to 3.0 lb. ai	0.5 to 1.0% <sup>(4)</sup>	1.0% <sup>(4)</sup>
CLEANSE 2 EC + ULTRA BLAZER®2 SL	6 to 8 fl. oz. + 1 to 1.5 pts.	6 to 8 fl. oz. + 1 to 1.5 pts.	0.5 to 1.0%	1.0%
CLEANSE 2 EC + FLEXSTAR® HL <sup>(6)</sup>	6 to 8 fl. oz. Refer to the FLEXSTAR® HL label for specific application rates.	8 to 16 fl. oz. Refer to the FLEXSTAR® HL label for specific application rates.	1.0%	1.0%
CLEANSE 2 EC + CLASSIC® 25 DG	8 to 10 fl. oz. + 0.5 to 0.75 oz.	10 to 16 fl. oz. + 0.5 to 0.75 oz.	1.0%	1.0%
CLEANSE 2 EC <sup>(5)</sup> + PURSUIT® 70 DG	6 to 8 fl. oz. + 1.44 oz.	8 to 16 fl. oz. + 1.44 oz.	1.0%	1.0%
CLEANSE 2 EC <sup>(5)</sup> + COBRA + CLASSIC® 25 DG	8 to 10 fl. oz. + 6 to 8 fl. oz. + 0.5 to 0.75 oz.	-	0.5%	1.0%

*(continued)*

**CLEANSE 2 EC TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEANS** *(continued)*

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>			
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
			GROUND	AIR
CLEANSE 2 EC <sup>(5)</sup> + COBRA + BASAGRAN® 4 SL	8 to 10 fl. oz. + 6 to 10 fl. oz. + 1 to 1.5 pts.	-	0.5%	1.0%
CLEANSE 2 EC <sup>(5)</sup> + COBRA + PURSUIT® 70 DG	8 to 10 fl. oz. + 6 to 10 fl. oz. + 1.44 oz.	-	0.5%	1.0%
CLEANSE 2 EC <sup>(5)</sup> + STORM®	8 to 10 fl. oz. + 1.5 pts.	-	0.5%	1.0%
CLEANSE 2 EC + RESOURCE® + PURSUIT® 70 DG	8 to 10 fl. oz. + 4 fl. oz. + 1.44 oz.	-	1.0%	1.0%
CLEANSE 2 EC + RESOURCE® + BASAGRAN®	8 to 10 fl. oz. + 4 fl. oz. + 1 pt.	-	1.0%	1.0%
CLEANSE 2 EC + RESOURCE® + CLASSIC®	8 to 10 fl. oz. + 4 fl. oz. + 0.5 oz.	-	1.0%	1.0%

*(continued)*

**CLEANSE 2 EC TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEANS** *(continued)*

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>			
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
			GROUND	AIR
CLEANSE 2 EC + COBRA® + RESOURCE®	6 to 8 fl. oz. + 6 fl. oz. + 4 fl. oz.	-	0.5%	1.0%
CLEANSE 2 EC <sup>(5)</sup> + FIRSTRATE®	6 to 8 fl. oz. + 0.3 oz./A	8 to 16 fl. oz. + 0.3 oz./A	1.0%	-
CLEANSE 2 EC <sup>(5)</sup> + COBRA® + FIRSTRATE®	6 to 8 fl. oz. + 6 to 8 fl. oz. + 0.3 oz./A	8 to 16 fl. oz. + 6 to 8 fl. oz. + 0.3 oz./A	1.0%	-
CLEANSE 2 EC <sup>(5)</sup> + RAPTOR® (1 AS)	6 to 8 fl. oz. + 4 to 5 fl. oz.	-	1.0%	-
CLEANSE 2 EC <sup>(5)</sup> + COBRA® + RAPTOR® (1 AS)	6 to 8 fl. oz. + 6 to 8 fl. oz. + 4 to 5 fl. oz.	-	1.0%	-
CLEANSE 2 EC <sup>(5)</sup> + SYNCHRONY® STS™	6 to 8 fl. oz. <sup>(7)</sup> + 0.5 oz. /A	-	1 qt.	-
CLEANSE 2 EC <sup>(5)</sup> + COBRA® + SYNCHRONY® STS™	6 to 8 fl. oz. <sup>(7)</sup> + 4 to 8 fl. oz. + 0.5 oz./A	-	1 pt.	-

*(continued)*

**CLEANSE 2 EC TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEANS** *(continued)*

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>			
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
			GROUND	AIR
CLEANSE 2 EC <sup>(5)</sup> + RESOURCE <sup>®</sup>	6 to 8 fl. oz. + 4 to 12 fl. oz.	-	1 qt.	-
CLEANSE 2 EC <sup>(5)</sup> + FRONTROW <sup>™</sup>	8 to 10 fl. oz. + Refer to FRONTROW <sup>™</sup> label for use rates.	-	1.0%	-
CLETHODIM 2 EC HERBICIDE + FIRSTRATE + FLEXSTAR HL <sup>®</sup>	6 to 8 fl. oz. + 0.3 oz. + Refer to the FLEXSTAR <sup>®</sup> HL label for specific application rates.	8 to 16 fl. oz. + 0.3 oz. + Refer to the FLEXSTAR <sup>®</sup> HL label for specific application rates.	1.0%	-

- (1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Do not tank mix in these situations.
- (2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC alone (without a tank mix herbicide), according to the appropriate size and rate directions.
- (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- (4) The addition of 2.5 lbs. of ammonium sulfate is required when CLEANSE 2 EC is tank mixed with glyphosate. If the glyphosate formulation has a stand alone built in adjuvant, add 0.125% v/v non-ionic surfactant in place of crop oil concentrate. If the glyphosate formulation does not have a built in adjuvant system, add 0.5 to 1.0% crop oil concentrate for ground application and 1.0% v/v for aerial application.
- (5) Add 1 to 2 qts./A of liquid fertilizer (10-34-0 or 32%N) when CLEANSE 2 EC is tank mixed with PURSUIT<sup>®</sup>, RESOURCE<sup>®</sup>, GALAXY<sup>™</sup> STORM<sup>®</sup>, FIRSTRATE<sup>®</sup>, SYNCHRONY<sup>®</sup>, RAPTOR<sup>®</sup>, FRONTROW<sup>™</sup>, COBRA<sup>®</sup> plus CLASSIC<sup>®</sup>, COBRA<sup>®</sup> plus BASAGRAN<sup>®</sup>, COBRA<sup>®</sup> plus PURSUIT<sup>®</sup>, COBRA<sup>®</sup> plus FIRSTRATE<sup>®</sup>, COBRA<sup>®</sup> plus SYNCHRONY<sup>®</sup>, and COBRA<sup>®</sup> plus RAPTOR<sup>®</sup>. An equivalent amount (2.5 to 4.0 lbs./A) of adjuvants are to be added in addition to the crop oil concentrate.
- (6) Refer to FLEXSTAR<sup>®</sup> HL label for geographic and rotational restrictions.
- (7) Annual grasses and sizes controlled with these tank mixtures are those that are identified in the **DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES** table.

**Table 10. REDUCED RATE CLEANSE 2 EC TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN** (Refer to table for reduced rate use in canola, dry bean, flax, mustard seed, soybean, and sugar beet directions for small annual grasses for specific grasses and growth stages)

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>			
	ANNUAL GRASSES <sup>(2)</sup>	PERENNIAL GRASSES	CROP OIL CONCENTRATE <sup>(3,4)</sup> (V/V)	
			GROUND	AIR
CLEANSE 2 EC + FIRSTRATE®	4 to 8 fl. oz. + 0.3 oz.	-	1.0%	1.0%
CLEANSE 2 EC + PURSUIT® 70 DG	4 to 6 fl. oz. + 1.44 oz.	-	1.0%	1.0%

(1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC alone (without a tank mix herbicide), according to the appropriate size and rate directions.  
(2) Annual grasses and sizes controlled with these tank mixtures are those that are identified in the **DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES** table.  
(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.  
(4) The addition of 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N) is required when CLEANSE 2 EC is tank mixed at reduced rates. An equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer. Fertilizer adjuvants are to be added in addition to the crop oil concentrate.

**Table 11. USE DIRECTIONS FOR ROUNDUP READY VOLUNTEER CORN CONTROL IN ROUNDUP READY SOYBEANS WITH CLEANSE 2 EC TANK MIX**

Roundup Ready Volunteer Corn Height (inches)	CLEANSE 2 EC Rate fl. oz./A	Glyphosate <sup>(1)</sup> rate for formulations with built in adjuvant	Adjuvant
<12	4	1.0 to 2.0 lbs. ai./A (Approximately equivalent to 22 to 44 fl. oz./A of ROUNDUP weather MAX)	Non-ionic surfactant @ 0.125 to 0.25% v/v plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals. of carrier
12 to 18	5		
18 to 24	6		

  

Roundup Ready Volunteer Corn Height (inches)	CLEANSE 2 EC Rate fl. oz./A	Glyphosate <sup>(1)</sup> rate for formulations without built in adjuvant	Adjuvant
<12	4	Up to 2.0 lbs. ai./A (Equivalent to 32 to 64 fl. oz./A of ROUNDUP Original)	Crop oil concentrate @ 1 pt./A plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals. of carrier
12 to 18	5		
18 to 24	6		

(1) Glyphosate formulation must be labeled for use on Roundup Ready soybeans.

## PEANUT (INCLUDING PERENNIAL)

**Table 12. CLEANSE 2 EC TANK MIXES WITH BROADLEAFHERBICIDES FOR PEANUT (INCLUDING PERENNIAL)**  
(Refer to the tables above for specific grasses and growth stages.)

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>			
	ANNUAL GRASSES <sup>(2)</sup>	PERENNIAL GRASSES	CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
			GROUND	AIR
CLEANSE 2 EC + BASAGRAN®	8 to 10 fl. oz. + 1.0 to 2.0 pts.	-	1.0%	1.0%
CLEANSE 2 EC + ULTRA BLAZER	8 to 10 fl. oz. + 0.5 to 1.5 pts.	-	1.0%	1.0%
CLEANSE 2 EC + STORM®	8 to 10 fl. oz. + 1.5 pts.	-	1.0%	1.0%

(1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Do not tank mix in these situations.  
 (2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC alone (without a tank mix herbicide), according to the appropriate size and rate specifications.  
 (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

### GRASS SUPPRESSION FOR HARVEST EFFICIENCY IN PEANUT (INCLUDING PERENNIAL) WITH CLEANSE 2 EC

GRASS SPECIES	WEED STAGE	RATE FL. OZ./ ACRE	HIGH RATE
Annual and perennial grasses that exceed height claimed for control on height charts "ANNUAL GRASSES" & "PERENNIAL GRASSES"	Up to and including grasses in the seed head stage.	16	32
<ul style="list-style-type: none"> <li>• Do not apply as part of a tank mix when applying CLEANSE 2 EC for grass suppression.</li> <li>• Add a crop oil concentrate at 1 qt./A by ground to the finished spray volume.</li> </ul>			

## SUGAR BEET

**Table 13. CLEANSE 2 EC TANK MIXED WITH STINGER® APPLIED TO SUGAR BEETS**

(Refer to the tables above for specific grasses and growth stages)

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	Annual Grasses	Perennial Grasses	Ground	Air
CLEANSE 2 EC + STINGER	6 to 8 fl. oz.	8 to 16 fl. oz.	1.0%	1.0%
See STINGER label for rates. Refer to the CLEANSE 2 EC label for weed height and species controlled.				

(1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Do not tank mix in these situations.  
 (2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC alone (without a tank mix herbicide), according to the appropriate size and rate directions.  
 (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

**Table 14. CLEANSE 2 EC TANK MIXED WITH BETAMIX® or BETANEX® APPLIED TO SUGAR BEETS**

PRODUCT <sup>(1)</sup>	WEEDS CONTROLLED		WEED HEIGHT (inches)	APPLICATION RATE/ACRE <sup>(2)</sup>
	COMMON NAME	SCIENTIFIC NAME		
CLEANSE 2 EC ® + BETAMIX® or BETANEX®	Barnyardgrass	<i>Echinochloa crus-galli</i>	1 to 3	8 fl. oz.
	Foxtail	<i>Setaria</i> spp.	1 to 3	
	Foxtail Millet	<i>Setaria italica</i>	1 to 3	
	Wild Oats	<i>Avena fatua</i>	1 to 3	
	Wild Proso Millet	<i>Panicum miliaceum</i>	See <b>BETAMIX® and BETANEX®</b> labels for rates to control broadleaf weeds. <b>Do not use additives in the tank mix.</b>	

(1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Do not tank mix in these situations.  
 (2) Do not use crop oil concentrate. Do not use additives in the tank mix.  
 (3) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC at full label rate with appropriate rate of crop oil concentrate.

**TABLE 15. CLEANSE 2 EC PLUS BETANEX® OR BETAMIX® TANK MIX FOR THREE SEQUENTIAL APPLICATIONS FOR ANNUAL GRASS CONTROL (MICRO-RATE APPLICATION)**

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>		METHYLATED SEED OIL <sup>(2)</sup> (V/V)	
	ANNUAL GRASSES	GRASSES CONTROLLED (inches)	GROUND	AIR
CLEANSE 2 EC + BETANEX® or BETAMIX®	2 to 3 fl. oz. + 8 to 12 fl. oz. <sup>(3)</sup> or 8 to 12 fl. oz. <sup>(3)</sup>	Green Foxtail (1-2) Yellow Foxtail (1-2) Barnyardgrass (1-2) Wild Oat (1-2) Volunteer Cereals (1-2)	1.5%	1.5%

(1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Do not tank mix in these situations.  
 (2) Always use a methylated seed oil at the listed rate (but not less than 1 pt./A) in the finished spray volume.  
 (3) Use 8 fl. oz./A rate when sugar beet are in the cotyledon to 4 leaf stage. Rate can be increased up to 12 fl. oz./A when the smallest sugar beet plants in the field are in the 4 true leaf stage or larger.

**Micro-Rate Applications to Sugar Beets**

Multiple micro-rate applications of CLEANSE 2 EC in tank mixtures with reduced rates of BETANEX® or BETAMIX® and methylated seed oils may be applied by air or ground equipment to sugar beets to control early germinating annual grasses listed above. The rate of BETANEX® or BETAMIX® must not exceed 0.12 lb. ai./A (broadcast application) when in combination with these spray adjuvants. Note that maximum rate allowed varies depending on crop growth stage. The use of wetting agents or spray adjuvants with conventional rates (0.73 to 1.22 lb. ai./A) or multiple low rate (0.24 to 0.73 lb. ai./A) applications of BETANEX® or BETAMIX® is prohibited on the BETANEX® and BETAMIX® master label. Favorable climatic conditions (good conditions for plant growth and development) are essential for adequate weed control. All use precautions and restrictions on the BETANEX® and BETAMIX® master labels must be followed.

**Micro-Rate Multiple Applications of CLEANSE 2 EC Tank Mixes**

Apply CLEANSE 2 EC in broadcast applications only at a rate of 2 to 3 fl. oz./A in tank mixtures with either BETANEX® or BETAMIX® following the directions for use on the tank mix partner label. A minimum of three sequential applications of 2 fl. oz./A or a minimum of 2 sequential applications of 3 fl. oz./A must be utilized for CLEANSE 2 EC tank mixtures. A minimum of 3 sequential applications of BETANEX® or BETAMIX® must be used. Accurate timing is essential; make initial application immediately after weeds emerge, and make repeat applications on 5 to 7 day intervals. If weed control is not adequate due to climatic conditions, spray coverage or other factors, return to conventional application rates of CLEANSE 2 EC (6 to 8 fl. oz./A) and add rates of BETANEX® or BETAMIX® as directed on their label. When using conventional rates of BETANEX® or BETAMIX® in tank mixtures with CLEANSE 2 EC, do not use a spray adjuvant.

**Use Precautions for Micro-Rate Applications: (See CLEANSE 2 EC, BETANEX® and BETAMIX® master label for further use precautions.)**

Not all weeds will be adequately controlled, even with favorable climatic conditions. Conventional rates of CLEANSE 2 EC, BETANEX® or BETAMIX® and/or hand labor may be required if multiple micro-rate applications do not adequately control weeds. Plugging of spray nozzles may be encountered due to the potential for formation of a precipitate in the spray solution that is often associated with



micro-rate applications. Direct Ag Source, LLC will not be responsible for any nozzle plugging that may occur with the use of multiple micro-rate applications. Methylated seed oils must not be added if the BETANEX® or BETAMIX® rate exceeds 0.12 lb. ai./A broadcast, as the addition of methylated seed oils could increase the possibility of crop injury at dosage rates greater than 0.12 lb. ai./A.

#### GROUND APPLICATION

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 10 gals. and a maximum of 20 gals. of spray solution per acre. Spray pressures must reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

#### AERIAL APPLICATION

Use of sufficient spray volumes is essential to ensure complete coverage. Use a minimum of 5 gals. and a maximum of 15 gals. of spray solution per acre.

**Table 16. TANK MIX APPLICATION OF CLEANSE 2 EC AND FUNGICIDES FOR CONTROL OF GRASS WEEDS AND DISEASES IN SUGAR BEET**

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>		
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)
CLEANSE 2 EC + EMINENT®	6 to 8 fl. oz. + 13 fl. oz.	8 to 16 fl. oz. + 13 fl. oz.	1.0%

(1) Refer to CLEANSE 2 EC and fungicide label for rates and weeds and diseases controlled.  
 (2) If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC alone (without a tank mix fungicide) according to the appropriate size and rate specifications.  
 (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

**Table 17. TANK MIX APPLICATION OF CLEANSE 2 EC AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN ALFALFA, COTTON, MINT, PEANUT (INCLUDING PERENNIAL), SOYBEANS AND SUNFLOWER**

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>			CROP					
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONCENTRATE (V/V) <sup>(3)</sup>	Alfalfa <sup>4</sup>	Cotton	Mint <sup>4,5</sup>	Peanut	Soybean	Sunflower
CLEANSE 2 EC + ORTHENE® 75 S or ORTHENE 97	6 to 8 fl. oz. + 0.33 to 1.33 lbs. + 0.25 to 1.0 lb.	8 to 16 fl. oz. + 0.33 to 1.33 lbs. + 0.25 to 1.0 lb.	1.0%		X	X	X		
CLEANSE 2 EC + ORTHENE® 90 S <sup>(6)</sup>	6 to 8 fl. oz. + 0.25 to 1 lb.	8 to 16 fl. oz. + 0.25 to 1 lb.	1.0%		X	X	X	X	
CLEANSE 2 EC + DANITOL® 2.4 EC	6 to 8 fl. oz. + 10 2/3 to 16 fl. oz.	8 to 16 fl. oz. + 10 2/3 to 16 fl. oz.	1.0%		X		X		
CLEANSE 2 EC + ASANA XL®	6 to 8 fl. oz. + Refer to ASANA XL® label	8 to 16 fl. oz. + Refer to ASANA XL® label	1.0%						X
CLEANSE 2 EC + WARRIOR®	6 to 8 fl. oz. + Refer to WARRIOR® label	8 to 16 fl. oz. + Refer to WARRIOR® label	1.0%						X
CLEANSE 2 EC + WARRIOR®	10 to 16 fl. oz. <sup>(7)</sup> + Refer to WARRIOR® label	10 to 16 fl. oz. + Refer to WARRIOR® label	1.0%	X					
CLEANSE 2 EC + BAYTHROID®	10 to 16 fl. oz. <sup>(7)</sup> + Refer to BAYTHROID® label	10 to 16 fl. oz. + Refer to BAYTHROID® label	1.0%	X					

(continued)

**Table 17. TANK MIX APPLICATION OF CLEANSE 2 EC AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN ALFALFA, COTTON, MINT, PEANUT (INCLUDING PERENNIAL), SOYBEANS AND SUNFLOWER (continued)**

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>			CROP					
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONCENTRATE (V/V) <sup>(3)</sup>	Alfalfa <sup>(4)</sup>	Cotton <sup>(4,5)</sup>	Mint <sup>(4,5)</sup>	Peanut	Soybean	Sunflower
CLEANSE 2 EC + DIMETHOATE®	10 to 16 fl. oz. <sup>(7)</sup> + Refer to DIMETHOATE® label	10 to 16 fl. oz. + Refer to DIMETHOATE® label	1.0%	X					
CLEANSE 2 EC + LORSBAN®	10 to 16 fl. oz. <sup>(7)</sup> + Refer to LORSBAN® label	10 to 16 fl. oz. + Refer to LORSBAN® label	1 to 2 pt. <sup>(8)</sup>	X					
CLEANSE 2 EC + POUNCE®	10 to 16 fl. oz. <sup>(7)</sup> + Refer to POUNCE® label	10 to 16 fl. oz. + Refer to POUNCE® label	1.0%	X					

(1) Refer to CLEANSE 2 EC and insecticide label for rates and weeds and insects controlled.  
 (2) If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC alone (without a tank mix insecticide) according to the appropriate size and rate specifications.  
 (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.  
 (4) Certain insecticides may cause temporary phytotoxic symptoms on alfalfa and mint foliage.  
 (5) The CLEANSE 2 EC rate must be 6 to 8 fl. oz./A for annual grass control in baby mint, minimum of 8 fl. oz./A for annual grass control in established mint and 8 to 16 fl. oz./A for perennial grass control. Add crop oil concentrate at the rate of 1.0 to 2.0 pts./A.  
 (6) Insecticide tank mix use with ORTHENE® 90 S in soybean is permitted only in a state having an approved Section 24(c) registration for ORTHENE® 90 S use in soybeans.  
 (7) The CLEANSE 2 EC rate must be 6 to 8 fl. oz./A for annual grass control in seedling alfalfa.  
 (8) For the CLEANSE 2 EC plus LORSBAN tank mix, reduce the adjuvant rate down to 1.0 pt./A when the LORSBAN rate is 1.0 pt./A or higher.

**THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.**

- Apply only to actively growing grass and broadleaf weeds at specified height or growth stage listed on each label.
- Apply under favorable soil moisture and humidity which exist a few days after rainfall or within seven days after irrigation.
- Tank mix applications may sometimes result in reduced grass control. If regrowth occurs, or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC, as specified in the respective size and rate tables.
- Do not tank mix CLEANSE 2 EC when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.
- This tank mix may be applied postemergence to ROUNDUP READY soybean up through the full flowering stage. Do not apply less than 60 days before harvest.

- Avoid contact with foliage, green stems, or fruit crops, or any desirable plants and trees, other than soybean with the ROUNDUP READY gene as severe injury or destruction will result.
- Do not allow the CLEANSE 2 EC plus ROUNDUP to mist, drip, drift or splash onto desirable vegetation as minute quantities of the tank mix can cause severe damage or destruction to the crops, plants or other areas on which treatment was not intended. The likelihood of injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions that allow spray drift to occur such as combinations of spray pressure and nozzle type that will result in fine particles (mist) that are likely to drift.

### FALLOW LAND

CLEANSE 2 EC may be used to control annual and perennial grasses in land that has been left fallow the previous year and other non-producing agricultural areas. Apply CLEANSE 2 EC at 6 to 8 fl. oz./A for annual grasses and 8 to 16 fl. oz./A for perennial grasses. When both grass and broadleaf weeds are the target pest, CLEANSE 2 EC may be tank mixed with 2,4-D ester or Dicamba for broad spectrum control. When both annual and perennial grasses occur in the same field, use a minimum of 8 fl. oz./A. CLEANSE 2 EC rate.

- Use a minimum spray volume of 5 gals./A for aerial applications and 15 gals./A for ground applications.
- Apply only to actively growing grasses when the first grass reaches the specified weed height as specified in the Annual and Perennial Grasses section of this label.
- Annual grasses which emerge after the CLEANSE 2 EC application will not be controlled, and a second application may be necessary.
- The control of perennial grasses may require more than 1 application in non-tilled areas.
- Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop.
- Do not apply to grasses that have tillered, formed seedheads or exceeded specified growth stage.
- Do not use flood jet nozzles.
- Do not apply to drought stressed grasses.
- Do not mow area for 2 weeks prior to or after the CLEANSE 2 EC application.

**TABLE 18. CLEANSE 2 EC IN TANK MIXES TO CONTROL ANNUAL AND PERENNIAL GRASSES IN FALLOW LAND**

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(2)</sup> (V/V)	
	Annual Grasses	Perennial Grasses	Ground	Air
CLEANSE 2 EC + 2,4-D ester or Dicamba	6 to 8 fl. oz.	8 to 16 fl. oz.	1.0%	1.0%
	0.5 lb./A			
	See Dicamba label for rates.			
<p>(1) Refer to CLEANSE 2 EC label for weed height and species control. Review Dicamba and 2,4-D labels for crop restrictions, use rates and weeds controlled.</p> <p>(2) Always use a crop oil concentrate or methylated seed oil containing at least 15% emulsifier at the listed rate (but not less than 1 pt./A) in the finished spray volume.</p>				

GRASS SUPPRESSION IN NON-CROP AREAS WITH CLETHODIM 2 EC HERBICIDE			
GRASS SPECIES	WEED STAGE	RATE FL. OZ./ ACRE	HIGH RATE
Annual and perennial grasses that exceed height claimed for control on height chart above.	Up to and including grasses in the seed head stage.	12	16
<ul style="list-style-type: none"> <li>Do not apply as part of a tank mix when applying CLEANSE 2 EC for grass suppression.</li> <li>Add a crop oil concentrate at 1 qt./A by ground to the finished spray volume.</li> </ul>			

**TABLE 19. CLEANSE 2 EC FOR THE CONTROL AND/OR SUPPRESSION OF TALL FESCUE IN NATIVE PRAIRIE WARM-SEASON GRASS RESTORATION PROJECTS**

PRODUCT	PRODUCT RATES	GRASS WEEDS CONTROLLED/SUPPRESSED		WEED STAGES
		Common Name	Scientific Name	
CLEANSE 2 EC	10 to 12 fl. oz./A	Tall Fescue	<i>Festuca arundinacea</i>	4 to 6 inches tall (40 to 60% green-up)
<b>Adjuvant:</b> CLEANSE 2 EC must be applied with crop oil concentrate at 1 qt./A, plus a spray grade ammonium sulfate at 2.5 to 4 lbs./A.				
<b>Mixing Order:</b> Thoroughly mix spray grade ammonium sulfate in water, add CLEANSE 2 EC, then add crop oil concentrate.				

**SPECIAL APPLICATION INSTRUCTIONS/PRECAUTIONS:**

- Burn or mow fields a minimum of 3 weeks prior to application to remove excess crop residue. Apply in the spring, at 40 to 60% green-up, prior to emergence of warm-season grasses. Do not mow area for 2 weeks after the CLEANSE 2 EC application.
- Apply in a minimum of 15 to 20 gals. of water per acre at a spray pressure of 40 to 60 psi at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood nozzles.
- Apply only to fields that have warm-season grasses established for 2 years. Applications of CLEANSE 2 EC to emerged warm-season grasses may cause injury. Do not apply to warm-season grasses grown for seed.
- Do not graze treated fields or feed treated forage and or hay to livestock. Do not plant any crop for 30 days after application, unless clethodim is registered for use in that crop.

**NOTE:** CLEANSE 2 EC applications are most effective if applied when average nighttime temperatures are consistently greater than or equal to 47°F.

**TABLE 20. CLEANSE 2 EC FOR THE SUPPRESSION OF TALL FESCUE SEED-HEADS IN NON-PRODUCING AGRICULTURAL AREAS**

Product	Product Rate	Suppression	Application Timing
CLEANSE 2 EC	1 1/2 to 2 fl. oz./A	Tall Fescue Seed-Heads ( <i>Festuca arundinacea</i> )	50 to 90% Tall Fescue green-up
<b>ADJUVANT:</b> CLEANSE 2 EC must be applied with crop oil concentrate at 1 qt./A, plus a spray grade ammonium sulfate at 2.5 to 4 lbs./A. <b>Mixing Order:</b> Thoroughly mix spray grade ammonium sulfate in water, add CLEANSE 2 EC then add crop oil concentrate.			

**SPECIAL APPLICATION INSTRUCTIONS/PRECAUTIONS:**

- Apply at 50 to 90% tall fescue green-up.
- Use the higher CLEANSE 2 EC rate if less tall fescue green matter is present.
- Do not mow area for two weeks after the CLEANSE 2 EC application.
- Apply in a minimum of 15 to 20 gals. of water per acre at a spray pressure of 40 to 60 psi at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood nozzles.
- 2-4-D ester may be added to this tank mix for broadleaf control (see 2,4-D ester label for weeds controlled).
- Do not graze treated fields or feed treated forage and/or hay to livestock. Do not plant any crop for 30 days after application, unless clethodim is registered for use in that crop.

**THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.**

- Apply only to actively growing grass and broadleaf weeds at specified height or growth stage listed on each label.
- Apply under favorable soil moisture and humidity which exist a few days after rainfall or within seven days after irrigation.
- Tank mix applications may sometimes result in reduced grass control. If regrowth occurs, or an additional flush of new grass emerges, make a second application of CLEANSE 2 EC, as specified in the respective size and rate tables.
- Do not tank mix CLEANSE 2 EC when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.
- This tank mix may be applied postemergence to ROUNDUP READY soybeans up through the full flowering stage. Do not apply less than 60 days before harvest.
- Avoid contact with foliage, green stems, or fruit crops, or any desirable plant and trees, other than soybeans with the ROUNDUP READY gene as severe injury or destruction will result.
- Do not allow the CLEANSE 2 EC plus ROUNDUP to mist, drip, drift or splash onto desirable vegetation as minute quantities of the tank mix can cause severe damage or destruction to the crops, plants or other areas on which treatment was not intended. The likelihood of injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions that allow spray drift to occur such as combinations of spray pressure and nozzle type that will result in fine particles (mist) that are likely to drift.

## ORNAMENTALS

For ornamental plant uses, CLEANSE 2 EC can be used to control labeled grass weeds in greenhouses, lathhouses, shadehouses, and around outdoor ornamentals, including nurseries, parks, roadside plantings, and structure landscapes.

### IMPORTANT

CLEANSE 2 EC successfully controls weeds in newly transplanted and established non-grassy ornamentals. Plant tolerance to CLEANSE 2 EC at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, the user must determine if herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of CLEANSE 2 EC have investigated the safety factor to ornamental plants not listed on the label.

The following plants have shown a tolerance for CLEANSE 2 EC applications:

### ORNAMENTAL TREES

COMMON NAME	SCIENTIFIC NAME
Alder, Red	<i>Alnus oregona</i>
Ash	<i>Fraxinus</i> spp.
Basswood	<i>Tilia</i> spp.
Birch, European White	<i>Betula pendula</i>
Birch, River	<i>Betula nigra</i>
Birch, White	<i>Betula papyrifera</i>
Crabapple, Flowering	<i>Malus halliana</i>
Dogwood, Flowering	<i>Cornus, florida</i>
Golden Chain Tree	<i>Laburnum anagyroides</i>
Maples	<i>Acer</i> spp.
Mulberry, White	<i>Morus alba</i>
Oaks	<i>Quercus</i> spp.
Olive, Wild	<i>Elaeagnus angustifolia</i>
Redbud	<i>Cercis Canadensis</i>
Sweet Gum, American	<i>Liquidambar styraciflua</i>

## GROUND COVERS

COMMON NAME	SCIENTIFIC NAME
Bugleweed, Carpet	<i>Ajuga reptans</i>
Ivy, English	<i>Hedera helix</i>
Japanese Spurge	<i>Pachysandra terminalis</i>
Lilyturf	<i>Liriope muscari</i>
Moneywort	<i>Lysimachia nummularia</i>
Mondo Grass, White	<i>Ophiopogon jaburan</i>
Mondo Grass Dwarf	<i>Ophiopogon japonicus</i>
Periwinkle, Common	<i>Vinca minor</i>

## GARDEN FLOWERS AND PLANTS

COMMON NAME	SCIENTIFIC NAME
Ageratum	<i>Ageratum</i> spp.
Alyssum*, Sweet	<i>Lobularia maritima</i>
Asparagus Fern	<i>Asparagus setaceus</i>
Bleeding Heart	<i>Dicentra spectabilis</i>
Cast Iron Plant	<i>Aspidistra elatior</i>
Chrysanthemum	<i>Chrysanthemum</i> spp.
Cinquefoil	<i>Potentilla</i> spp.
Coleus	<i>Coleus</i> spp.
Coralbells	<i>Heuchera sanguinea</i>
Cranesbill	<i>Geranium</i> spp.
Dahlia	<i>Dahlia</i> spp.
Daisy, Trailing African	<i>Osteospermum fruticosum</i>
Daylily	<i>Hemerocallis</i> spp.
Dusty Miller	<i>Senecio cineraria</i>
Euonymus	<i>Euonymus</i> spp.
Gazania	<i>Gazania</i> spp.
Geranium, House	<i>Pelargonium hortorum</i>

(continued)



**GARDEN FLOWERS AND PLANTS** (continued)

COMMON NAME	SCIENTIFIC NAME
Heather, False	<i>Cuphea hyssopifolia</i>
Hosta	<i>Hosta fortunei</i>
Iris	<i>Iris</i> spp.
Jasmine Tobacco	<i>Nicotiana alata</i>
Loosestrife	<i>Lythrum salicaria</i>
Marigold	<i>Tagetes</i> spp.
Partridgeberry	<i>Mitchella repens</i>
Petunia*	<i>Petunia hybrida</i>
Phlox	<i>Phlox</i> spp.
Pinks	<i>Dianthus</i> spp.
Portulaca	<i>Portulaca grandiflora</i>
Salvia	<i>Salvia</i> spp.
Saxifrage	<i>Saxifraga</i> spp.
Sedum	<i>Sedum</i> spp.
Selloum	<i>Philodendron selloum</i>
Snapdragon*	<i>Antirrhinum majus</i>
Sweet Flag	<i>Acorus gramineus</i>
Tickseed	<i>Coreopsis grandiflora</i>
Touch-Me-Not	<i>Impatiens</i> spp.
Verbena	<i>Verbena</i> spp.
Violet	<i>Viola</i> spp.
Yarrow, Common	<i>Achillea millefolium</i>
Zinnia	<i>Zinnia elegans</i>
*Slight foliage or flower speckling has been observed on these species.	

## SHRUBS

COMMON NAME	SCIENTIFIC NAME
Abelia	<i>Abelia</i> spp.
Anise, Purple	<i>Illicium floridanum</i>
Aucuba	<i>Aucuba</i> spp.
Azalea*	<i>Rhododendron</i> spp.
Bamboo	<i>Bambusa</i> spp.
Barberry, Japanese	<i>Berberis thunbergii</i>
Barberry, Magellan	<i>Berberis buxifolia</i>
Bayberry	<i>Myrica pensylvanica</i>
Bottlebrush	<i>Callistemon citrinus</i>
Boxwood, Common	<i>Buxus sempervirens</i>
Camellia, Common	<i>Camellia japonica</i>
Candytuft	<i>Iberis sempervirens</i>
Cleyera	<i>Cleyera japonica</i>
Coralberry	<i>Ardisia crenata</i>
Crape Myrtle	<i>Lagerstroemia indica</i>
Coyote Brush	<i>Baccharis pilularis</i>
Fig, Creeping	<i>Ficus pumila</i>
Gardenia	<i>Gardenia</i> spp.
Holly	<i>Ilex</i> spp.
Honeysuckle	<i>Lonicera pileate</i>
Indian Hawthorn	<i>Raphiolepis indica</i>
Jasmine	<i>Jasminum</i> spp.
Jasmine, Asiatic	<i>Trachelospermum asiaticum</i>
Jasmine, Star	<i>Trachelospermum jasminoides</i>
Juniper	<i>Juniperus</i> spp.
Lantana	<i>Lantana</i> spp.
Nandina* Bamboo Heavenly	<i>Nandinia domestica</i>
Oleander, Common	<i>Nerium oleander</i>

(continued)

**SHRUBS** (continued)

COMMON NAME	SCIENTIFIC NAME
Oregon Grape	<i>Mahonia aquifolium</i>
Photinia	<i>Photinia</i> spp.
Pittosporum	<i>Pittosporum</i> spp.
Podocarpus	<i>Podocarpus</i> spp.
Privet	<i>Ligustrum</i> spp.
Pyracantha	<i>Pyracantha</i> spp.
Rhododendron	<i>Rhododendron</i> spp.
Rose	<i>Spiraea bumalda</i>
Sweet Olive	<i>Osmanthus fragrans</i>
Viburnum	<i>Viburnum tinus</i>
Wisteria	<i>Wisteria</i> spp.
Yellow Sage/Shrub Verbena	<i>Lantana camara</i>
*Slight foliage or flower speckling has been observed on these species.	

**ANNUAL GRASSES IN ORNAMENTALS**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

GRASS SPECIES	SCIENTIFIC NAME	WEED* HEIGHT (INCHES)	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
Barnyardgrass	<i>Echinochloa crus-galli</i>	2 to 8	8	16
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	2 to 6	8	16
Brome				
California	<i>Bromus carinatus</i>	2 to 6	8	16
Cheatgrass	<i>Bromus secalinus</i>	2 to 6	8	16
Downy	<i>Bromus tectorum</i>	2 to 6	8	16
Ripgut	<i>Bromus diandrus</i>	2 to 6	8	16
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	8	16
Crabgrass				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	8	16

(continued)

**ANNUAL GRASSES IN ORNAMENTALS** *(continued)*

GRASS SPECIES	SCIENTIFIC NAME	WEED* HEIGHT (INCHES)	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
Large	<i>Digitaria sanguinalis</i>	2 to 6**	8	16
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	8	16
Southern	<i>Digitaria ciliaris</i>	2 to 6**	8	16
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	8	16
Fall Panicum	<i>Panicum dichotomiflorum</i>	2 to 8	8	16
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	8	16
Foxtail				
Giant	<i>Setaria faberi</i>	2 to 12	8	16
Green	<i>Setaria viridis</i>	2 to 8	8	16
Yellow	<i>Setaria glauca</i>	2 to 8	8	16
Foxtail Barley	<i>Hordeum jubatum</i>	2 to 6	8	16
Goosegrass	<i>Eleusine indica</i>	2 to 6**	8	16
Itchgrass	<i>Rottboellia exaltata</i>	2 to 6	8	16
Junglerice	<i>Echinochloa colona</i>	2 to 6	8	16
Lovegrass (Stinkgrass)	<i>Eragrostis ciliaris</i>	2 to 6	8	16
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	8	16
Red Rice	<i>Oryza sativa</i>	1 to 3	8	16
Ryegrass				
Hardy	<i>Lolium remotum</i>	2 to 6	8	16
Italian	<i>Lolium multiflorum</i>	2 to 6	8	16
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	8	16
Shattercane	<i>Sorghum bicolor</i>	6 to 18	8	16
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	8	16
Sprangletop				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	8	16
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	8	16
Mexican	<i>Leptochloa uninervia</i>	2 to 6	8	16
Red	<i>Leptochloa filiformis</i>	2 to 6	8	16

*(continued)*

**ANNUAL GRASSES IN ORNAMENTALS** (continued)

GRASS SPECIES	SCIENTIFIC NAME	WEED* HEIGHT (INCHES)	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
Texas Panicum	<i>Panicum texanum</i>	2 to 6	8	16
Volunteer Cereals				
Barley	<i>Hordeum vulgare</i>	2 to 6	8	16
Oats	<i>Avena sativa</i>	2 to 6	8	16
Rye	<i>Secale cereale</i>	2 to 6	8	16
Wheat	<i>Triticum aestivum</i>	2 to 6	8	16
Volunteer Corn	<i>Zea mays</i>	4 to 12	6	8
Volunteer Corn	<i>Zea mays</i>	12 to 24	8	16
Volunteer Grain Sorghum	<i>Sorghum bicolor</i>	8 to 12	8	16
Wild Oats	<i>Avena fatua</i>	2 to 6	8	16
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	8	16
Witchgrass	<i>Panicum capillare</i>	2 to 8	8	16
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	8	16

\*Generally occurs between 3-leaf stage and tillering.

\*\*Length of lateral growth.

(1) 8 fl. oz./A = approximately 0.2 fl. oz./1000 sq. ft.

(2) 16 fl. oz./A = approximately 0.4 fl. oz./1000 sq. ft.

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).

**ANNUAL BLUEGRASS CONTROL WITH CLEANSE 2 EC IN ORNAMENTALS**

GRASS SPECIES	WEED STAGE	RATE FL. OZ./ACRE	HIGH RATE
Annual Bluegrass ( <i>Poa annua</i> )	to 4-leaf	6	16
<ul style="list-style-type: none"> <li>• Apply under favorable soil moisture and humidity which exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).</li> <li>• Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.</li> <li>• Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.</li> <li>• Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).</li> </ul>			

## PERENNIAL GRASSES

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (inches)	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
<b>Bermudagrass</b>	<i>Cynodon dactylon</i>			
First Application		3 (or up to 6" runners)	8	16
Repeat Application(s) (if regrowth occurs)		3 (or up to 6" runners)	8	16
<b>Quackgrass</b>	<i>Elytrigia repens</i>			
First Application		4 to 8	8	16
Repeat Application(s) (if regrowth occurs)		4 to 8	8	16
<b>Rhizome Johnsongrass</b>	<i>Sorghum halepense</i>			
First Application		12 to 24	8	16
Repeat Application(s) (if regrowth occurs)		6 to 18	6	8
<b>Wirestem Muhly</b>	<i>Muhlenbergia frondosa</i>			
First Application		4 to 8	8	16
Repeat Application(s) (if regrowth occurs)		4 to 8	8	16
(1) 8 fl. oz./A = approximately 0.2 fl. oz./1000 sq. ft.				
(2) 16 fl. oz./A = approximately 0.4 fl. oz./1000 sq. ft.				
Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).				

## STORAGE AND DISPOSAL

### PROHIBITIONS

Do not contaminate water, food or feed by storage, or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place in original container.

**Emergency Response:** For help with any spill, leak, fire or exposure involving this material, call Chemtrec day or night 1-800-424-9300.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

(continued)

## **STORAGE AND DISPOSAL** *(continued)*

**CONTAINER HANDLING: Nonrefillable Container.** Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**Containers 5 gallons or less:** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Containers larger than 5 gallons:** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable Containers larger than 55 gallons: Refill this container with pesticide only. Do not reuse this container for any other purpose.** Cleaning this container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## **LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read this Limitation of Warranty And Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Direct Ag Source, LLC. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants.

Direct Ag Source, LLC does not agree to be an insurer of these risks beyond what is expressly warranted by this label. When you buy or use this product, you agree to accept these risks.

Direct Ag Source, LLC warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

To the extent consistent with applicable law, Direct Ag Source, LLC makes no other express or implied warranty of fitness or of merchantability or any other express or implied warranty.

To the extent consistent with applicable law, in no event shall Direct Ag Source, LLC or seller be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. Buyer's or user's bargained-for expectation is crop protection.

To the extent consistent with applicable law, the exclusive remedy of the user or buyer and the exclusive liability of Direct Ag Source,

LLC or seller, for any and all claims, losses, injuries, or damages (including claims based on breach of warranty or contract, negligence, tort, or strict liability), whether from failure to perform or injury to crops or other plants, and resulting from the use or handling of this product, shall be the return of the purchase price of the product, or at the election of Direct Ag Source, LLC or seller, the replacement of the product.

To the extent consistent with applicable law that allows such requirement, Direct Ag Source, LLC or its Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify Direct Ag Source, LLC or a Direct Ag Source, LLC Retailer of any claims, whether based on contract, negligence, strict liability, or other tort or otherwise be barred from any remedy.

This Limitation of Warranty And Liability may not be amended by any oral or written agreement.

ASANA® - Reg. TM of DuPont

BANVEL® - Reg. TM of BASF Corporation

BASAGRAN® - Reg. TM of BASF AG.

BAYTHROID® - Reg. TM Bayer Corporation

BETAMIX® - Reg. TM of Aventis CropScience

BETANEX® - Reg. TM of Aventis CropScience

BLAZER® - Reg. TM of United Phosphorus, Inc.

BRONATE®, BRONATE ADVANCED™ - Reg. TM of Bayer Crop Science

BUCTRIL® - Reg. TM of Bayer Corporation

CLASSIC® - Reg. TM of E.I. duPont de Nemours & Co., Inc.

COBRA® - Reg. TM of Valent U.S.A. Corporation

DANITOL® - Reg. TM of Sumitomo Chemical Co., Ltd.

DUAL MAGNUM® - Reg. TMs of Syngenta Corporation

EMINENT® - Reg. TM of Sipcam Agro USA, Inc.

FIRSTRATE® - Reg. TM of Dow AgroSciences LLC

FLEXSTAR® - Reg. TM of Syngenta Corporation

FRONTROW™ - TM of Dow AgroSciences LLC

IMI-CORN® - Reg. TM of BASF Corporation

LIBERTY LINK® - Reg. TM of Aventis CropScience

LORSBAN® - Reg. TM of Dow AgroSciences LLC

ORTHENE® - Reg. TM of OMS Investments, Inc.

POUNCE® - Reg. TM FMC Corporation

PROWL® - Reg. TM of BASF Corporation

PURSUIT® and PURSUIT DG® - Reg. TMs of BASF Corporation

RAPTOR® - Reg. TM of BASF Corporation

RESOURCE® - Reg. TM of Valent U.S.A. Corporation

RHONOX® - Reg. TM of Nufarm Americas Inc.

ROUNDUP READY® and ROUNDUP ULTRA® - Reg. TMs of Monsanto Company

SENCOR® - Reg. TM of Bayer Corporation

STINGER® - Reg. TM of Dow AgroSciences LLC

STORM® - Reg. TM of BASF Corporation

SYNCHRONY® STS™ - TM and Reg. TM of E.I. duPont de Nemours & Co. Inc.

WARRIOR® - Reg. TM of Syngenta Corporation



# CLEANSE™ 2 EC

Herbicide

<b>ACTIVE INGREDIENT:</b>	<b>BY WT.</b>
*Clethodim .....	26.4%
<b>OTHER INGREDIENTS:</b> .....	73.6%
<b>TOTAL:</b> .....	100.0%

Contains Petroleum Distillates

\*(E)-2-[1-[[[3-chloro-2-propenyloxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one

Contains 2.0 lbs. clethodim per gal.

## KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See booklet for additional Precautionary Statements and Directions for Use.

### PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

**WARNING.** Causes substantial but temporary eye irritation. Avoid contact with skin. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid breathing vapors or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

EPA Reg. No. 83222-30

EPA Est. No. 37429-GA-001  
37429-GA-002  
42750-MO-001



**DIRECT AG  
SOURCE™**

Manufactured For:  
Direct Ag Source, LLC  
30473 260th Street • Eldora, IA 50627



**FIRST AID** - If in **eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. Call a poison control center or doctor for treatment advice. If on **skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If **swallowed:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. **HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 7:30 AM - 3:30 PM PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information. **NOTE TO PHYSICIANS:** Ingestion of this product or subsequent vomiting can result in aspiration of light hydrocarbon liquid, which can cause pneumonitis. If ingested, probable mucosal damage may contraindicate the use of gastric lavage.

## STORAGE AND DISPOSAL

### PROHIBITIONS

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in cool, dry place in original container.

**Emergency Response:** For help with any spill, leak, fire, or exposure involving this material, call Chemtrec day or night 1-800-424-9300.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING: Nonrefillable container.** Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**Containers 5 gallons or less:** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

CLEANSE-N010515-REVA012915

Net Contents: 1 Gallon (3.79 L)