

Restricted Use Pesticide Due to eye and skin effects. For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

GROUP 12C ACARICIDE

COMITE® II

AGRICULTURAL MITICIDE

NOT FOR USE IN CALIFORNIA, IDAHO, OREGON OR WASHINGTON

INGREDIENTS:	% BY WT.
ACTIVE INGREDIENT:	
Propargite: (Sulfurous acid, 2-[4-(1,1-dimethyl-ethyl) phenoxy] cyclohexyl-2-propynyl ester)*	69.6%
OTHER INGREDIENTS:	30.4%
TOTAL:	100.0%

*Contains 6 lbs active ingredient per gallon. Contains petroleum distillates.

EPA Reg. No. 70506-567

KEEP OUT OF REACH OF CHILDREN

DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

Complete First Aid affixed to front panel. See booklet for complete Precautionary Statements and Directions for Use.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN: Contains petroleum distillate. Vomiting may cause aspiration pneumonia. Probable mucosal damage may contraindicate the use of gastric lavage.	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY: 1-866-673-6671. FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC: 1-800-424-9300 or +1-703-527-3887.	

For Product Use Information Call 1-800-438-6071

Net Contents: _____ Gallons

Produced For: UPL NA Inc. • 630 Freedom Business Center, Suite 402
King of Prussia, PA 19406 U.S.A. • 1-800-438-6071



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Contains petroleum distillate. Corrosive. Causes irreversible eye damage or skin burns. May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Do not breath spray mist. Do not get in eyes, on skin, or on clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All Mixers, Loaders, Applicators, and Other Handlers Must Wear:

- Coveralls over long-sleeved shirt and long pants.
- Chemical-resistant footwear and socks.
- Chemical-resistant gloves made of barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or viton ≥ 14 mils, except for flaggers and applicators using closed cabs.
- Chemical-resistant apron for mixers and loaders, and persons exposed to the concentrate.
- For overhead exposure wear chemical-resistant headgear.
- Wear a NIOSH-approved: Full face respirator with organic vapor cartridges and any R or P combination filters (TC-84A); or Gas mask with an organic vapor canister (TC-14G); or Powered air purifying respirator with an organic vapor cartridge and HE filter (TC-23C).

See **ENGINEERING CONTROLS** for further requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

Mixers and loaders supporting aerial applications to corn and cotton must: use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)] **and must:**

- Wear the personal protective equipment required in **PPE** for mixers/loaders.

Applicators using airblast spray equipment must:

- Wear the personal protective equipment required in **PPE** for mixers and loaders.
- Take off any PPE that was worn in the treated area before reentering the cab if the applicators are operating in a closed cab.
- Store all such PPE in a chemical-resistant container, such as plastic bag, to prevent contamination of the inside of the cab or other enclosed area.

Direct flaggers need to wear the personal protective equipment required above for mixers/loaders, and applicators/handlers, unless they are in an enclosed vehicle which need not be pressurized. When handlers and/or applicators use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Then wash thoroughly and put on clean clothing.
- Remove contaminated clothing and wash before reuse.
- Remove clothing/PPE immediately if pesticide gets inside.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may cause reproductive effects in wildlife. Do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store the product near heat or open flame.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow workers to enter treated areas during the restricted entry interval (REI) listed except as provided for by the WPS. The REI and Exceptions are listed in the Directions for Use associated with the crop. Notify workers of these exceptions (including when entry is permitted for each of the tasks named in the exception).

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, appears below.

- coveralls over long-sleeved shirt and long pants,
- chemical-resistant gloves made of barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or viton ≥ 14 mils, except for flaggers and applicators using closed cabs,
- chemical-resistant footwear plus socks, and
- protective eyewear.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Keep this product in its tightly closed original container. Do not use or store this product near heat or open flame. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING

Nonrefillable container equal to or less than 5 gallons. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows. Empty the remaining contents into application equipment or mix tank and drain container for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain container for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If pressure rinsing: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

RECYCLING

Once cleaned, some agricultural pesticide plastic containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or Ag Container Recycling Council (ACRC) at 877-952-2272 or <http://www.acrecycle.org>. If recycling is not available, dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

SPRAY DRIFT PRECAUTIONS

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS:

- Do not apply by ground within 50 feet or by air within 75 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.
- The above excludes irrigation canals and wasteways as well as man-made irrigation conveyance structures and impoundments, unless an exclusion contains water year-round.
- Risk of exposure to sensitive aquatic areas can be reduced by making applications when the wind direction is away from the aquatic area.
- Consider inclusion of a drift control agent in the spray tank.

Wind Direction and Speed

- Only apply this product if the wind direction favors on target deposition.
- Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

- Do not make aerial or ground applications into temperature inversions.
- Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

- For ground and aerial applications apply the coarsest droplet size spectrum that provides sufficient coverage and mite control.

Additional Requirements for Ground Applications

- For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.
- For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

- The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.
- Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.
- When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

FOR CHEMIGATION APPLICATION, refer to CHEMIGATION USE PRECAUTIONS FOR CORN section; DOSAGE INSTRUCTIONS table exhibits application rate range.

RUNOFF PRECAUTIONS

Under some conditions, propargite may have a high potential for runoff into surface water for several days after application. Do not apply in the following areas:

- frequently flooded areas (excluding crop irrigation areas).
- areas where intense or sustained rainfall is forecasted to occur within 48 hours.

Care should be taken when making applications where the following conditions exist:

- poorly draining or wet soils with readily visible slopes toward adjacent surface water.
- areas with in-field canals or ditches that drain to surface water.
- areas not separated from adjacent surface waters with vegetated filter.

Consult your soil conservation service for recommendations in your use area.

USE RESTRICTIONS

Do not apply in spray solution above pH 10. It is best to use COMITE® II at pH 7 or lower.

Unless the rotated crop is a registered use for propargite, the following applies:

Do not plant any root crop in rotation within 6 months after last application of propargite to labeled crops.

Do not plant other crops in rotation within 2 months after last application of propargite to labeled crops.

Registered-labeled field and vegetable uses for propargite are:

Beans, dry	Mint	Seed crops only:
Corn (sweet, field and popcorn)	Peanuts	Alfalfa
Cotton	Potatoes	Carrot
Hops	Sorghum	Clover
Jojoba		Sugar Beets

PRODUCT USE INSTRUCTIONS

COMITE II is a liquid emulsifiable concentrate for preparing sprays to control mites.

COMITE II is not systemic in action, therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. COMITE II can be used in concentrate sprays as directed by label.

COMITE II performance is best when day temperatures average above 70°F.

MIXING INSTRUCTIONS

After spray tank is three-quarters full of water, add recommended dosage (from table) of COMITE II to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

COMPATIBILITIES

Do not use penetrating surfactants. Under certain conditions a non-penetrating surfactant can be added. Rely on your prior use experience or consult your local surfactant supplier for specific recommendations.

Do not apply in combination with petroleum based foliar spray oils as phytotoxicity may occur.

COMITE II is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of COMITE II will be reduced.

Except as noted, COMITE II is compatible with many pesticides. Observe all cautions and limitation of labeling of all products used in mixtures. However, due to variations in water quality, e.g. hardness and pH, it is required that users conduct small scale trials under local conditions to ensure compatibility prior to large scale use.

Contact your supplier or the company representative for additional information on compatibility.

DOSAGE INSTRUCTIONS

CORN (Field - Grain or Silage) (Seed, Sweet, Popcorn)		
MITES CONTROLLED	APPLICATION RATE (fl ozs/A)	APPLICATION INFORMATION
Two-spotted spider Banks grass	36 - 54	Apply a single high rate application to areas with high mite infestation and to areas which historically have had high mite infestations. At least 48 fl oz per acre should be used under conditions of severe mite pressure. Treat early. Best results are achieved when mite populations are low, before the mites have damaged the plants. Treatment is recommended when mite populations are just beginning to build, and the corn is still small enough (usually 3 to 4 feet) to allow good coverage. Ensure application is made only when corn leaves are dry. This is especially true for white corn varieties which are the most susceptible to plant injury. See COMPATIBILITIES section.
<p>Ground Application: Apply in a minimum of 20 gallons total volume per acre.</p> <p>Aerial Application: Apply in a minimum of 3 gallons total volume per acre.</p> <p>Chemigation Application: Apply in 0.1 to 0.2 acre inches of water. (See CHEMIGATION USE PRECAUTIONS FOR CORN.)</p>		
<p>RESTRICTIONS FOR CORN (Field - Grain or Silage) (Seed, Sweet, Popcorn):</p> <ul style="list-style-type: none"> • Do not apply more than 54 fl ozs of COMITE II per acre per calendar year (2.5 lbs propargite ai per acre per calendar year). • Do not make more than 1 application per season. • The Restricted Entry Interval (REI) is 13 days. See AGRICULTURAL USE REQUIREMENTS for general REI exemptions. There is no exception allowed for hand weeding. • Pre-harvest Interval: Do not harvest within 30 days of the last application. • Similarly, the grazing or cutting for silage within 30 days after treatment is prohibited. 		

COTTON
(East of the Rocky Mountains)

MITES CONTROLLED	APPLICATION RATE (fl ozs/A)	APPLICATION INFORMATION
Two-spotted spider Pacific spider Strawberry spider	Early: 20 Mid-Season to Layby: 24 - 36 Layby to Boll Opening: 36	Apply high rate to areas with high mite infestation and areas which historically have had high mite infestations. COMITE II may be phytotoxic to young cotton under certain conditions. Age and climatic conditions result in a physiological hardening when plants are generally between 6 and 12 inches tall. Applications can begin after plants are hardened. COMITE II may be used in concentrate sprayers designed for concentrate row crop applications. Maintain COMITE II dosage rate per acre. Commercially acceptable mite control may be difficult to obtain where synthetic pyrethroids are or have been used. See COMPATIBILITIES section.

Ground Application: Apply in 15 to 30 gallons total volume per acre for early cotton; 25 to 40 gallons for mid-season to layby cotton; 25 to 50 gallons for layby to boll opening cotton.

Aerial Application: Apply in 5 to 15 gallons total volume per acre from mid-season to boll opening; aerial application is not recommended for early cotton.

RESTRICTIONS FOR COTTON (East of the Rocky Mountains):

- Do not apply more than 70 fl ozs of COMITE II (3.3 lbs propargite ai) per acre per calendar year. Apply only before bolls open.
- Do not apply COMITE II after bolls have opened.
- Do not feed treated foliage or cotton trash to livestock.
- Do not make more than 2 applications per season.
- The minimal spray interval is 21 days.
- The Restricted Entry Interval (REI) is 6 days. See **AGRICULTURAL USE REQUIREMENTS** for general REI exemptions. There is no exception allowed for hand weeding.
- **Pre-harvest Interval:** Do not harvest within 50 days of the last application.

ORANGES and GRAPEFRUIT (Bearing)
(Florida and Texas only)

MITES CONTROLLED	APPLICATION RATE (fl ozs/A)	APPLICATION INFORMATION
Citrus red Texas citrus Six-spotted Citrus rust False spider Broad spider	36 - 52	Apply high rate to areas with high mite infestation and to areas which historically have had high mite infestations. Do not apply petroleum oils 30 days prior to or 40 days after applying COMITE II as phytotoxicity may occur on leaves and or fruit. Do not apply COMITE II mixed with petroleum oils or any material containing petroleum based solvents as phytotoxicity may occur on leaves and/or fruit. Do not mix COMITE II with or apply prior to, or after, alkaline materials such as lime, Bordeaux mixture or lime sulfur. Avoid COMITE II applications to immature leaves and fruit, and prevent spray solution from puddling, dripping, or running off leaves and/or fruit. See COMPATIBILITIES section.

Ground Application: Apply in a minimum of 25 gallons total volume per acre. Apply in sufficient water for thorough coverage.

Aerial Application: Apply in a minimum of 10 gallons total volume per acre. Apply in sufficient water for thorough coverage.

RESTRICTIONS FOR ORANGES and GRAPEFRUIT (Bearing) (Florida and Texas only):

- Do not apply more than 87 fl ozs of COMITE II (4.1 lbs propargite ai) per acre per calendar year.
- Do not make more than 2 applications per season.
- The minimal spray interval is 28 days.
- The Restricted Entry Interval (REI) is 20 days. See **AGRICULTURAL USE REQUIREMENTS** for general REI exemptions. There is no exception allowed for hand weeding.
- **Pre-harvest Interval:** Do not harvest within 21 days of the last application.

PEANUTS		
MITES CONTROLLED	APPLICATION RATE (fl ozs/A)	APPLICATION INFORMATION
Two-spotted spider	36	When temperatures are greater than 90°F with high humidity some leaf phytotoxicity may occur. See COMPATIBILITIES section.
Ground Application: Apply in a minimum 20 gallons total volume per acre.		
Aerial Application: Apply in a minimum of 5 gallons total volume per acre.		
RESTRICTIONS FOR PEANUTS:		
<ul style="list-style-type: none"> Do not apply more than 72 fl ozs of COMITE II per acre per calendar year (3.3 lbs propargite ai per acre per calendar year). Do not make more than 2 applications per season. The minimal spray interval is 14 days. Do not graze or feed livestock on treated areas or cut treated forage for hay. The Restricted Entry Interval (REI) is 2 days. See AGRICULTURAL USE REQUIREMENTS for general REI exemptions. There is no exception allowed for hand weeding. Pre-harvest Interval: Do not harvest within 14 days of the last application. 		

SORGHUM (East of Rocky Mountains)		
MITES CONTROLLED	APPLICATION RATE (fl ozs/A)	APPLICATION INFORMATION
Two-spotted spider Banks grass	24 - 36	Apply high rate to areas with high mite infestation and to areas which historically have had high mite infestations. Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check for potential phytotoxicity before using COMITE II on a large area. See COMPATIBILITIES section.
Ground Application: Apply in a minimum of 20 gallons total volume per acre.		
Aerial Application: Apply in a minimum of 5 gallons total volume per acre.		
RESTRICTIONS FOR SORGHUM (East of Rocky Mountains):		
<ul style="list-style-type: none"> Do not apply more than 36 fl ozs of COMITE II per acre per calendar year (1.7 lbs propargite ai per acre per calendar year). Do not make more than 1 application per season. The Restricted Entry Interval (REI) is 2 days. See AGRICULTURAL USE REQUIREMENTS for general REI exemptions. There is no exception allowed for hand weeding. Pre-harvest Interval: Do not harvest silage within 30 days of the application. Do not harvest grain within 60 days of the application. 		

CHEMIGATION USE PRECAUTIONS FOR CORN

- A. Apply this product only through sprinkler systems, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make the necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.

- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application.
- N. Inject the product with a positive displacement pump into the main line ahead of a right angle turn, to insure adequate mixing.
- O. Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness.
- P. Do not apply when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.
- Q. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of more dilute mixture per hour. Pesticide should be applied continuously for the duration of the water addition.
- R. Where sprinkler irrigation patterns do not overlap sufficiently unacceptable mite control may result. Where sprinkler distribution patterns overlap excessively crop injury may result.
- S. Check with state lead agencies for state specific chemigation requirements.

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of UPL NA Inc., and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. All such risks shall be assumed by the user or buyer.

UPL NA Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to UPL NA Inc., and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC., MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF UPL NA INC. IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF UPL NA INC., MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT UPL NA INC.'S ELECTION, THE REPLACEMENT OF THE PRODUCT.

UPL, the UPL logo and COMITE are trademarks of a UPL Corporation Limited Group Company.

©2022 UPL Corporation Limited Group Company

ESL072017-9730-061422