



FOR CONTROL OF LISTED BROADLEAF WEEDS IN WHEAT, BARLEY, OATS, TRITICALE AND RYE

ACTIVE INGREDIENT:

Octanoic acid ester of bromoxynil (3,5-dibromo-4-hydroxybenzotrile)*	30.1%*
Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid**	31.2%**
OTHER INGREDIENTS	38.7%
TOTAL	100.0%

Contains xylene range/petroleum distillates.

*Bromoxynil octanoate equivalent to 22.9% of bromoxynil or not less than 2.0 pounds of bromoxynil per gallon.

**2,4-Dichlorophenoxyacetic acid equivalent to 20.69% or not less than 1.9 pounds per gallon. Isomer specific by AOAC method.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**For Chemical Spill, Leak, Fire, Exposure or Accident,
Call CHEMTREC 1-800-424-9300
For Medical Emergencies Only, Call 1-877-325-1840**

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor for treatment advice. Do not give **any** liquid to the person. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or a doctor for treatment advice. **IF INHALED:** Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN

May pose an aspiration pneumonia hazard. Contains petroleum distillate.

SEE INSIDE LEAFLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND COMPLETE DIRECTIONS FOR USE.

NET CONTENTS:

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION - PRECAUCIÓN

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not store near fertilizers or seeds. Store at temperatures above 32°F. If allowed to freeze, remix before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Non-refillable (<5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Non-refillable (>5 gallons):** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. **Refillable Container:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

EPA Reg. No. 5905-552

EPA Est. No. 228-IL-1

AD 022712

Patent No. 6,232,272

Manufactured For
HELENA CHEMICAL COMPANY
225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE, TENNESSEE 38017

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

12642000

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION – PRECAUCIÓN

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are barrier laminate, butyl rubber, nitrile rubber or viton. If you want more options, follow the instructions for category G on an EPA chemical resistant category selection chart.

All mixer, loader, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes and socks, plus
- Protective eyewear
- Chemical-resistant gloves except for pilots
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE Separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

If you will handle a total of 60 gallons or more of this product per day, you must use a mechanical transfer system for all mixing and loading operations. If this product is packaged in a 30 gallon drum, you must use a mechanical transfer system which terminates in a drip-free hard coupling which may be used only with a spray or mix tank which has been filled with a compatible coupling. If you do not presently own or have access to a mechanical transfer system with this type of coupling, contact your dealer for information on how to obtain such a system or to modify your present system. When using a mechanical transfer system, do not remove or disconnect the pump or probe from the container until the container has been emptied and rinsed. The pump or probe system must be used to rinse the empty container and to transfer the rinsate directly to the mixing or spray tank.

To reduce exposure to residues, wash the spray rig, tractor, and all other equipment used to handle or apply this product with water daily or before using the equipment for any other purpose.

See "Engineering Controls" for additional requirements.

Engineering Controls

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

APPLICATION BY CHEMIGATION must be done by fixed pipe, overhead sprinkler systems or hand moved pipe. If hand-moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle.

DURING AERIAL APPLICATION, human flaggers are prohibited unless in enclosed vehicles. Aerial application is prohibited within 300 feet of residential areas (e.g., homes, schools, hospitals, shopping areas, etc.).

USER SAFETY

Users should:

- Wash hands before eating, drinking, chewing
- Remove clothing immediately if pesticide gets on skin, wash immediately
- Remove PPE immediately after handling the concentrate. As possible, wash thoroughly and change if possible.

ENVIRONMENTAL

This pesticide is toxic to fish and aquatic invertebrates present, or to intertidal areas below the mean low water mark of organisms in water adjacent to treated areas. Do not discharge into water or rinsate.

This chemical has properties and characteristics that may contribute to groundwater contamination in areas where soils are permeable. Application around groundwater.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with the label directions for use.

Do not apply this product in a way that will contribute to the drift of pesticides to protected handlers may be in the area during application. Consult the agency responsible for pesticide regulation.

AGRICULTURAL USES

Use this product only in accordance with its label directions. This standard contains requirements for use in greenhouses, and handlers of agricultural pesticides. The statements on this label about personal protective equipment in this box only apply to uses of this product.

Do not enter or allow worker entry into treated areas until the REI is 24 hours.

PPE required for early entry to treated areas involves contact with anything that has been treated. PPE includes long pants, chemical-resistant gloves, and protective eyewear.

PHYSICAL AND CHEMICAL

Do not use or store near heat or open flame.



USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI). For all crops on this label the REI is 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

STO

Do not contaminate water, food, or feed. Do not use. **PESTICIDE STORAGE:** Do not store in cold, freeze, remix before using.

PESTICIDE DISPOSAL: Wastes resulting from use should be disposed of at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable: Empty the container. Offer for recycling, if available. Do not reuse.

Non-Refillable (<5 gallons): Triple rinse or a mix tank and drain for 10 seconds after each use. Shake for 10 seconds. Pour rinsate into disposal container. Drain for 10 seconds after each use.

Non-Refillable (>5 gallons): Triple rinse or a mix tank. Fill the container 1/4 full with water. Roll it back and forth, ensuring at least one full revolution. Empty the rinsate into a disposal container. Repeat this procedure two more times.

Pressure rinse as follows: Empty the container. Triple rinse for 10 seconds after the final use. Rinse the container, then the remaining contents from this container, and rinse at about 40 PSI.

Refillable Container: Refill this container only after it has been cleaned. Cleaning the container before refilling is the responsibility of the user. Cleaning before refilling is the responsibility of the user. Cleaning before refilling is the responsibility of the user. Cleaning before refilling is the responsibility of the user. Cleaning before refilling is the responsibility of the user.

C

DOUBLE UP® B+D is formulated as a suspension concentrate containing the equivalent of 2 pounds of bromoxynil per gallon of 1.9 pounds per gallon.

DOUBLE UP® B+D is a selective pre-emergent herbicide for small grains wheat, barley, oats, rye applied to actively growing weed seeds. It is a selective pre-emergent herbicide, therefore thorough coverage of the soil is necessary.

DOUBLE UP® B+D has little residual activity. Generally, crops that require pre-emergent herbicide should be planted to shade the subsequent flushes of weed growth.





UTIONS

ing the toilet.
ughly and put on clean clothing. If
of gloves before removing. As soon

IS

o water, to areas where surface water
runoff may be hazardous to aquatic
en disposing of equipment washwa-

detected in groundwater. The use of
water table is shallow, may result in
in contamination of drinking water or

its labeling. Read entire label before

s, either directly or through drift. Only
is specific to your State or Tribe, con-

MENTS

Protection Standard, 40 CFR Part
orkers on farms, forests, nurseries,
rements for training, decontamina-
ctions and exceptions pertaining to
and restricted-entry interval. The
by the Worker Protection Standard.
entry interval (REI). For all crops on

orker Protection Standard and that
or water, is coversalls over a long-
roof material, shoes plus socks and

ZARDS

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not store near fertilizers or seeds. Store at temperatures above 32°F. If allowed to freeze, remix before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Non-Refillable (<5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Non-Refillable (>5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable Container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

GENERAL INFORMATION

DOUBLE UP® B+D is formulated as an emulsifiable concentrate of octanoic acid ester of bromoxynil containing the equivalent of 2 pounds of bromoxynil per gallon and 2,4-Dichlorophenoxyacetic acid containing the equivalent of 1.9 pounds per gallon.

DOUBLE UP® B+D is a selective postemergence herbicide for control of important broadleaf weeds infesting small grains wheat, barley, oats, rye and triticale. Optimum weed control is obtained when **DOUBLE UP® B+D** is applied to actively growing weed seedlings. **DOUBLE UP® B+D** contains a contact herbicide and a systemic herbicide, therefore thorough coverage of the weed seedlings is essential for optimum control.

DOUBLE UP® B+D has little residual activity. Therefore subsequent flushes of weeds will not be controlled by the initial treatment. Generally, crops that form a good canopy will help shade subsequent weed flushes. However, certain crops or short-straw varieties (for example, Yaccora Rojo wheat) may not develop the crop canopy fast enough to shade the subsequent flushes of weeds.

Use a standard herbicide b
equipped with screens no fin

Select a spray volume and d
spray distribution and throu
of 40–60 psi are recomme
droplets may not provide ad
flood nozzles are not recom

In general, a spray volume o
minimum of 5 GPA with a mi
used with higher speed, low
spray distribution. When usi
conditions cause excessive
applications made when dry,
Applications using less than

When weed infestations are
uniform weed coverage. Wh
nozzles should be used to ob
consult your local Extension

Do not apply when winds are
movement.

Avoid use of small diameter
ers, tomatoes, beans, sugart
spray at all in the vicinity nee
of herbicides. State agricultu
that the use of this product c

If tank mixing, a compatibility
ibility, use a small container a
ratio as the anticipated use.
spraying. Indications of inco
mum crop safety and weed
used in the tank mixture.

Use orifice discs, cores and
and maximum coverage. In
recommended.

Do not apply during inversion
erage and/or off-target spray
volume per acre and not app



DISPOSAL

Apply at temperatures above 32°F. If allowed to

the product may be disposed of on site or at an

30 & 55 gallon): Do not reuse or refill this container (or equivalent) promptly after

transferring contents into application equipment. Rinse the tip. Fill the container 1/4 full with water and empty into a mix tank or store rinsate for later use or repeat this procedure two more times.

Transferring contents into application equipment. Rinse the tip. Fill the container on its side and empty into a mix tank or store rinsate for later use or disposal. Stand the container on its end and tip it back and forth several times or store rinsate for later use or disposal.

Use application equipment or a mix tank and container upside down over application equipment. Rinse the pressure rinsing nozzle in the side of the container for 10 seconds after the flow begins to drip.

Do not reuse this container for any other purpose. The responsibility of the person disposing of the container. Rinse the container before final disposal, empty into a mix tank. Fill the container about 10% with water and run the pump for 2 minutes. Pour or pump rinsate at this rinsing procedure two more times.

ATTENTION

Double Up B+D is a 2,4-D choline salt containing 2,4-D choline salt containing the equivalent

Control of important broadleaf weeds infesting crops is obtained when **DOUBLE UP® B+D** is used. It maintains a contact herbicide and a systemic herbicide for optimum control.

Double Up B+D flushes of weeds will not be controlled by the double up B+D. However, weeds may not develop the crop canopy fast enough

GROUND APPLICATION

Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with screens no finer than 50 mesh in the nozzle tips and in-line strainers.

Select a spray volume and delivery system that will ensure thorough and uniform spray coverage. For optimum spray distribution and thorough coverage use of flat fan nozzles (maximum tip size 8008) with a spray pressure of 40–60 psi are recommended. Other nozzle types and lower spray pressures that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control. **Raindrop®** nozzles and flood nozzles are not recommended as weed control with **DOUBLE UP® B+D** may be reduced.

In general, a spray volume of 10 to 20 gallons per acre (GPA) is recommended for optimum spray coverage. A minimum of 5 GPA with a minimum spray pressure of 50 psi and a maximum ground speed of 10 mph may be used with higher speed, low volume ground application if ground terrain, crop and weed density allow effective spray distribution. When using higher speed equipment, a maximum ground speed of 10 mph is suggested if field conditions cause excessive boom movement during application which results in poor spray coverage. Ground applications made when dry, dusty field conditions exist may provide reduced weed control in wheel track areas. Applications using less than 10 gallons per acre may result in reduced weed control.

When weed infestations are heavy, use of higher spray volumes and spray pressure will be helpful in obtaining uniform weed coverage. When corn or grain sorghum are large enough to interfere with the spray pattern, drop nozzles should be used to obtain uniform weed coverage. If you are unsure of the infestation level or size of crop, consult your local Extension Service.

Do not apply when winds are gusty or when other conditions favor poor spray coverage and/or off-target spray movement.

USER PRECAUTIONS

Avoid use of small diameter spray nozzles. Avoid spray drifts onto susceptible plants such as vegetables, flowers, tomatoes, beans, sugarbeets, sunflowers, and other legumes. Coarse sprays are less likely to drift. Do not spray at all in the vicinity near susceptible plants to prevent any vapor drift. Local conditions may affect the use of herbicides. State agricultural authorities in many states issue recommendations to fit local conditions. Be sure that the use of this product conforms to all applicable conditions.

If tank mixing, a compatibility test is recommended to ensure satisfactory spray preparation. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing. To ensure maximum crop safety and weed control, follow all cautions and limitations on this label and the labels of products used in the tank mixture.

AERIAL APPLICATION

Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage. In general a minimum spray volume of 5 GPA and a maximum pressure of 40 psi are recommended.

Do not apply during inversion conditions, when winds are gusty or when other conditions favor poor spray coverage and/or off-target spray movement. Off-target spray movement can be minimized by increasing the spray volume per acre and not applying when winds exceed 10 mph.



SPRAY DRIFT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

FOR THE CONTROL OF LISTED BROADLEAF

Use 3/4 to 2 pints of this herbicide per acre in evenly 10–20 gallons per acre by ground equipment grains from the fully tillered stage of growth but before weed and wild buckwheat. Use the 2 pints rate for cool, dry conditions. Heavy weed populations require

This product controls the following weeds: mugwort, gumweed, pigweed, ragweed, wild buckwheat, smartweed, lambsquarter and other broadleaf weeds whichever comes first.

RESTRICTIONS AND PRECAUTIONS:

- Do not permit dairy animals or meat animals be in the field for 3 days after treatment.
- Do not apply when crop canopy covers the weeds.
- Do not apply when crops are under moisture stress.
- Reduced weed control may occur when weeds are present.
- Do not plant rotational crops within 30 days following application.
- Limited to 1.75 lbs. ae per acre per crop cycle.
- Do not feed treated straw to livestock.
- The preharvest interval (PHI) is 14 days.

Postemergence:

- Limited to one postemergence application per crop cycle.

Preharvest:

- Limited to one preharvest application per crop cycle.
- Maximum of 2 pints/acre per application.



FOR THE CONTROL OF LISTED BROADLEAF WEEDS IN WHEAT, BARLEY, OATS, TRITICALE AND RYE

Use 3/4 to 2 pints of this herbicide per acre in enough spray volume to provide uniform coverage of weeds, usually 10–20 gallons per acre by ground equipment and 5 gallons by aircraft. Application should be made to cereal grains from the fully tillered stage of growth but before the jointing stage. Use the 2 pints rate for mustards, pigweed and wild buckwheat. Use the 2 pints rate for kochia and tansy mustard or when weeds are growing under cool, dry conditions. Heavy weed populations require a higher rate and complete spray coverage.

This product controls the following weeds: mustards, frenchweed, nightshade, arrowhead, field bindweed, gumweed, pigweed, ragweed, wild buckwheat, plantain, sunflower, canola, cow cockle, Russian thistle, annual smartweed, lambsquarter and other broadleaf weeds. Apply to weeds up to the 4-leaf stage or 2 inches in height, whichever comes first.

RESTRICTIONS AND PRECAUTIONS:

- Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 45 days after treatment.
- Do not apply when crop canopy covers the weeds as poor weed control will result.
- Do not apply when crops are under moisture stress.
- Reduced weed control may occur when weeds are stressed from lack of moisture or cold temperatures.
- Do not plant rotational crops within 30 days following application.
- Limited to 1.75 lbs. ae per acre per crop cycle.
- Do not feed treated straw to livestock.
- The preharvest interval (PHI) is 14 days.

Postemergence:

- Limited to one postemergence application per crop cycle. Maximum of 5 pints/acre per application.

Preharvest:

- Limited to one preharvest application per crop cycle.
- Maximum of 2 pints/acre per application.

CONDITIONS AND LIMITATIONS

Read the Conditions of Sale – Warranty of the product. If the terms are not accepted, the product will be refunded.

The directions on this label are believed to be true and/or injury to the crop to which the product is applied under weather conditions or the failure to follow the control of Helena Chemical Company may cause injury to crops, animals, man or property. The chemical description on the label and the factors noted above which are warranties or representations of any kind, are not a part of merchantability or fitness for any particular purpose.

The exclusive remedy against the Company shall be limited to, at Helena Chemical Company's discretion:

1. Refund of the purchase price paid by the customer.
2. Replacement of the product used.

To the extent allowed by law, the Company shall not be liable for special, indirect, incidental, or consequential damages, including loss of profits or income. The Company's liability is limited to the purchase price of the product and is subject to the foregoing conditions of sale.

© Copyright Helena Holding Company, 2012

RAINDROP® is a registered trademark of Helena Chemical Company. **DOUBLE UP®** is a registered trademark of Helena Chemical Company.



**RYE, OATS, TRITICALE AND RYE**

uniform coverage of weeds, usual application should be made to cereal at the 2 pints rate for mustards, pigweed or when weeds are growing under the spray coverage.

arrowhead, field bindweed, wild cockle, Russian thistle, annual bluegrass 4-leaf stage or 2 inches in height,

in large treated grain fields within 45

days result.

in low temperatures or cold temperatures.

at 1 pint/acre per application.

CONDITIONS OF SALE – LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale – Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2012.

RAINDROP® is a registered trademark of Delavan Corporation.
DOUBLE UP® is a registered trademark of Helena Holding Company.

FOR CONTROL OF LISTED**ACTIVE INGREDIENT:**

Octanoic acid ester of bromoxynil (3, Isooctyl (2-ethylhexyl) Ester of 2,4-Di

OTHER INGREDIENTS

TOTAL

Contains xylene range/petroleum dis
*Bromoxynil octanoate equivalent to
**2,4-Dichlorophenoxyacetic acid eq

**KEEP
CAUTION**

If used no entien
(If you do n

**For Chemical Spill, L
For**

IF IN EYES: Hold eye open and rinse with water for first 5 minutes, then continue rinsing with water.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not drink milk or eat fatty foods.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Wash skin thoroughly with soap and water. Stop and get medical attention if irritation occurs.

IF INHALED: Remove person to fresh air. If breathing is difficult, call a doctor immediately. Do not use mouth-to-mouth resuscitation if possible. Call a doctor immediately.

Have the product container or label with you when calling a poison control center or doctor for product information. In Canada, call 1-800-424-9300 for emergency information.

May pose an aspiration pneumonia hazard if inhaled.

AD 022712
Patent No. 6,232,272

**H
225 SCHILLING BLVD**

Job #59985



WARRANTY AND REMEDIES


ability and Remedies before using this product, and the full purchase price

lowed carefully. Insufficient control of pests in the occurrence of extraordinary or unusual application practices, all of which are beyond in addition, failure to follow label directions on any warrants that this product conforms to use referred to in the directions for use sub-company. The Company makes no other warranty on the product, including no implied warranty. Any warranty shall be implied by law.

relating to the handling or use of this product as follows:

weight, or

any and all claims against the Company are of no force or effect, in whole or in part, in the absence of any nature, including, but not limited to, the product and the buyer and user accept it, including any and all remedies.



FOR CONTROL OF LISTED BROADLEAF WEEDS IN WHEAT, BARLEY, OATS, TRITICALE AND RYE

ACTIVE INGREDIENT:

Octanoic acid ester of bromoxynil (3,5-dibromo-4-hydroxybenzoxirone)* 30.1%*
Isocetyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid** 31.2%**

OTHER INGREDIENTS 38.7%

TOTAL 100.0%

Contains xylene range/petroleum distillates.

*Bromoxynil octanoate equivalent to 22.9% of bromoxynil or not less than 2.0 pounds of bromoxynil per gallon.

**2,4-Dichlorophenoxyacetic acid equivalent to 20.69% or not less than 1.9 pounds per gallon. Isomer specific by AOAC method.

KEEP OUT OF REACH OF CHILDREN CAUTION – PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

**For Chemical Spill, Leak, Fire, Exposure or Accident, Call CHEMTREC 1-800-424-9300.
For Medical Emergencies Only, Call 1-877-325-1840.**

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or a doctor for treatment advice.

IF INHALED: Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN

May pose an aspiration pneumonia hazard. Contains petroleum distillate.

AD 022712
Patent No. 6,232,272

EPA Reg. No. 5905-552
EPA Est. No. 228-IL-1

Manufactured For
HELENA CHEMICAL COMPANY
225 SCHILLING BOULEVARD, SUITE 300 • COLLIERVILLE, TENNESSEE 38017

Job #59985