

# DRYACIDE® CIK

## Kills Crawling Insects

### ACTIVE INGREDIENT:

Diatomaceous Earth ..... 100%  
(including silicon dioxide, other oxides and moisture)

## KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 69261-4-1381

EPA Est. No. 69261-OR-1

1/1015/4

DRYACIDE® CIK consists of the skeletal remains of fresh water diatoms and contains amorphous silicon dioxide. This product is intended for indoor and outdoor use to control ants, bedbugs, spiders, fleas, houseflies, little houseflies, German cockroaches, mites, earwigs and the other insect listed on this label.

### FIRST AID

**IF IN EYES:** Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call 1-877-424-7452 for further medical information.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Use adequate ventilation and avoid breathing dust. For prolonged exposure to dust, use a suitable dust mask approved by NIOSH/MSHA.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a ready-for-use insecticide dust which has given effective knockdown and control of the pests listed.

For use in residential and commercial pest control, in and around buildings and structures and their immediate surroundings, and on modes of transport.

Intended for application with hand or power duster, or by other suitable means, to cracks, crevices, voids, hiding and runway areas, and places where pests are found.

In living areas, make applications in such a manner as to avoid deposits on exposed surfaces or introducing the material into the air. Treat a small area of the surface to be treated and look for staining of other adverse reaction before making regular application.

To apply insecticide directly into cracks and crevices, use a bulbous duster or other suitable equipment. Apply lightly and uniformly to infested areas. Pay particular attention to cracks, crevices and voids where insects may be harboring, traveling,

breeding or entering the structure. Examples of these areas are service ducts; false floors and ceilings; wall voids; around electrical and telephone fittings and equipment; around water and sewer pipes; under and behind cabinets, refrigerators and sinks; around window and door frames; along baseboards; in attics and crawl spaces; landscaped and mulched areas. The amount of product to be applied will vary with the from site to site but will usually be in the range of 0.15 – 0.6 oz/10 ft<sup>2</sup> or 1 – 5 oz./100 ft<sup>2</sup>. To treat against Black Widow Spiders and Pharaoh Ants, apply at a rate of 0.633 oz/10 ft<sup>2</sup> or 6.33 oz/100 ft<sup>2</sup>. Repeat treatments as necessary to maintain adequate control.

**Asian Lady Beetles, Boxelder Bugs and Cluster Flies:** Apply where these insects may be harboring, traveling, breeding or entering the structure. Food Areas of Food/Feed Handling Establishments: Food/feed handling establishments are places other than private residences in which food/feed is held, processed, prepared or served. In food handling areas, or areas where food or feed products are prepared or stored, apply as a Crack & Crevice® or spot treatment. As a Crack & Crevice treatment, apply sparingly into cracks and crevices using equipment designed to place powder into specific locations. Limit spot treatments to walls or floors where pests are suspected of hiding. No individual spot treatments should exceed 2 ft<sup>2</sup>. Remove or cover all food or feed items in the treatment area before applying product. Treat open spaces below floors, above ceilings, within walls, under stairs, underneath and behind equipment or appliances, around pipes and sinks, in utility closets and around garbage and waste cans. For best results, apply to dry areas. Repeat treatment on dry surfaces, as needed, to maintain control of problem pests.

### OUTDOOR USE

Apply as a treatment in and around cracks, crevices or voids associated with windows, doors, porches, screens, eaves, patios, garages, under stairways, around foundations, sewers, animal burrows, mulched areas, manure or compost piles and in crawl spaces and other areas where insects may be harboring, traveling, breeding or entering the structure.

**\* In the home, exposed food, food processing and utensils in the treatment areas should be covered, cleaned or removed.**

**\*\* Do not apply to classrooms when in use. Do not apply to institutions (including libraries, sport facilities, etc.) in the immediate area when occupants are present.**

**† Remove patients before using. After treatment, ventilate rooms for 30 minutes before returning patients to rooms.**

**†† Do not apply in aircraft cabins.**

### LIVESTOCK HOUSING STRUCTURES

For use in and around livestock housing structures, swine houses, cattle barns, calf hutches, loafing barns, horse stables and pet kennels. To control ants, spiders, fleas, houseflies, little houseflies, cockroaches, mites, litter beetles, hide beetles, earwigs and other crawling insects, apply dust evenly at a rate of 1 pound per 150 square feet. Apply dust using hand or power duster or other suitable equipment. Apply uniformly to infested areas of walls and floors. Pay particular attention to cracks and crevices, bedding, wall voids and around window and door frames. Applications are permitted when animals are present. Do not apply directly to animal feed or watering

equipment. Repeat treatments as necessary to maintain adequate control.

### POULTRY HOUSES

To control bedbugs, mites, litter beetles (darkling beetles) and hide beetles, apply dust to litter at a rate of 4 pounds per 1000 square feet of surface area. Uniformly cover floors and walls, paying particular attention to cracks and crevices, corners and underneath feed and watering equipment and lines. Avoid contamination of feed and water. Applications can be made up to 75 pounds per 1000 square feet for severe infestations and in moist areas. If litter is greater than 3 inches deep, apply up to an additional 25 pounds per 1000 sq. ft. per additional inch of litter. Applications are permitted mid-flock while birds are present. Litter should be retreated when replaced and before each new batch of poultry is introduced to the facility.

### INDOOR & OUTDOOR TREATMENTS

**Crawling Insects:** Apply to cracks, crevices and voids where insects may be harboring, traveling, breeding or entering the structure. For example: around sinks and storage areas; behind baseboards; around doors and windows; behind and under refrigerators, cabinets and desks, sinks and stoves, and attics and crawl spaces. In commercial areas, apply to cracks, crevices and voids; office areas; stainless steel equipment; shelving; machinery; pallets; and other areas where these insects may be harboring, traveling or entering the structure.

**Carpenter Ants:** Apply to cracks, crevices and voids where insects may be harboring, traveling, breeding or entering the structure. If possible, apply dust into infested wood through existing openings or newly drilled small holes. When possible, locate and treat nests, trails and surrounding areas.

**Fleas:** Apply dust to infested areas such as in and around pet beds and resting quarters; nearby cracks and crevices; under the edges of rugs, carpets and floor coverings. Apply a thin coat of dust to rugs and carpets. Rake (not sweep) nap back and forth, causing dust to enter rug or carpet. Leave overnight. Repeat applications and vacuum every 3-4 days as necessary for effective flea control. Dryacide CIK will kill live fleas. Treat cracks, crevices and voids around and under upholstered furniture, and other areas where these pests frequent. When treating furniture, rugs and fabrics, treat a small area first to check for staining before treating large areas. Do not treat animals with this product.

### Bees, Wasps, Hornets, Paper Wasps and Yellowjackets:

It is generally advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Wear protecting clothing, if deemed necessary, to avoid stings. Using hand or power duster, or other suitable means (with extension tubes if necessary), thoroughly dust nest, nest entrance and surrounding areas where insects alight. Nests in voids should be located and treated. If necessary, drill a hole into the void(s), blow dust in, and reseal. For ground wasps and yellowjackets, dust entrance and surrounding areas. For best results, carefully check nests 1 – 2 days after treatment to ensure complete kill. When treating overhead nests, apply in such a manner as to avoid drift onto the applicator.

**Bed Bugs:** Apply at a minimum rate of 1.6 oz/100 ft<sup>2</sup> to cracks, crevices and voids associated with mattresses, boxsprings, bedframes and other furniture where these insects may harbor, travel, breed or enter the room. Treat mattresses, especially tufts, folds and edges. After four hours vacuum the dust from the mattress. Cover with clean bedding before using. Bed Bugs may also harbor in areas of the room away from the bed or furniture. Apply product to wall voids, behind wall hangings, floor molding, window casings, carpet's edge, popcorn ceilings and other cracks, crevices or voids.

**Carpet Beetles (Black, Furniture and Varied) Webbing Clothes Moths and Spiders:** Apply where these insects may be harboring, traveling breeding or entering the structure. For example: on and under edges of floor coverings; under rugs and furniture; in closets and associated wall voids. When treating furniture, rugs and fabrics, treat a small area first to check for staining before treating a large area.

**Fruit and Phorid Flies:** Apply where these insects harbor, travel, breed or enter the structure. For example: under cabinets or shelving; behind or under equipment or drains; protected exterior areas such as eaves, trash corrals and dumpsters; and other areas where organic debris may accumulate.

**IN CASE OF SPILL.** Vacuum or dampen with water and sweep. Dispose of in trash or at local sanitary landfill.

#### **STORAGE AND DISPOSAL**

**Do not contaminate water, food or feed by storage or disposal.**

**PESTICIDE STORAGE AND SPILL PROC-DURES:** Store in original container in a dry location. In case of spill, vacuum or dampen with water and sweep up.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Empty container by using the product according to the label directions. Do not reuse! Place empty container in trash or offer for recycling if available. If container is partly filled, contact your local solid waste agency for disposal instructions. **NEVER PLACE UNUSED PRODUCT DOWN ANY INDOOR OR OUTDOOR DRAIN!**

#### **WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

#### **LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSE-

QUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

DRYACIDE® is a registered trademark of Winfield Solutions, LLC.

**Manufactured for:**  
Winfield Solutions, LLC  
P.O. Box 64589, St. Paul, MN 55164  
**NET WT. 50 lbs.**