

Dyna-Shield Tibet™

FOR USE ONLY BY COMMERCIAL SEED TREATERS
FOR CONTROL OF SPECIFIED DISEASES ON CORN, WHEAT, BARLEY, TRITICALE AND OATS

ACTIVE INGREDIENT:

Tebuconazole:

Alpha-[2-(4-Chlorophenyl)ethyl]-alpha-

(1,1 -dimethylethyl)-1H-1,2,4-triazole-1-ethanol 28.3%

OTHER INGREDIENTS: 71.7%

TOTAL: 100.0%

Contains 2.6 pounds tebuconazole per gallon.

STOP — READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Medical Emergencies Only, Call 1-866-944-8565

EPA REG. NO. 34704-1110
EPA EST. NO. 34704-XX-XXX

FORMULATED FOR
Loveland Products, Inc.
P.O. Box 1286
Greeley, Colorado 80632-1286



NET CONTENTS 2.5 GAL (9.46 L)

081817 V1D 08G17

FIRST AID

IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth- to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-866-944-8565 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wear long sleeved shirt, long pants or coveralls, shoes and socks, and chemical-resistant gloves made of waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

Groundwater Advisory: Tebuconazole is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

Not for use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

For use by commercial seed treaters only.

When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

IMPORTANT: Read these entire Directions and Warranty before using this product

SEED TREATMENT

This product controls or suppresses certain designated seed and soilborne diseases of corn, wheat, barley, triticale, and oats.

CROP	DISEASE	RATE Fl. oz./100 lbs. of Seed	APPLICATION DIRECTIONS
Corn Field corn, Field corn grown for seed, popcorn, sweet corn	Soilborne and Seedborne <i>Fusarium</i> *	0.075-0.1	Apply this product as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment. Uniform application to seed is necessary to ensure seed safety and best disease protection. Seed must be sound and well cured before treatment. Dilute product with sufficient water to ensure seed is completely covered. Add dye to the resulting slurry. Consult a seed treatment specialist regarding slurry rates specified for the crop to be treated. The length of control will vary depending on the rate used.
	Soilborne and Seedborne Head Smut (<i>Sphacelotheca reiliana</i>)*	0.37-0.74	
Wheat Triticale	Stinking Smut*, Flag Smut*, Loose Smut*, Early Season <i>Septoria</i> Disease Complex, Early Season <i>Rhizoctonia</i> Root Rot, Early Season Common Foot Rot*, Early Season <i>Fusarium</i> Foot Rot, Early Season Suppression of Powdery Mildew, Early Season Suppression of Wheat Leaf Rust	0.08 to 0.10	Apply this product as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment. Prepare a slurry by mixing the specified dosage in 9.0 to 20.0 fluid ounces of water per 100 pounds of seed. Add dye to the resulting slurry. Mix the slurry thoroughly with the seeds to provide uniform coverage.

*Not approved for this use in California

SEED TREATMENT (Continued)

This product controls or suppresses certain designated seed and soilborne diseases of corn, wheat, barley, triticale, and oats.

CROP	DISEASE	RATE Fl. oz./100 lbs. of Seed	APPLICATION DIRECTIONS
Barley	Covered Smut*, Loose Smut*, Barley Stripe (suppression), Early Season <i>Rhizoctonia</i> Root Rot, Early Season Common Root Rot, Early Season <i>Fusarium</i> Foot Rot*, Early Season Suppression of Powdery Mildew, Early Season Suppression of Barley Leaf Rust	0.08 to 0.10	Apply this product as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment. Prepare a slurry by mixing the specified dosage in 9.0 to 20.0 fluid ounces of water per 100 pounds of seed. Add dye to the resulting slurry. Mix the slurry thoroughly with the seeds to provide uniform coverage.
Oats	Loose Smut*, Pythium Damping-off, Early Season <i>Rhizoctonia</i> Root Rot, Early Season Common Root Rot, Early Season <i>Fusarium</i> Foot Rot*, Early Season Suppression of Powdery Mildew, Early Season Suppression of Crown Rust		

*Not approved for this use in California

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with Dyna-Shield Tibet™, a fungicide containing tebuconazole.
- Do not use treated seed for feed, food, or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with tebuconazole:

- Store treated seed away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Barley, wheat and oat, green forage may be grazed or harvested for hay 31 days after seeding.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.
- Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Note to Reviewer: The following statement will be included on all Final Printed Labels bearing multiple Container Handling statements] **NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type / size."

[Nonrefillable Containers 5 Gallons or Less]

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

[Nonrefillable containers larger than 5 gallons]

Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure-rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

[Refillable containers]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC 1 (800) 424-9300

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

For non-emergency (e.g., current product information), call Loveland Products, Inc.
1-888-574-2878.

DYNA-SHIELD Tibet is a trademark of Loveland Products, Inc.

All other trademarks are the property of their respective owners.

Formulated for:
Loveland Products, Inc.
P.O. Box 1286
Greeley, Colorado 80632-1286