

Endow™ 25C

Azoxystrobin Group 11 Fungicide



For control of listed diseases on Turf; Golf courses; Lawns and landscape areas around residential, institutional, public, commercial and industrial buildings; Parks; Recreational areas; Athletic fields; Ornamentals; Conifers including Christmas trees; and Roses, commercial rose production

ACTIVE INGREDIENT:

Azoxystrobin: methyl (E)-2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate* 22.93%

OTHER INGREDIENTS: 77.07%

TOTAL: 100.00%

*IUPAC

Contains 2.08 lbs. Azoxystrobin per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Emergency Phone Numbers: (800) 424-9300 CHEMTREC (transportation and spills)
(800) 222-1222 Poison Control Center

See additional Precautionary Statements and Directions for Use inside booklet.

NET CONTENTS: 2.5 Gallons (9.46 L)

Manufactured for:
SIPCAM AGRO USA, INC.
2525 Meridian Parkway
Durham, NC 27713

2.5G

EPA Reg. No. 60063-59

EPA Est. No.: 60063-GA-001 – Lot number begins with VL

EPA Est. No.: 70815-GA-001 – Lot number begins with CB

EPA Est. No.: 86555-MO-001 – Lot number begins with AF



SIPCAM AGRO
USA, INC.

EPA20210701 (9/21)

**READ THE LABEL
CAREFULLY
BEFORE OPENING
THE CONTAINER**

FUNGICIDE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning. May be fatal if swallowed. Harmful if absorbed through the skin. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Human flagging is prohibited.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils
- In addition, mixers/loaders/applicators using mechanically pressurized handwands except when applying to Christmas tree farms, nursery ornamentals, landscaping, must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Respirator fit testing, medical qualification, and training

Using a program that conforms to OSHA's requirements (see 29 CFR Part), employers must verify that any handler who uses a respirator is:

- Fit-tested and fit-checked,
- Trained, and
- Examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn. A qualified medical practitioner is a physician or other licensed health care professional who will evaluate the ability of a worker to wear a respirator. The initial evaluation consists of a questionnaire that asks about medical conditions (such as a heart condition) that would be problematic for respirator use. If concerns are identified, then additional evaluations, such as a physical exam, might be necessary. The initial evaluation must be done before respirator use begins. Handlers must be reexamined by a qualified medical practitioner if their health status or respirator style or use conditions change.

Upon request by local/state/federal/tribal enforcement personnel, employers must provide documentation demonstrating how they have complied with these requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer.

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. **DO NOT** discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Sipcam Agro USA, Inc. immediately if you observe any adverse environmental effects due to use of this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use of this product through aerial application equipment on grapes is prohibited in the following townships and boroughs of Erie County, Pennsylvania: North East, Harborcreek, Lawrence Park, Erie, Presque Isle, Millcreek, Fairview, Girard, and Springfield.

This prohibition is intended to help eliminate phytotoxicity problems with apples observed in this geographic location.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval and notification to workers. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area being treated must be vacated by unprotected persons.

DO NOT treat while unprotected humans or domestic animals are present in the treatment areas. Because certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

DO NOT allow entry into treatment area until the area that was treated with this product is dried.

PRODUCT INFORMATION

This product is a broad spectrum, preventative fungicide with systemic and curative properties. This product may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered crop protection products. All applications must be made according to the use directions that follow.

RESTRICTIONS

- **DO NOT** graze or feed clippings from treated turf areas to animals.
- This product is extremely phytotoxic to certain apple varieties. **AVOID SPRAY DRIFT.** Extreme care must be used to prevent injury to apple trees and apple fruit. **DO NOT** spray this product where spray drift may reach apple trees. **DO NOT** use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.
- Aerial and/or chemigation application to sod is prohibited.

SPRAY DRIFT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft. above the ground or crop canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzles that deliver Medium to coarse spray droplets in accordance with ASABE Standard S-572.1.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Groundboom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Airblast Applications:

- Sprays must be directed into the canopy.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer rows.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

- For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

- Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

- Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

- Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

MIXING, LOADING AND APPLYING

This product is intended to be diluted into water and then applied to crops by typical agricultural spraying techniques. Always apply this product in sufficient water to obtain thorough, uniform coverage of foliage and crop surfaces intended to be protected from disease. Spray volume to be used will vary with crop and amount of plant growth. Spray volume must normally range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute ground sprays and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are acceptable unless specific directions are given for a crop.

Slowly invert container several times to assure uniform mixture. Measure the required amount of this product and pour into the spray tank during filling. Keep agitator running when filling spray tank and during spray operations.

It is necessary to thoroughly apply the product in order to provide good disease control. **DO NOT** prepare more spray solution than is needed for application. Avoiding spray overlap will reduce the potential for crop injury.

Tank Mixing

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

When tank mixing this product with other pesticides, observe the more restrictive label limitations and precautions. **DO NOT** exceed any label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

DO NOT combine this product in the sprayer tank with pesticides, surfactants or fertilizers, unless prior use has shown the combination to be physically compatible, non-injurious and effective under similar use conditions. **DO NOT** combine this product with other products that contain *Bacillus thuringiensis* subsp. *kurstaki* strain ABTS-351, as the combination may result in phytotoxicity when applied to the crops listed on this label. **DO NOT** tank mix this product with oil or with any adjuvants which contain oil as their principal ingredient.

When mixed with EC (emulsifiable concentrate) formulations, this product may be phytotoxic to other crops listed on this label, especially when applied during cool, cloudy conditions that last for several days. Adjuvants containing silicone could also have phytotoxic effects. When an adjuvant is used with this product, use a Council of Producers and Distributors of Agrotechnology (CPDA) certified adjuvant.

Applications through Sprinkler Irrigation Systems (Chemigation)

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set and portable (wheel move, side roll, end tow, or hand move) irrigation system(s). **DO NOT** apply this product through any other type of irrigation system. Use only on crops specifically designated in the **Crop Use Directions**.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

DO NOT apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation must be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject this product into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

This product may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a metering pump, such as a positive displacement injection pump of either diaphragm or piston type, constructed of materials that are compatible with pesticides, fitted with a system interlock, and capable of injection at pressures approximately 2 to 3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required, amount of water injected, and acreage covered. Thoroughly mix the specified amount of this product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run but continue to operate irrigation system until this product has been cleared from the last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a 30 – 45 minute period. Mix desired amount of this product for acreage to be covered with water so that the total mixture of this plus water in the injection tank is equal to the quantity of water used during calibration and operate entire system at normal pressures specified by the manufacturer of injection equipment used for the amount of time established during calibration. No agitation should be required. This product can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until this product has been cleared from the last sprinkler head.

INTEGRATED PEST / DISEASE MANAGEMENT

This product provides excellent control of fungal diseases when used according to label directions for control of a broad spectrum of plant diseases. This product is for use in programs that are compatible with the principals of Integrated Pest Management (IPM), including the use of disease resistant crop varieties, cultural practices, pest scouting, and disease forecasting systems which reduce unnecessary applications of pesticides.

RESISTANCE MANAGEMENT

For resistance management, Endow 2SC contains azoxystrobin, a QoI Group 11 fungicide. Any fungal population may contain individuals naturally resistant to Endow 2SC and other QoI Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Fungal isolates with acquired resistance to Group 11 may eventually dominate the fungal population if Group 11 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. Cross resistance has been shown between all members of the QoI fungicides. Since QoI fungicides are a high risk for resistance, this may result in partial or total loss of control of those species.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 11 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance - management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your local Sipcma Agro representative. You can also contact your pesticide distributor or university extension specialist to report resistance.

Follow the crop specific resistance management guidance listed in the Crop Use Directions table. If resistance management guidance is not specified in the Crop Use Directions table, then follow the guidance provided in the table below.

Total fungicide applications planned per crop	1	2	3	4	5	6	7	8	9	10	11	12
Applications of QoI fungicides applied alone	1	1	2	2	2	2	2	3	3	3	3	4
Applications of QoI fungicides applied in mixture (tank-mix or formulated)	1	2	2	2	2	3	3	4	4	5	5	6

When multiple applications are required during the year, spray programs for Group 11 (QoI) fungicides must be developed. When two sequential applications of Group 11 fungicides are made, they must be alternated with two or more applications of a fungicide that is not a Group 11 fungicide. If more than 12 applications are made during the year, observe these guidelines:

- When applying Group 11 (QoI) fungicides alone, the number of applications must not exceed more than 1/3 of the total number of fungicide applications per year.
- When applying Group 11 (QoI) fungicides in tank mixes or premixes with mixing partners of different modes of action, the number of QoI containing applications must not exceed more than ½ of the total number of fungicide applications per year.
- When applying Group 11 (QoI) fungicides both alone and in mixtures, the number of QoI containing applications must not exceed 50% of the total number of fungicide applications per year.

When applying a Group 11 fungicide to seed or soil, wait at least 3 weeks before making another application with a Group 11 fungicide.

TURF

This product is for control of labeled diseases on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Integrated Pest (Disease) Management

Integrate this product into an overall disease and pest management strategy whenever the use of a fungicide is required. Follow cultural practices known to reduce disease development, including varieties with disease resistance, removal of plant debris in which inoculums overwinter, and proper timing and placement of irrigation. Consult your agricultural authorities for additional IPM strategies established for your area. This product may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development.

Resistance Management

Some turf diseases are known to have developed resistance to products used repeatedly for their control. Apply this product in a tank mix or alternation program with other registered fungicides with different modes of action and to which pathogen resistance has not developed.

DO NOT make more than two sequential applications of this product for control of *Pythium* spp. and Gray Leaf Spot. For all other diseases when *Pythium* spp. is not present, **DO NOT** make more than three sequential applications of this product.

Application Directions

Apply this product prior to disease development. Mix with the required amount of water and apply as a dilute spray application in 2-4 gallons of water per 1000 sq. ft. (87–174 gallons per acre). Repeat at specified application intervals until disease is under control or until the maximum treatment rate has been reached. See **Rate Conversion Chart for Turf** for maximum number of applications at each of the specified rates.

For spot treatments, use 0.4 fl.oz. of this product per 1 to 2 gallons of water.

Apply the higher rate and shorter application intervals when prolonged disease conditions exist.

Note: This product does not control Dollar Spot. It is compatible in tank mixes with many other fungicides that do control Dollar Spot. Always tank mix this product with another fungicide that controls Dollar Spot when this disease is present.

Follow directions under Tank Mixes below.

RESTRICTIONS

- Apply by ground (includes any hand-held and mechanically pressurized handwands) only.
- **DO NOT** use more than the maximum allowed use rate down in the "Application Directions for Turf" chart below.
- **DO NOT** apply more than 8.8 quarts of product per acre per year (6.46 fl.oz. product/1000 sq. ft./year) (0.52 lbs./acre/year). See **Rate Conversion Chart for Turf** for maximum number of applications at each of the specified rates.
- **DO NOT** make more than two sequential applications of this product for control of *Pythium* spp. For all other diseases, **DO NOT** make more than three sequential applications of this product.
- Follow the Application Interval requirements in the "Application Directions for Turf" chart below.

Tank Mixes

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

When mixed with EC (emulsifiable concentrate) formulations, this product may be phytotoxic, especially when applied during cool, cloudy conditions that last for several days. Adjuvants containing silicone could also have phytotoxic effects.

APPLICATION DIRECTIONS FOR TURF DISEASES

Target Diseases	Use Rate (fl.oz./1000 sq. ft.)	Application Interval (days)	Application Directions
Anthracnose <i>(Colletotrichum graminicola)</i> Brown Patch <i>(Rhizoctonia solani)</i> Fusarium Patch <i>(Microdochium nivale)</i> Gray Leaf Spot <i>(Pyricularia grisea)</i> Red Thread <i>(Laetisaria fuciformis)</i> Southern Blight <i>(Sclerotium rolfsii)</i> Summer Patch <i>(Magnaporthe poae)</i>	0.38-0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Cool Weather Brown Patch Yellow Patch <i>(Rhizoctonia cerealis)</i>	0.77	28	Make one or two applications in fall or when conditions are favorable for disease development. Minimum Retreatment Interval: 28 days
Fairy Ring <i>(Lycoperdon spp., Agroclybe pediades, and Bovistra plumbea)</i>	0.77	28	Apply this product as soon as possible after fairy ring symptoms develop. Irrigate with at least 0.5 inches water within 24 hours after application if rainfall is not expected. Severely damaged or thin turf may require reseeded. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. Make an additional application 14-28 days if needed. Minimum Retreatment Interval: 28 days
Gray Snow Mold* Typhula Blight <i>(Typhula incarnata, T. ishikariensis)</i>	1.33	Single Application	Make a single application of 1.33 fl.oz. or two applications of 0.67 fl.oz. at 14-day intervals in late fall just before snow cover. When disease pressure is severe, tank mix with another snow mold fungicide to enhance control. Minimum Retreatment Interval: 28 days
	0.67	28	*NOT for use in California.
Leaf Spot <i>(Bipolaris sorokiniana)</i> Melting Out <i>(Drechslera poae)</i>	0.38 – 0.77	14 – 21	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Leaf Rust <i>(Puccinia spp.)</i>	0.38 – 0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Necrotic Ring Spot <i>(Leptosphaeria korrae)</i>	0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Pink Patch <i>(Limonomysces roseipellis)</i>	0.38 - 0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Pink Snow Mold* <i>(Microdochium nivale)</i>	1.33	Single Application	Make a single application of 1.33 fl.oz. or two applications of 0.67 at 14-day intervals in late fall just before snow cover. When disease pressure is severe, tank mix with another snow mold fungicide to enhance control. Minimum Retreatment Interval: 14 days
	0.67	14	*NOT for use in California.
Powdery Mildew <i>(Blumeria graminis)</i>	0.38 – 0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Pythium Blight Pythium Root Rot <i>(Pythium aphanidermatum, Pythium spp.)</i>	0.77	10 – 14	Apply this product before disease is present. During periods of prolonged favorable conditions, treat on the 10 day application interval. This product can be used on newly seeded as well as established turf. Minimum Retreatment Interval: 10 days

(continued)

APPLICATION DIRECTIONS FOR TURF DISEASES *(continued)*

Target Diseases	Use Rate (fl.oz./1000 sq. ft.)	Application Interval (days)	Application Directions
Pythium Root Dysfunction (<i>Pythium volutum</i>)	0.77	21 – 28	Apply preventatively when mean daily soil temperatures are between 55°F and 70°F. Irrigate with 0.1 to 0.2 inches within 24 hours after application to facilitate movement into the root zone. Minimum Retreatment Interval: 21 days
Rhizoctonia Large Patch** (<i>Rhizoctonia solani</i>)	0.38 - 0.77	14 – 28	Make one or two applications in fall or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days **Not for use at the lower rate in California.
Rhizoctonia Leaf Spot (<i>Rhizoctonia zeae</i>)	0.77	14 – 28	Apply when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Spring Dead Spot (<i>Leptosphaeria korrae</i>) or (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>) or (<i>Ophiosphaerella herpotricha</i>)	0.77	28	Make one or two applications in fall or when conditions are favorable for disease development. Minimum Retreatment Interval: 28 days
Take-all Patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)	0.77	28	Make two applications at 28-day intervals in the spring and two applications at 28-day intervals in the fall. Minimum Retreatment Interval: 28 days
Zoysia Patch (<i>Rhizoctonia solani</i> and/or <i>Gaeumannomyces incurstana</i>)	0.38 - 0.77	14 – 28	Make one or two applications in late fall before snow cover or when conditions are favorable for disease development. DO NOT apply on top of snow. Minimum Retreatment Interval: 14 days

Rate Conversion Chart for Turf

Fluid Ounces Product/1000 sq. ft.	Ounces AI /1000 sq. ft.	Fluid Ounces Product/Acre	Pints of Product / Acre	Maximum number of applications at specified rate
0.38	0.100	16.6	1.03	17
0.5	0.130	21.8	1.4	13
0.6	0.156	26.1	1.6	12
0.67	0.174	29.2	1.8	10
0.77	0.200	33.5	2.1	8
1.33	0.34	57.9	3.6	5

Amount to Mix per 100 Gallons for Turf Applications

Spray Volume (gallons/1000 square feet)				
Use Rate (fl.oz.)	2.0 gals.(fl.oz.)	3.0 gals. (fl.oz.)	4.0 gals. (fl.oz.)	
0.38	19	12.7	9.5	
0.5	25	17	13	
0.6	30	20	15	
0.67	33.5	22.4	16.8	
0.77	38.5	25.7	19.3	
1.33	66.5	44.3	33.2	

ORNAMENTALS

This product is for control of certain pathogens causing foliar, aerial, and root diseases, including leaf, tip and flower blights, leaf spots, downy mildew, powdery mildew, anthracnose, and rusts of ornamental plants. Use this product to control certain diseases of container, bench, flat, plug, bed or field-grown ornamentals in greenhouses, shade-houses, outdoor nurseries, retail nurseries and other landscape areas.

Integrated Pest (Disease) Management

Integrate this product into an overall disease and pest management strategy whenever the use of a fungicide is required. Follow cultural practices known to reduce disease development, including selection varieties with disease resistance, optimum plant populations, proper fertilization, winter and/or spring pruning, plant residue management and proper timing and placement of irrigation. Consult your agricultural authorities for additional IPM strategies established for your area. This product may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development.

Application Directions

Apply this product as a broadcast or banded spray, targeting the foliage or crown of the plant. Apply to runoff in sufficient water, ensuring complete coverage of the target plant. Best control of targeted diseases is attained with sufficient coverage and wetting of foliage. Refer to the specific use directions for control of certain diseases. Repeat at specified application intervals (plus alternations for resistance management) until disease is under control or until the maximum treatment rate has been reached. (see **Application Rates** chart below).

Begin applications of this product prior to disease development and continue throughout the season at the specified intervals following resistance management guidelines. Use this product as part of a preventative disease management program.

This product may not provide adequate disease control when applied after disease outbreak.

Application Rates

See charts below for the specific rate range for this product. This product can be applied every 7 – 28 days (or as otherwise specified on this label for a specific plant or disease). Adding a non-silicone based wetting/sticking agent at the specified use rate may enhance coverage on hard-to-wet plant foliage.

Field-grown or Nursery ornamentals (ground and chemigation applications)

Conditions	Rate Range	Retreatment Intervals	Maximum Number of Applications
For most conditions and for most diseases	3.85 – 5.75 fl.oz. / 100 gallons	7 – 14 days	DO NOT apply more than 0.36 gallons (46 fl.oz. or 0.75 lbs.AI) of this product per acre per year or 8 applications per year.
	1.9 – 2.88 fl.oz. / 50 gallons	7 – 14 days	
When disease pressure is light to moderate	1.9 – 2.88 fl.oz. / 100 gallons	7 – 14 days	
	0.95 – 1.44 fl.oz. / 50 gallons	7 – 14 days	
	5.75 fl.oz. / 100 gallons	14 – 28 days	
	2.85 – 2.88 fl.oz. / 50 gallons	14 – 28 days	
When environmental conditions are favorable to severe disease development	5.75 fl.oz. / 100 gallons	7 – 14 days	
	2.85 – 2.88 fl.oz./50 gallons	7 – 14 days	

Nursery ornamentals broadcast application and Landscape (plants, flowers, trees, turf), (any hand-held applications)

Greenhouse ornamentals (mechanically pressurized handwand applications)

Conditions	Rate Range	Retreatment Intervals	Maximum Number of Applications
For most conditions and for most diseases	3.85 – 5.75 fl.oz. / 100 gallons	7 – 14 days	DO NOT apply more than 0.0012 gallons (0.15 fl.oz. or 0.0025 lbs.AI) of this product per acre per year or 8 applications per year.
	1.9 – 2.88 fl.oz. / 50 gallons	7 – 14 days	
When disease pressure is light to moderate	1.9 – 2.88 fl.oz. / 100 gallons	7 – 14 days	
	0.95 – 1.44 fl.oz. / 50 gallons	7 – 14 days	
	5.75 fl.oz. / 100 gallons	14 – 28 days	
	2.85 – 2.88 fl.oz. / 50 gallons	14 – 28 days	
When environmental conditions are favorable to severe disease development	5.75 fl.oz. / 100 gallons	7 – 14 days	
	2.85 – 2.88 fl.oz./50 gallons	7 – 14 days	

Ornamental Uses (except for Field-grown or Nursery ornamentals (ground and chemigation applications), any hand-held applications and mechanically pressurized handwand applications)

Conditions	Rate Range	Retreatment Intervals	Maximum Number of Applications
For most conditions and for most diseases	3.85 – 7.7 fl.oz. / 100 gallons	7 – 14 days	DO NOT apply more than 2.4 gallons (5.0 lbs.Al) of this product per acre per year or 8 applications per year.
	1.9 – 3.85 fl.oz. / 50 gallons	7 – 14 days	
When disease pressure is light to moderate	1.9 – 3.85 fl.oz. / 100 gallons	7 – 14 days	
	0.95 – 1.9 fl.oz. / 50 gallons	7 – 14 days	
	5.75 – 7.7 fl.oz. / 100 gallons	14 – 28 days	
	2.85 – 3.85 fl.oz. / 50 gallons	14 – 28 days	
When environmental conditions are favorable to severe disease development	5.75 – 7.7 fl.oz. / 100 gallons	7 – 14 days	
	2.85 – 3.85 fl.oz. / 50 gallons	7 – 14 days	

Application Rate Conversion Chart

Fl.oz. Rate	Lbs.Al	Fl.oz. Rate	Lbs.Al
0.38 fl.oz.	0.006	3.85 fl.oz.	0.063
0.95 fl.oz.	0.015	5.75 fl.oz.	0.093
1.44 fl.oz.	0.023	7.7 fl.oz.	0.125
1.75 fl.oz.	0.028	15.4 fl.oz.	0.250
1.9. fl.oz.	0.031	30.75 fl.oz.	0.500
2.85 fl.oz.	0.046	46.0 fl.oz.	0.748
2.88 fl.oz.	0.047		

Drench Application

Apply this product as a preventative, drench treatment prior to infection to control soilborne, seedling, and crown diseases of production ornamentals (greenhouse, shadehouse, and container grown). Good coverage of the pre-infection area (root zone, root ball, crown, etc.) is necessary for satisfactory control.

This product may be applied as a drench to container grown ornamentals using 0.38 – 1.75 fl.oz. / 100 gallons of water prior to infection as healthy roots are necessary to optimize product uptake, systemic translocation and disease protection. Apply 1-2 pints of the solution per square foot surface area at 7-28 day intervals.

Drench applications may cause phytotoxicity in small bedding plants in the seedling/plug stage. Test this product on a small number of plants before applying on a larger scale.

Drip Irritation

Apply this product through drip irrigation systems to potted ornamentals or to bedded, field grown ornamentals for soil-borne disease control. Apply at the rate of 3.85 – 30.75 fl.oz. per acre as a preventative disease application. Ensure that the soil or potting media has adequate moisture before making the drip application.

Drip irrigation must be terminated when the fungicide is depleted from the main feed supply tank or 6 hours after starting irrigation, whichever is shorter. Delay a subsequent irrigation (water only) for at least 24 hours following the drip application to ensure maximum efficacy.

ORNAMENTAL USE PRECAUTIONS

- This product may be applied to certain varieties of crabapple for control of apple scab. It has been shown to be safer when applied to the species and varieties listed in Table 3, however, due to the large number of genera, species, and varieties of crabapple, it is impossible to test every one for tolerance to this product. The professional user must conduct small scale testing to ensure plant safety prior to broad scale commercial use on plant genera and species.
- Use only surfactants approved for use on ornamental plants in combination with this product. Prior to broad scale use, test tank mixes on a small group of representative plants.
- This product may not provide adequate disease control when applied after disease outbreak.
- Test this product on a small number of plants before applying on a larger scale.

ORNAMENTAL USE RESTRICTIONS

- For all ornamental Uses (NOT for Field-grown or Nursery ornamentals (ground and chemigation applications), any hand-held applications and mechanically pressurized handwand applications):
 - **DO NOT** apply more than 2.4 gallons (5.0 lbs.Al) of this product per acre per year.
 - **DO NOT** apply more than 8 applications per year.
 - The Maximum Single Application rate is indicated in the chart by ^m following the rate (for example 7.7^m fl.oz.)
 - The Minimum Retreatment Interval is indicated in the chart by ^R following the interval range (for example 7^R – 28 days).
- For Field-grown or Nursery ornamentals (ground and chemigation applications):
 - **DO NOT** apply more than 0.36 gallons (46 fl.oz. or 0.75 lbs.Al) of this product per acre per year.
 - **DO NOT** apply more than 8 applications per year.
- For Nursery ornamentals broadcast application and Landscape (plants, flowers, trees, turf), (any hand-held applications) and Greenhouse ornamentals (mechanically pressurized handwand applications):
 - **DO NOT** apply more than 0.0012 gallons (0.15 fl.oz. or 0.0025 lbs.Al) of this product per acre per year.
 - **DO NOT** apply more than 8 applications per year.

- **DO NOT** apply more than 600 gallons spray volume per acre for foliar applications.
- **DO NOT** apply more than 2 pints per square foot for drench and crown applications.
- Apply by ground (includes any hand-held and mechanically pressurized handwands) **only**.
- **DO NOT** apply this product to apple or cherry trees (flowering, Yoshina variety) due to possible phytotoxicity. **DO NOT** use spray equipment that has applied this product for use on these sensitive crops due to possible phytotoxicity from residue remaining in the sprayer.
- **DO NOT** use silicone based products with this product due to possible phytotoxicity.
- **DO NOT** tank mix this product with other pesticides, fertilizers, adjuvants, etc., unless testing or local knowledge indicates that the tank mixture is safe when used on ornamental plants.
- **DO NOT** make more than three sequential drench applications of this product before alternating with a fungicide of a different mode of action.
- **DO NOT** alternate with other strobilurin fungicides.

TABLE 1 - DISEASES CONTROLLED

When used according to the label directions, this product provides control of the following diseases of ornamental plants:

Target Diseases	Use Rates and Application Directions	
	8 oz. and larger Containers fl.oz. product per 100 gallons	4 oz. Containers fl.oz. product per 50 gallons
1. Conifer Blights		
1a. <i>Phomopsis</i> Blight (<i>Phomopsis juniperovora</i>)	Apply 1.9 – 7.7 ^M fl.oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl.oz. at 7 ^R to 28-day intervals.
1b. Tip Blight (<i>Sirococcus strobilinus</i>)		
2. Leaf Blights/Leaf Spots		
2a. Alternaria Leaf Spot (<i>Alternaria</i> spp.) 2b. Anthracnose (<i>Colletotrichum</i> spp., <i>Elsinoe</i> spp.) 2c. Entomosporium Leaf Spot (<i>Entomosporium mespilii</i>) 2d. Leaf spot (<i>Cladosporium echinulatum</i>) 2e. Downy Mildew of Bedding Plants (<i>Peronospora</i> spp.) 2f. Cercospora Leaf Spot (<i>Cercospora</i> spp.)	Apply 1.9 – 7.7 ^M fl.oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl.oz. at 7 ^R to 28-day intervals.
2h. Marssonina Leaf Spot (<i>Marssonina</i> spp.)	Apply 1.9 – 7.7 ^M fl.oz. at 14 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl.oz. at 14 ^R to 28-day intervals.
2i. Downy Mildew of Rose (<i>Peronospora sparsa</i>)	Apply 3.85 – 7.7 ^M fl.oz. at 7 ^R to 21-day intervals during periods of active plant growth and prior to dormancy or severe infection.	Apply 1.95 – 3.85 ^M fl.oz. at 7 ^R to 21-day intervals during periods of active plant growth and prior to dormancy or severe infection.
2j. Iris Leaf Spot (<i>Mycosphaerella macrospora</i>) 2k. Myrothecium leaf spot (<i>Myrothecium</i> spp.)	Apply 3.85 – 7.7 ^M fl.oz. at 7 ^R to 21-day intervals.	Apply 1.95 – 3.85 ^M fl.oz. at 7 ^R to 21-day intervals.
2l. Rose Blackspot (<i>Diplocarpon rosea</i>)	Apply 7.7 – 15.4 ^M fl.oz. at 7 ^R to 14-day intervals. If disease pressure is light, apply at 7-day intervals. This product may be tank-mixed with another Rose Blackspot fungicide if disease conditions are severe. DO NOT exceed 46 fl.oz. of product per acre.	Apply 3.85 – 7.7 ^M fl.oz. every 7 ^R -14 days. If disease pressure is light, apply at 7-day intervals. This product may be tank-mixed with another Rose Blackspot fungicide if disease conditions are severe. DO NOT exceed 46 fl.oz. of product per acre.
2m. Scab (<i>Venturia inaequalis</i>)	Apply 1.9 – 7.7 ^M fl.oz. at 10 ^R to 28-day intervals. DO NOT apply to apple trees. Refer to Table 3 for tolerant species of crabapples.	Apply 0.95 – 3.85 ^M fl.oz. at 10 ^R to 28-day intervals. DO NOT apply to apple trees. Refer to Table 3 for tolerant species of crabapples.
3. Powdery Mildew		
3a. <i>Erysiphe pannosa</i> , E. spp. 3b. <i>Microsphaera azalea</i> 3c. <i>Sphaerotheca pannosa</i>	Apply 1.9 – 7.7 ^M fl.oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl.oz. at 7 ^R to 28-day intervals.
4. Rusts		
4a. Needle Rust (<i>Melampsora occidentalis</i>) 4b. <i>Phragmidium</i> spp. 4c. <i>Puccinia</i> spp. 4d. <i>Gymnosporangium</i> spp.	Apply 1.9 – 7.7 ^M fl.oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl.oz. at 7 ^R to 28-day intervals.
5. Flower Blights		
5a. Anthracnose (<i>Colletotrichum</i> spp., <i>Elsinoe</i> spp.)	Apply 1.9 – 7.7 ^M fl.oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl.oz. at 7 ^R to 28-day intervals.
5b. Botrytis Slight (<i>Botrytis cinerea</i>) – Suppression only	Apply 7.7 – 15.4 ^M fl.oz. at 7 ^R to 21-day intervals. DO NOT exceed 46 fl.oz. of product per acre.	Apply 3.85 – 7.7 ^M fl.oz. at 7 ^R to 21-day intervals. DO NOT exceed 46 fl.oz. of product per acre.

(continued)

TABLE 1 - DISEASES CONTROLLED (continued)

Target Diseases	Use Rates and Application Directions	
	8 oz. and larger Containers fl.oz. product per 100 gallons	4 oz. Containers fl.oz. product per 50 gallons
6. Shoot/Stem Diseases		
6a. Aerial/Shoot Blight (<i>Phytophthora</i> spp.)	Apply 1.9 – 3.85 ^M fl.oz. at 7 th to 28-day intervals.	Apply 0.95 – 1.9 ^M fl.oz. at 7 th to 28-day intervals.
7. Soilborne Diseases (Directed Spray)	Refer to the Soilborne/Seedling Disease section for application guidelines.	
7a. <i>Rhizoctonia solani</i> 7b. <i>Sclerotium rolfsii</i> 7c. <i>Rosarium</i> spp.	Apply 1.9 – 7.7 ^M fl.oz. at 7 th to 21-day intervals.	Apply 0.95 – 3.85 ^M fl.oz. at 7 th to 21-day intervals.
8. Soilborne Diseases (Drench)	Refer to the Drench Application section above for additional application directions.	
8a. <i>Rhizoctonia solani</i> 8b <i>Sclerotium rolfsii</i> 8c. <i>Fusarium</i> spp.	Apply 0.35 – 1.75 ^M fl.oz., 1 -2 pints of the solution per square foot of surface area, at 7 th to 28-day intervals.	Apply 0.19 – 0.95 ^M fl.oz., 1-2 pints of the solution per square foot of surface area, at 7 th to 28-day intervals.

^M The Maximum Single Application rate is indicated in the chart by ^M following the rate (for example 7.7^M fl.oz.)

^R The Minimum Retreatment Interval is indicated in the chart by ^R following the interval range (for example 7^R – 28 days).

PLANT SAFETY

This product has been shown to be safe when applied to the ornamental plants listed in Tables 2 and 3. However due to the large number of genera, species and varieties of ornamental and nursery plants; it is impossible to test every one for tolerance to this product. Neither the manufacturer nor the seller has determined whether or not this product can be used safely on all genera, species, or varieties of ornamental and nursery plants. The professional user must conduct small scale testing to ensure plant safety prior to broad scale commercial use on plant genera and species not listed in this label.

In addition, **DO NOT** tank mix this product with other pesticides, fertilizers, adjuvants, etc, unless local experience indicates that the tank mix is safe to ornamental plants.

DO NOT apply or use spray equipment used to make applications of this product to certain apple, crabapple or cherry trees and other sensitive crops due to possible phytotoxicity.

Table 2 - Tolerant Ornamental Plants and Diseases Controlled

When applied to the plants listed in Tables 2 and 3 at the listed rates and according to the application directions on this label, this product has been found to be safe and effective at controlling the listed diseases.

Common Name	Botanical Name	Diseases/Pathogens (Refer to Table 1)
Abelia	Abelia spp.	2
Alder (White), Clethra	Clethra alnifolia	2
Arborvitae	Thujaopsis spp.	2
Aspen trees	Populus spp.	2
Aster, Starwort	Aster, spp.	4
Azalea, Glacier	Rhododendron spp.	2b, 3, 6, 7
Azaleas, Rhododendron	Rhododendron spp.	2b, 3, 6, 7
Barberry	Berberis thunbergii	3, 4
Australian Laurel	Pittosporum spp.	3, 4
Baby Rubber-plant	Peperomia spp.	2, 7
Begonia	Begonia spp. (except Reiger begonia)	2, 3
Birch (River)	Betula nigra	3, 4
Black-eyed Susan	Rudbeckia hirta	2
Blanket-Flower	Gaillardia spp.	2
Bougainvillea	Bougainvillea spp.	2
Boxwood	Buxus sempervirens	2, 7a
Bradford's Pear	Pyres calleryana	3
Buddleia, Butterfly Bush	Buddleia davidii	2
Bugle, Bugleweed	Ajuga reptans	3
Burning Bush	Euonymus alatus	2
Caladium	Caladium spp.	7

(continued)

Table 2 - Tolerant Ornamental Plants and Diseases Controlled (continued)

Common Name	Botanical Name	Diseases/Pathogens (Refer to Table 1)
Camellia	Camellia japonica	2
Carnation	Dianthus caryophyllus	3, 4
Cedar (Atlas)	Cedrus atlantica	2, 4
Cedar (Red)	Juniperus virginiana	1a, 4
Cedar, Western Red	Thuja plicata	4
Cedar (White)	Cedrus spp.	2, 4
Cherry	Prunus pumila	2, 5
Chinese evergreen	Aglaonema spp.	2, 4
Chrysanthemums	Chrysanthemum spp.	2, 7c
Cinquefoil	Potentilla spp.	2
Cotoneaster (Creeping)	Cotoneaster adpressus	7
Cotoneaster (Variegated Rockspray)	Cotoneaster horizontalis	7
Crabapple (See Table 3 for variety list)	Malus spp.	2i
Cranesbill	Geranium spp.	5b
Crapemyrtle	Lagerstroemia indica	2, 3
Creeping thyme	Thymus sagahyifam	4
Cyclamen	Cyclamen spp.	7c
Cyperus	Cyperus spp.	1
Cypress (Sawara)	Chamaecyparis pisifera	1
Cypress, Leyland cypress	Chamaecyparis spp.	1
Daisy (Gerber, Transvaal)	Gerbera jamesonii	3
Dogwood	Cornus florida	2b, 3
Dogwood, Pink Dogwood, Flowering Dogwood	Cornus spp.	2b, 3
Dwarf Pampas Grass	Phelans spp.	3
Dumb cane	Dieffenbachia spp.	2
Euonymus (Dwarf Winged)	Euonymus alata	2
Euonymus (Evergreen)	Euonymus japonicas	2
Fatsia (Japanese), Paper-plant	Fatsia japonica	2
Fig	Ficus spp.	2
Fir, Douglas	Pseudotsuga spp.	1, 4
Fir (Fraser)	Abies fraseri	1, 4
Fir (Noble)	Abies procera	1, 4
Floss-flower	Ageratum spp.	3, 4
Forsythia	Forsythia viridissima	2
Foxglove	Digitalis spp.	2, 3
French hydrangea	Hydrangea macrophylla	2, 3
Gardenia	Gardenia jasminoides	3
Geranium	Pelargonium spp.	3, 4, 5b
Grass	Permisetum alopecuriodes	2
Hydrangea	Hydrangea spp.	2, 3
Heather	Erica dareyensis	2
Hibiscus	Hibiscus moscheutos	2, 3
Hemlock	Tsuga spp.	4

(continued)

Table 2 - Tolerant Ornamental Plants and Diseases Controlled (continued)

Common Name	Botanical Name	Diseases/Pathogens (Refer to Table 1)
Hibiscus	Hibiscus rosa-sinensis	2, 3
Holiday cactus	Schlumbergera	2, 7
Holly, Winterberry, Yaupon	Ilex spp.	3
Hosta	Hosta spp.	2
Impatiens, Balsam	Impatiens spp.	2a, 7a
Indian Hawthorn	Phaphiopsisindica	2, 3, 4
Iris (African, Butterfly)	Diets iridiodes	4c
Iris (bulbous, Spanish, Dutch)	Iris xiphium	2e
Ivy (Algerian)	Hedera algeriensis	2
Ivy (English)	Hedera helix	2
Ivy, Swedish Coleus	Plectranthus spp.	2
Japanese Andromeda	Pieris japonica	2, 7
Japanese aucuba, Japanese laurel	Aucuba japonica	7
Juniper	Juniperus procumbens	1a, 4
Juniper	Juniperus scopulorum	1a, 4
Juniper	Juniperus spp.	1a, 4
Larkspur	Delphinium spp.	2
Laurel	Lauras nobilis	3
Lilac (wild)	Ceanothus sanguineus	3
Lily (Asiatic)	Lilium spp.	2
Lily-turf	Liriope muscari	2
Live-forever, House-Leek	Sempervivum spp.	2
Magnolia, Southern	Magnolia grandiflora	2
Magnolia, Saucer	Magnolia soulangiana	2
Magnolia	Magnolia spp.	2
Maple (Japanese)	Acer palmatum	2
Maple (Sugar)	Acer saccharum	2
Marigold	Tagetes spp.	2a
Mock-orange	Philadelphus	3, 4
Muhgo pine	Pinus muhgo	1b, 4
Mugwort, Sagebrush	Artemisia spp.	2
Nandina	Nandina domestica	2
Oak, pin	Quercus palustris	2, 3
Oak, red	Quercus falcate	2, 3
Oleander, Rose-bay	Nerium oleander	2
Orpine, Stonecrop	Sedum spp.	21
Palm, date	Phoenix dactylifera	2, 7
Palm (Parlor)	Chamaedora elegans	7
Palm, Queen	Syagrus romanzollianum	2
Palm, Roebelin's	Phoenix roebelenii	2, 7
Palm (Sago)	Caryota urens	2, 7
Pampas Grass	Cortaderia selloana	3
Peace lily	Spathiphyllum floribundium	2, 7

(continued)

Table 2 - Tolerant Ornamental Plants and Diseases Controlled (continued)

Common Name	Botanical Name	Diseases/Pathogens (Refer to Table 1)
Periwinkle	Vinca spp.	2, 6a
Petunia	Petunia spp.	6a
Philodendron	Philodendron spp.	2j
Phlox	Phlox spp.	3
Pine, Black	Pinus nigra	1b, 4
Pine, Eastern White	Pinus strobes	1b, 4
Pine, Scotch	Pinus silvestris	1, 4
Pine	Pinus spp.	1b, 4
Pink	Dianthus spp.	3, 4
Plum, Flowering; Purple-leaf	Prunes spp.	2, 5
Poinsettia	Euphorbia spp.	2a
Poplar	Populus trichocarpa	4
Pothos	Epipremnum spp.	2
Primrose	Primula spp.	2
Pussy's Foot	Ageratum spp.	3, 4
Redbud (Western)	Cercis occidentalis	2
Red tip photinia	Photinia glabra	2, 3, 4
Ribbon Grass	Setaria spp.	2, 3
Rose	Rosa spp.	2a, 2c, 3c, 4b
Rose of Sharon	Hibiscus syriacus	2, 3
Rosemary (prostrate)	Rosmarinus spp.	2
Rubber-tree, Umbrella-tree	Brassaia actinophylla	2, 7
Sage	Salvia spp.	3, 4j
Snapdragon	Antirrhinum spp.	3, 4
Snowball, Ceanothus, California lilac	Ceanothus spp.	3
Spirea	Spirea budalda	3
Spirea	Spirea japonica	3
Spreading yew	Taxus baccata	7
Spruce, Blue	Picea purtgens	1
Spruce, Norway	Picea abies	1
Spruce, White	Picea glauca	1
Sweet Alyssum	Lobularia maritima	7
Verbena	Verbena spp.	3
Vervain	Verbena spp.	3
Viburnum	Viburnum spp.	2, 3, 4
Vinca	Catharanthus roseus	2
Viola, Pansy	Viola spp.	1, 2
Virginia Willow	Itea virginica	3, 4
Western hemlock	Tsuga heiarophylla	4
Wiegela (Pink)	Wiegela florida	2
Wormwood	Artemisia spp.	2
Yucca	Yucca spp.	7
Zebra Plant	Aphelandra spp.	2
Zinnia	Zinnia sp.	2a, 3

* **DO NOT** exceed 3.85 fl.oz./100 gallons on these species.

TABLE 3. Tolerant Varieties of Crabapple Species (Genus Malus)

Arkansas Black	Eleyi	Mary Potter	Seiboldii
Atrosanguinea	Enterprise	Molten Lava	Selkirk
Baccafa	Evereste	New Centennial	Sentinel
Baccata var. jackii	Eyeiynn	Ormiston Roy	Silver Moon
Baccata var. mandshurica	Floribunda	Pink Satin	Siiverdrift
Callaway	Gloriosa	Prairie Maid	Sinai Fire
Candyment Sargent	Golden Delicious	Prairifire	Spectabfis
Christmas Holly	Golden Raindrops	Profusion	Sugar Tyme
Coronaria	Hopa	Pumila	Van Eseltine
David	Indian Magic	Ralph Shay	White Angel
Dolgo	Island	Red Jade	Williams Pride
Donald Wyman	Katherine	Red Baron	Winter Gold
Dorothea	Lancelot	Sargent	Yellow Delicious
Doubloons	Louisa	Sargentii	Zumi Calocarpa

TABLE 4. Intolerant Plants. DO NOT apply this product to these species or varieties.

COMMON NAME	BOTANICAL NAME
Apple	Malus domestics
Crabapple - Flame variety	Malus spp.
Crabapple - Brandywine variety	Mals spp.
Crabapple - Novamac variety	Malus spp.
Cherry, Flowering - Yoshina variety	Prunus yedoensis
Leatherleaf Fern and Other Ferns for cut foliage	Rumohra adianformis and other species for cut foliage
Privet	Ligusirum spp.

CONIFERS INCLUDING CHRISTMAS TREES, COMMERCIAL PRODUCTION ROSES

Use this product to control diseases on conifers in production (indoor and outdoor) and in landscape situations. Please see the Ornamental Section above for more detailed directions for use in landscape situations.

CONIFERS, INCLUDING CHRISTMAS TREES		
Target Diseases	Rate per Acre (lbs.AI/A)	Application Directions
<p>Diplodia tip blight (<i>Diplodia pinea</i>)</p> <p>Lophodermium Needlecast (<i>Lophodermium pinastr</i>)</p> <p>Swiss Needlecast (<i>Phaeocryptopus gaumannii</i>)</p>	<p>6.0 – 15.5 (0.10 – 0.25)</p>	<p>Apply this product before disease outbreak and continue throughout the season at 7- to 21-day intervals following resistance management guidelines.</p> <p>Apply this product by ground, air or chemigation. If an adjuvant is used, add it at the manufacturer's specified rates.</p> <p>Include this product in an IPM program, which includes alternating fungicides with different modes of action and selections of varieties with disease resistance and removal of plant debris in which inoculum may overwinter.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl.oz. (0.25 lbs.AI) of this product per acre per application. • DO NOT apply more than 123 fluid ounces (2.0 lbs.AI) of this product per acre per year. • DO NOT make more than eight applications of this product per acre per year at the high rate (0.25 lbs.AI). • Minimum Retreatment Interval: 7 days 		

CHRISTMAS TREES (Any hand-held applications)		
Target Diseases	Rate per Acre (lbs.AI/A)	Application Directions
<p>Diplodia tip blight (<i>Diplodia pinea</i>)</p> <p>Lophodermium Needlecast (<i>Lophodermium pinastr</i>)</p> <p>Swiss Needlecast (<i>Phaeocryptopus gaumannii</i>)</p>	<p>0.005 – 0.01 fl.oz. (0.0001 – 0.0002)</p>	<p>Apply this product before disease outbreak and continue throughout the season at 7- to 21-day intervals following resistance management guidelines.</p> <p>If an adjuvant is used, add it at the manufacturer's specified rates.</p> <p>Include this product in an IPM program, which includes alternating fungicides with different modes of action and selections of varieties with disease resistance and removal of plant debris in which inoculum may overwinter.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 0.01 fl.oz. (0.0002 lbs.AI) of this product per acre per application. • DO NOT apply more than 0.08 fluid ounces (0.00125 lbs.AI) of this product per acre per year. • DO NOT make more than eight applications of this product per acre per year at the high rate (0.0002 lbs.AI). • Minimum Retreatment Interval: 7 days • Not for use in California 		

ROSES, COMMERCIAL ROSE PRODUCTION

Target Diseases	Rate per Acre fl.oz./Acre (lbs.AI/A)	Application Directions
Downy Mildew (Peronospora sparsa) Powdery Mildew (Sporotheca pannosa) Rust (Phragmidium mucronatum, P. tubercalutum, and other Phragmidium spp.) Septoria Leaf Spot (Septoria rosea) Alternaria Leaf Spot (Alternaria alternate)	3.0 – 15.5 (0.05 – 0.25)	<p>Apply this product before disease outbreak and continue throughout the year at 7– to 21 day intervals following resistance management guidelines.</p> <p>Apply this product by ground, air or chemigation. If an adjuvant is used, add it at the manufacturer's specified rates.</p> <p>Include this product in an IPM program, which includes alternate fungicides with different modes of action and selection of varieties with disease resistance, proper fertilizer application, winter and/or spring pruning, management of plant residue, and proper irrigation timing and application.</p> <p>Azoxystrobin has been shown to be safe when applied to roses, however, all varieties of roses have not been tested. Test the product first on a smaller scale to ensure its safety prior to making a broadscale application.</p> <p>DO NOT tank mix this product with other pesticides, fertilizers, etc. unless testing or local knowledge indicates that the tank mixture is safe when used on roses.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>

Restrictions:

- **DO NOT** apply more than 15.5 fl.oz. (0.25 lbs.AI) of this product per acre per application.
- **DO NOT** apply more than 123 fluid ounces (2.0 lbs.AI) of this product per acre per year.
- **DO NOT** make more than eight applications of this product per acre per year at the high rate (0.25 lbs.AI).
- **Minimum Retreatment Interval:** 7 days

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

CONTAINER HANDLING [Bulk/Mini-Bulk] [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER HANDLING [Bulk/Mini-Bulk] [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container or pressure rinse promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full of water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

WARRANTY AND LIMITATION OF DAMAGES

CONDITIONS OF SALE: To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. SIPCAM AGRO USA, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED. To the extent consistent with applicable law, SIPCAM AGRO USA, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. SIPCAM AGRO USA, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.

Endow™ 25C

Azoxystrobin Group 11 Fungicide



For control of listed diseases on Turf; Golf courses; Lawns and landscape areas around residential, institutional, public, commercial and industrial buildings; Parks; Recreational areas; Athletic fields; Ornamentals; Conifers including Christmas trees; and Roses, commercial rose production

ACTIVE INGREDIENT:

Azoxystrobin: methyl (E)-2-[2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate* 22.93%

OTHER INGREDIENTS: 77.07%

TOTAL: 100.00%

*IUPAC

Contains 2.08 lbs. Azoxystrobin per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Emergency Phone Numbers: (800) 424-9300 CHEMTREC (transportation and spills)
(800) 222-1222 Poison Control Center

See additional Precautionary Statements and Directions for Use inside booklet.

NET CONTENTS: 2.5 Gallons (9.46 L)

Manufactured for:
SIPCAM AGRO USA, INC.
2525 Meridian Parkway
Durham, NC 27713

2.5G

EPA Reg. No. 60063-59

EPA Est. No.: 60063-GA-001 – Lot number begins with VL

EPA Est. No.: 70815-GA-001 – Lot number begins with CB

EPA Est. No.: 86555-MO-001 – Lot number begins with AF



EPA20210701 (9/21)

**READ THE LABEL
CAREFULLY
BEFORE OPENING
THE CONTAINER**

PEEL BACK BOOK HERE

FUNGICIDE