## MAX POWER ORGANICS TM

# ENVIREPEL-38<sup>TM</sup>

# **BUG KILLER**

# For Use on Crops, Plants and Trees

### **Does Not Persist in the Environment**

#### **ACTIVE INGREDIENTS:**

Clove Oil	12.5%
Garlic	38.0%
INERT INGREDIENTS*	49.5%
Total	100%

\*Water, Citric Acid and Acetic Acid Contains 1.82 lbs. of Garlic Juice Extracts and 0.45 lbs. of Clove oil per Gallon.

# Keep Out of Reach of Children CAUTION

See back panel for additional precautionary statements.

Si usted no entiende la etizueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

**Net Contents: 2.5 Gallons** 

#### Made in the USA

The Contents of this Container will Cover Up to 20 Acres.

# PRECAUTIONARY STATEMENTS & FIRST AID CAUTION

Avoid contact with eyes. If in eyes flush with plenty of water.

#### **Personal Protective Work Clothing**

Applicators and other handlers must wear: Long-sleeved shirt and long pants, shoes plus socks.

#### **DIRECTIONS FOR USE**

#### AGRICULTURAL USE REQUIREMENTS

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR 170. This standard con tains requirements for the protection of agricultural workers on farms, forests, nurse ries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI).

Do not enter or allow workers entry into treated areas during the REI of 12 hours. PPE and work clothing required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, waterproof gloves and shoes plus socks.

For any requirements specific to your State or Tribe, consult the agency respons ible for pesticide regulation.

#### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

**Pesticide Storage:** The optimal storage temperature is between 35°F and 135°F to prevent product degradation. Do not freeze. Do not store in direct sunlight. Do not use or store near heat, open flame or hot surfaces. Keep out of reach of children or anim als. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposa 1 facility.

**Container Disposal:** Triple rinse (or equivalent). The offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out o f smoke.

In the event of a major spill, fire or other emergency, call (760) 944-7922 day or night.

#### **USE RESTRICTIONS**

Do not use this product if it cannot be applied according to the use patterns on this label. Do not let spray mixture stand in tank overnight. For use as a combination Insecticide and Repellent for controlling and repelling certain important insect pests.

#### SITES

**Avocados:** For controlling and repelling Aphids, Fruitflies, Mites, Thrips, Scale, and Whiteflies.

<u>Cereal Grains</u> (Barley, Buckwheat, Corn, Millet, Oats, Popcorn, Rye, Wheat and Wild Rice): For controlling and repelling Aphids, Armyworms, Beetles, Borers, Cutworms, Earworms, Fruitflies, Maggots, Mites, Rootworms, Sawflies, and Wireworms.

<u>Citrus</u> (Calamondin, Grapefruits, Kumquats, Lemons, Limes, Oranges, Tangelos and Tangerines): For controlling and repelling Aphids, Cutworms, Fruitflies, Leafrollers, Mites, Scale, Thrips, Psyllid and Whiteflies.

<u>Cotton:</u> For controlling and repelling Aphids, Armyworms, Boll Weevils, Bollworms, C otton Leafperforator, Cutworms, Fleahoppers, Fruitflies, Loopers, Lygus Bugs, Mites, Saltmarsh Caterpillar, Thrips, Webworms and Whiteflies.

<u>Curcurbit Crops</u> (Cantaloupe, Casaba Melons, Cucumbers, Gherkins, Gourds, Honeydew Melons, Mango Melons, Muskmelons, Pumpkins, Summer Squash, Watermelons and Winter Squash): For controlling and repelling Aphids, Beetles, Fruitflies, Leafhopp ers, Leafminers, Loopers, Mites, Pickleworms, Thrips, Whiteflies and Wireworms.

<u>Deciduous Nursery Stocks (Trees, Shrubs and Flowers) and Greenhouse Crops:</u> For controlling and repelling Aphids, Bagworms, Beetles, Borers, Chafers, Curculios, Cutworms, Fruitflies, Leafhoppers, Leafminers, Leafrollers, Maggots, Mealybugs, Mites, P lant Bugs, Sawflies, Scale, Spittlebugs, Thrips, Webworms and Whiteflies.

**Forage Crops** (Alfalfa, Clovers, Grasses, Timothy and Vetch): For controlling and repelling Aphids, Armyworms, \*Caterpillars, \*Cinch Bugs, Cutworms, Fleas, Japanese Beetles, Leafhoppers, Lygus Bugs, Mites, Spittlebugs, Thrips, Webworms, We evils and Whiteflies. **Kiwi:** For controlling and repelling Aphids, Fruitflies, Japanese Beetles, Loopers, Mealy Bugs, Mites, Thrips, Scale and Whiteflies.

# SITES (Continued)

<u>Nut Trees</u> (Almonds, Beech Nut, Brazil Nuts, Butternut, Chestnuts, Chinquapin, F ilberts, Hickory Nuts, Macadamia Nuts, Pecans, Pistachios and Walnuts): For controlling and repelling Aphids, Beetles, Borers, Bud Moths, Carpenterworms, Casebearers, Catocalas, Curculios, Mites, Gall Wasps, Fruitflies, Leafminers, Phylloxera, Plant Bugs, Sawflies, Scale, Shuckworms, Spittlebugs, Twig Girdlers, Walnut Caterpillars, Webworms, Weevils and Whiteflies.

<u>Ornamentals</u> (Flowers, Shrubs and Trees): For controlling and repelling Aphids, Bagworms, Beetles, Borers, Chafers, Curculios, Cutworms, Fruitflies, Mites, Leafhoppers, Leafminers, Leafrollers, Maggots, Mealybugs, \*Plant Bugs, Sawflies, Scale, \*Spittlebugs, Thrips, \*Webworms and Whiteflies.

Peanuts: For controlling and repelling Aphids, Armyworms, \*Beetles, \*Cutworms, \*Earworms, Leafhoppers, Mit es, \*Rootworms, Thrips and Whiteflies.

**<u>Pineapples:</u>** For controlling and repelling Aphids, Mites, Thrips and Whiteflies.

Root and Tuber Vegetables (Artichokes, Beets, Burdock, Carrots, Cassava, Chicory, Ginger, Ginseng, Horseradish, Parsley, Parsnip, Potatoes, Radish, Rutabaga, Salsify, Sweet Potatoes, Turnips and Yams): For controlling and repelling Armyworms, Aphids, Beetles, Chinch Bugs, Fruitflies, Leafhoppers, Leafminers, Mites, Plant Bugs, Psyllids, Rust Flies, Symphylans, Thrips, Tuberworms, Weevils, Whiteflies, and Wireworms.

**Tobacco:** For controlling and repelling Aphids, Armyworms, Budworms, Mites, Moths, Thrips and Whiteflies.

MIXING DIRECTIONS: Shake container well or stir contents thoroughly before adding **Bug Killer** to the spray tank. Spray tank pH level must be adjusted to a maximum of 6.0 before adding **Bug Killer** to the spray tank. Optimum pH range for tank mix before mixing **Bug Killer** is 5.5 to 6.0. Recommended buffering agents are **Tomahawk II**, citric acid, acetic acid, phosphoric acid or alliphatic polycarboxylate-type materials. Do not buffer with propionic acid or similar products that affect biologically active products. After mixing, ag itate spray tank mix before and during application to prevent settling of product.

**TANK MIXING WITH OTHER PRODUCTS:** Mix a solution of **Bug Killer** with the products to be tank mixed at their recommended label rates in a glass container to determine if there is any separation. It is strongly recommended to try a small test plot if n ew combinations of products are being used for the first time.

**APPLICATION TIMING AND PEST PRESSURE:** Apply **Bug Killer** at the first signs of insect presence. If tank mixing **Bug Killer** with other pesticides, use the recommended label rates of the other pesticides until exact tank mixed product performance is determined. Repeat application timing should be between 10 to 21 days, determined by field scouting reports of pest pressure, plant nutrition and temperature. Do not make more than 20 applications of **Bug Killer** per each growing season (60 applications total per year).

PRE-HARVEST INTERVAL: Do not apply Bug Killer within 24 hours of crop harvest.

**SPRAY TEMPERATURE:** The optimum temperature range to apply **Bug Killer** is between 50°F (10°C) and 90°F (32.2°C). Minimum recommended application temperature is 50°F (10°C). Maximum recommended application temperature is 100°F (37.7°C). For applications above 100°F (37.7°C) higher application spray rates (more than 40 gallons per acre) are recommended with slow drying surfactants after there has been a minimum of a 15°F drop in air temperature from maximum daytime temperature.

**SPRAY ADJUVANTS: Bug Killer** should not be used with dormant spray oils, heavy oils, spreader -stickers or any surfactants that will coat the surface of the plant foliage and prevent absorption of the product. Recommended surfactants like **Invader**, or surfactants made from Methylated seed oil, Ethoxylated seed oil, Methylated seed oil organo silicone (MSO/OS), Ethoxylated seed oil organo silicon e (ESO/OS), standard non-ionic surfactants, crop oil concentrates with a secondary spreading non ionic surfactant, or narrow range paraffinic oils with a secondary spreading non ionic surfactant are recommended to increase product coverage, absorption, and efficacy.

APPLICATION RATES: 16 to 64 ounces per acre [0.85 to 3.4 lb. Garlic Juice Extracts and 0.12 to 0.46 lb. of Clove oil per acre]. Tank Mix Application Rates: By Air: 7 to 20 gallons of spray solution per acre; By Ground Rig: 10 to 135 gallons of spray solution per acre. [California Tank Mix Application Rates: Apply at 7 to 150 gallons of spray solution per acre for row, field, vine and orchard crops.] Applying this product at rates below 1/2 ounce product per gallon of spray solution applied will result in longer periods of time to obtain product performance and efficacy. Do not spray to the point of runoff! Spray tank pH level must be adjusted to a maximum of 6.0 prior to a dding any chemicals. Heavy water rate applications using should use two quarts of Bug Killer I per 150 gallons of buffered water applied.

**APPLICATION METHODS:** Bug Killer can be applied using conventional application equipment, with little or no change to the operation of the equipment. Aerial or ground rigs are the preferred methods. Do not apply this product through any type of pivot irrigation system.

**POLLINATION WARNING: MAY REPEL BEES: Bug Killer** is an insect control and repellent chemistry, and may repel bees. Do not apply **Bug Killer** less than two (2) days prior to pollination.

**RAIN TIMING:** Do not apply if rain is forecasted or overhead water is to occur within 12 hours.

#### CONDITIONS OF SALE AND WARRANTY

**IMPORTANT:** Read the entire *Directions for Use* and the *Conditions of Sale and Warranty* before using this product. If the terms are not acceptable, return the unopened product container at once.

The *Directions for Use* are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently as sociated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of **Cal Crop USA II**, **L.L.C.** or the Seller. All such risks shall be assumed by the Buyer.

Cal Crop USA II, L.L.C. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the *Directions for Use* subject to the inherent risks referred to above. Cal Crop USA II, L.L.C. makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Cal Crop USA II, L.L.C. or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Cal Crop USA II, L.L.C. and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing *Conditions of Sale and Warranty*, which may be varied only by agreement in writing and signed by a duly authorized representative of Cal Crop USA II, L.L.C..

Manufactured by:

CAL CROP USA II, L.L.C. P.O. Box 698 Bonsall, CA 92003

Envirepel, Tomahawk II, Invader, Bug Killer and Max Power Organics are Trademarks of Cal Crop USA II, L.L.C.