

# RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms. For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

# F4092-3

<b>BIFENTHRIN</b>	<b>GROUP</b>	<b>3A</b>	<b>INSECTICIDE</b>
<b>B. VELEZENSIS STRAIN RTI301</b> <b>B. SUBTILIS STRAIN RTI477</b>	<b>GROUP</b>	<b>NC</b>	<b>FUNGICIDES</b>



For protection against listed soil insects and diseases in agricultural crops.

**EPA Reg. No. 279-9651**

**EPA Est. No. 279-NY-1**

Active Ingredients:	By Wt.
Bifenthrin *:	15.7%
<i>Bacillus velezensis</i> strain RTI301 **:	2.5%
<i>Bacillus subtilis</i> strain RTI477 **:	2.5%
<b>Other Ingredients:</b>	<b>79.3%</b>
	<b>100.0%</b>

\*Cis isomers 97% minimum, trans isomers 3% maximum

\*\* Contains a minimum of 5.0 x 10<sup>9</sup> colony-forming units (cfu) per milliliter of each strain.

This product contains 1.5 lbs. bifenthrin per gallon and a total of 0.48 lbs. per gallon of the *Bacillus* strains.

## KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)**

See other panels for additional precautionary information.

FIRST AID	
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"> <li>- Take off contaminated clothing.</li> <li>- Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>- Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>- Call a poison control center or doctor immediately for treatment advice.</li> <li>- Have person sip a glass of water if able to swallow.</li> <li>- Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>- Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"> <li>- Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>- Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Inhaled:</b>	<ul style="list-style-type: none"> <li>- Move person to fresh air.</li> <li>- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>- Call a poison control center or doctor for treatment advice.</li> </ul>
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
This product contains a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.	

**Net Contents: 2.5 Gallons**

**Sold by:**



FMC Corporation  
2929 Walnut Street  
Philadelphia PA 19104

**Batch/Lot code:** \_\_\_\_\_

**[For nonrefillable containers only. Will be located on this label or on the physical container.]**

# PRECAUTIONARY STATEMENTS

## Hazards to Humans and Domestic Animals

### WARNING

Causes skin irritation. Do not get on skin or on clothing. Wear the Personal Protective Equipment listed below. Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid contact with eyes. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### Personal Protective Equipment (PPE)

Applicators, and other handlers must wear:

- Coveralls worn over short-sleeved shirt and short pants
- Waterproof gloves
- Chemical-resistant footwear
- Socks
- Protective eyewear

Mixers, Loaders and Applicators must wear a minimum of:

A NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air-purifying respirator with an HE filter. (Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.)

When mixing and loading and when cleaning equipment, wear a chemical-resistant apron.

### User Safety Requirements

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meet the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d) and (e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

### User Safety Recommendations

#### Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove clothing/PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not make applications when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. **Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms.**

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county, contact the local extension service for procedures and precautions to use to protect endangered species.

### DIRECTIONS FOR USE

#### Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

F4092-3 can be applied in-furrow at plant, pre-plant incorporated, pre-emergent, and lay-by. Do not make foliar applications. Do not apply this product through any type of irrigation system.

## Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls worn over short-sleeved shirt and short pants, Waterproof gloves, Protective eyewear, Chemical-resistant footwear and socks.

## Resistance Management

For resistance management, F4092-3 contains a Group 3A insecticide. Any insect population may contain individuals naturally resistant to F4092-3 and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of F4092-3 or other Group 3A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - o Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - o Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - o When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pests.
  - o Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - o The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticidal activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

To reduce the potential for pesticide resistance, use this product in a rotation program with other classes of chemistry and modes of action. Always apply this product at the recommended rates and in accordance with the use directions.

## Application and Mixing Instructions

Shake well before using.

F4092-3 is an insecticide/fungicide that contains 1.5 pounds per gallon of the insecticide bifenthrin together with the biopesticides *Bacillus velezensis* strain RT1301 and *Bacillus subtilis* strain RT1477 each at a minimum of  $5.0 \times 10^9$  colony-forming units (cfu) per milliliter. *Bacillus velezensis* strain RT1301 and *Bacillus subtilis* strain RT1477 are naturally occurring beneficial bacteria used for suppression of fungal plant diseases. For suppression of diseases, F4092-3 must be applied in-furrow with the seed. F4092-3 can be mixed directly with liquid fertilizer or with water. The rate of application is variable according to pest pressure, timing of treatments and field scouting. Use lower rates under light to moderate pest infestations, and higher rates under heavier pest pressure. In arid climates application rates are generally higher. Fill the tank one-half full with liquid fertilizer or water and begin spray tank agitation. Add the proper amount of F4092-3, and then add the rest of the fertilizer or water. Maintain agitation until the mixture has been applied.

Agitate the F4092-3 spray solutions in nurse tanks prior to moving the solution to the spray system.

Apply in a minimum of 5 gallons of finished spray per acre unless indicated otherwise under crop specific use directions on this label.

## VEGETATIVE FILTER STRIPS

Construct and maintain a vegetative filter strip, according to the width specified below, of grass or other permanent vegetation between the field edge and nearby down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; streams; marshes; or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing bifenthrin onto fields where a maintained vegetative filter strip of **at least 25 feet** exists between the field edge and where a down gradient aquatic habitat exists. This minimum required width of 25 feet may be reduced or removed under the following conditions:

- For Western irrigated agriculture, a maintained vegetative filter strip of at least 10 feet wide is required. Western irrigated agriculture is defined as irrigated farmland in the following states:  
WA, OR, CA, ID, NV, UT, AZ, MT, WY, CO, NM, and TX (west of I-35).
  - o For Western irrigated agriculture, if a sediment control basin is present, a vegetative filter strip is not required.
- In all other areas, a vegetative filter strip with a minimum width of 25 feet is required, unless the following conditions are met. The vegetative filter strip requirement may be reduced from 25 feet to 15 feet if at least one of the following applies:
  - o The area of application is considered prime farmland (as defined in 7 CFR § 657.5)
  - o Conservation tillage is being implemented on the area of application. Conservation tillage is defined as any system that leaves at least 30% of the soil surface covered by residue after planting. Conservation tillage practices can include mulch-till, no-till, or strip-till.
  - o A functional terrace system is maintained on the area of application.
  - o Water and sediment control basins for the area of application are functional and maintained.
  - o The area of application is less than or equal to 10 acres.

For further guidance on vegetated filter strips, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services.

<https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0331-0175>

F4092-3 can be applied in-furrow with the seed, as a T-band (band over the open furrow), as a broadcast application, as a band over the row or as a transplant-water drench during setting. Refer to the individual crop use directions for pest control or suppression instructions.

F4092-3 can be mixed with commonly used liquid starter or pop-up fertilizers. Follow liquid fertilizer recommendations regarding seed safety and use guidelines. Conduct a preliminary jar test using the appropriate ratio of fertilizer and F4092-3. Do not allow a tank mixture to set overnight, but if this occurs agitate tank mixture prior to application.

Use rates in the individual crop sections listed as Fluid oz/1000 Linear feet are based on 30-inch row spacings. For conversion to rates applicable to other row spacings, see the table below.

<b>F4092-3 Required Per Acre for Various Row Spacings</b>					
<b>Row Spacing</b>	<b>36"</b>	<b>30"</b>	<b>20"</b>	<b>15"</b>	<b>Twin Row 30" centers</b>
Linear row ft/acre	14,520 ft	17,424 ft	26,136 ft	34,848 ft	34,848 ft
<b>Conversion</b>					
0.2 Fluid oz/1000 Linear ft =	2.9 fl oz/A	3.5 fl oz/A	5.2 fl oz/A	7.0 fl oz/A	7.0 fl oz/A
0.24 Fluid oz/1000 Linear ft =	3.5 fl oz/A	4.2 fl oz/A	6.3 fl oz/A	8.4 fl oz/A	8.4 fl oz/A
0.3 Fluid oz/1000 Linear ft =	4.4 fl oz/A	5.2 fl oz/A	7.8 fl oz/A	10.5 fl oz/A	10.5 fl oz/A
0.39 Fluid oz/1000 Linear ft =	5.7 fl oz/A	6.8 fl oz/A	10.2 fl oz/A	13.6 fl oz/A	13.6 fl oz/A
0.49 Fluid oz/1000 Linear ft =	7.1 fl oz/A	8.5 fl oz/A	12.8 fl oz/A	17.1 fl oz/A	17.1 fl oz/A
0.73 Fluid oz/1000 Linear ft =	10.6 fl oz/A	12.7 fl oz/A			
0.78 Fluid oz/1000 Linear ft =	11.3 fl oz/A	13.6 fl oz/A			
0.98 Fluid oz/1000 Linear ft =	14.2 fl oz/A	17.1 fl oz/A			
1.47 Fluid oz/1000 Linear ft =	21.3 fl oz/A	25.6 fl oz/A			

## Crop Rotation Restrictions

Crops for which bifenthrin tolerances exist may be rotated at any time. All other crops may be rotated 30 days following an application of F4092-3.

## Tank-Mixtures

F4092-3 may be applied in tank mixtures with other products approved for use on registered crops. Observe all restrictions and precautions which appear on the labels of these products. To ensure successful applications, product compatibility tests should be conducted.

## Maximum Allowable F4092-3 Use per Acre per Year

Refer to the individual crop sections for maximum allowable F4092-3 usage per acre per year. The maximum allowable use for the bifenthrin active ingredient must include all registered use patterns including at-plant, soil applied and/or foliar applications of all bifenthrin containing products for a 12-month period. The 12-month period is to begin upon the initial application to the acre.

## BUFFER ZONES TO WATER BODIES

### Ground Application

- Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

In New York State this product may not be applied within 100 feet (using ground equipment) of coastal marshes or streams that drain into coastal marshes.

## Mandatory Spray Drift Management

### Ground Boom Applications:

- User must only apply with the nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to select nozzle and pressure that deliver medium or coarser droplets (ASABE S572).
- Do not apply when wind speeds exceed 15 mph at the application site.
- Do not apply during temperature inversions.

### Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### BOOM HEIGHT – Ground Boom

- For ground equipment, the boom should remain level with the crop and have minimal bounce.

### SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### TEMPERATURE AND HUMIDITY

- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### TEMPERATURE INVERSIONS

- Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicate an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### WIND

- Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### NON-TARGET ORGANISM ADVISORY STATEMENT (Environmental Hazards):

- This product is highly toxic to bees and other pollinating insects exposed to direct treatment or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.

## CANOLA, CRAMBE, RAPESEED

### At-Plant

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound bifenthrin/acre	
Army cutworm, Armyworm species, Cutworm species, Grape colaspis, Grubs, Root aphids, Seedcorn beetle, Seedcorn maggot, Stalk borer, Sugarcane beetle, True armyworm, Wireworm  <sup>1</sup> Suppression of: "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , or <i>Phytophthora</i>	6.83	0.08	Apply as a 5 to 7-inch band (T-band) over an open furrow, or in-furrow with the seed. For suppression of diseases, F4092-3 must be applied in-furrow with the seed.

<sup>1</sup> Under moderate to severe disease pressure, or when environmental conditions are conducive to rapid disease development, mix F4092-3 with other fungicides labeled for this use for improved performance.

#### Canola, Crambe, and Rapeseed Restrictions:

- Do not apply more than 0.08 pound bifenthrin per acre per year including PPI, at-plant, PRE, and foliar applications of this and other bifenthrin containing products.

## CORN

Field Corn, Popcorn, Field Corn Grown for Seed, Sweet Corn, Sweet Corn Grown for Seed

### At-Plant

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid oz/1000 Linear ft. <sup>1</sup>	Pound bifenthrin/acre	
Corn rootworm larvae (Northern, Southern, and Western)	0.49 – 0.98	0.10 – 0.20	Apply as a 5 to 7-inch band (T-band) over an open furrow, or in-furrow with the seed.
Army cutworm, Armyworm species, Cutworm species, Grape colaspis, Grubs, Root aphids, Seedcorn beetle, Seedcorn maggot, Stalk borer, Sugarcane beetle, True armyworm, Wireworm  <sup>2</sup> Suppression of: "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , or <i>Phytophthora</i>	0.20 – 0.98	0.04 – 0.20	For Army cutworm, Stalk borer, Cutworm species, True armyworm or Armyworm species, apply as a 5 to 7-inch band over the row on the soil surface, a 5 to 7-inch band over the open furrow (T-band), in-furrow with the seed, or broadcast to the soil surface. For suppression of diseases, F4092-3 must be applied in-furrow with the seed.  Heavy Corn Rootworm Pressure Management Program: In areas where large corn rootworm populations are present, a multi-approach system may be needed for optimal pest management. However, if the population level is not known and if a corn rootworm adult scouting program along with threshold adult control measures were not completed during the previous growing season, then utilize a maximum dosage seed treatment program or genetically modified corn rootworm resistant hybrid in addition to F4092-3.  <sup>1</sup> See rate table on page [4] for amount of F4092-3 required per acre for various crop row spacings.

<sup>2</sup> Use highest application rate for better disease management; however, under moderate to severe disease pressure, or when environmental conditions are conducive to rapid disease development, mix F4092-3 with other fungicides labeled for this use for improved performance.

## Pre-Plant Incorporated (PPI) & Pre-Emergence (PRE)

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound bifenthrin/acre	
Cutworm species, Grape colaspis, White grubs, Wireworm, Seedcorn maggot, Armyworm species, Seedcorn beetle	<b>PPI</b> 4.0 – 5.3	<b>PPI</b> 0.047 – 0.062	F4092-3 can be tank mixed and applied with PPI herbicides. Incorporation of F4092-3 should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth.
Cutworm species, Armyworm species, Stalk borer, Seedcorn beetle	<b>PRE</b> 3.4	<b>PRE</b> 0.04	F4092-3 can be tank mixed and applied with PRE herbicides.

### Corn Restrictions

- Do not apply more than 0.2 pound bifenthrin per acre per season as an at-plant application.
- For field corn: Do not apply more than 0.3 pound bifenthrin per acre per year including PPI, at-plant, PRE, and foliar applications of this and other bifenthrin products.
- For sweet corn: Do not apply more than 0.2 pound bifenthrin per acre per year including PPI, at-plant, PRE, and foliar applications of this and other bifenthrin products.

## COTTON

### At-Plant

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid Linear ft. <sup>1</sup>	Pound bifenthrin/acre	
Cutworm species, White grubs, Wireworm, Grape colaspis, Root maggot, Seedcorn maggot  <sup>2</sup> Suppression of "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , or <i>Phytophthora</i>	0.10 – 0.49	0.02 – 0.10	Apply as a 5 to 7-inch band (T-band) over an open furrow, or in-furrow with the seed. For suppression of diseases, F4092-3 must be applied in-furrow with the seed.  <sup>1</sup> See rate table on page [4] for amount of F4092-3 required per acre for various crop row spacings.

<sup>2</sup> Use highest application rate for better disease management; however, under moderate to severe disease pressure, or when environmental conditions are conducive to rapid disease development, mix F4092-3 with other fungicides labeled for this use for improved performance.

### Cotton Restrictions:

- Do not apply more than 0.5 pound bifenthrin per acre per year including PPI, at-plant, PRE, and foliar applications of this and other bifenthrin containing products in all states except in California.
- For California, do not apply more than 0.3 pound bifenthrin per acre per year including PPI, at-plant, PRE, and foliar applications of this and other bifenthrin containing products.
- Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include but are not limited to Ambush®, Ammo®, Asana® XL, Baythroid®, Baythroid XL®, Brigade®, Capture®, Danitol®, Declare®, Discipline®, Fanfare®, Karate®, and Mustang®.

## DRIED BEANS AND PEAS (excluding Soybean) - Crop Subgroup 6C

Dried cultivars of bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (*Vigna* spp.), (includes adzuki bean, blackeyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar, lablab bean (hyacinth bean); lentil; pea (*Pisum* spp.), (includes field pea, pigeon pea, purple hulled peas)

### At-Plant

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid Linear ft. <sup>1</sup>	Pound bifenthrin/acre	
Army cutworm, Armyworm species, Cutworm species, Grape colaspis, Grubs, Root maggot, True armyworm, Wireworm  <sup>2</sup> Suppression of: "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , or <i>Phytophthora</i>	0.20 – 0.49	0.04 – 0.10	Apply as a 5 to 7-inch band over the row on the soil surface, 5 to 7-inch band (T-band) over an open furrow, or in-furrow with the seed. Apply broadcast to the soil surface for control of Army cutworm, Cutworm species, True armyworm, or Armyworm species. For suppression of diseases, F4092-3 must be applied in-furrow with the seed.  <sup>1</sup> See rate table on page [4] for amount of F4092-3 required per acre for various crop row spacings.

<sup>2</sup> Use highest application rate for better disease management; however, under moderate to severe disease pressure, or when environmental conditions are conducive to rapid disease development, mix F4092-3 with other fungicides labeled for this use for improved performance.

### Dried Beans and Peas (excluding Soybean)- Crop Subgroup 6C Restrictions:

- Do not apply more than 0.2 pound bifenthrin per acre per year to peas or 0.3 pound bifenthrin per acre per year to beans including PPI, at-plant, PRE, and foliar applications of this and other bifenthrin containing products.

## PEANUT

### At-Plant

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid Linear ft. <sup>1</sup>	Pound bifenthrin/acre	
Thrips, Leafhoppers, Aphids, Wireworms  <sup>2</sup> Suppression of: "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , or <i>Phytophthora</i>	0.39 – 0.49	0.08 – 0.10	Apply as a 5 to 7-inch band (T-band) over an open furrow, or in-furrow with the seed. For control/suppression of diseases, F4092-3 must be applied in-furrow with the seed.  <sup>1</sup> See rate table on page [4] for amount of F4092-3 required per acre for various crop row spacings.

<sup>2</sup> Use highest application rate for better disease management; however, under moderate to severe disease pressure, or when environmental conditions are conducive to rapid disease development, mix F4092-3 with other fungicides labeled for this use for improved performance.

### Peanut Restrictions:

- Do not apply more than 0.5 pound bifenthrin per acre per year including PPI, at-plant, PRE, and foliar applications of this and other bifenthrin containing products.



## SOYBEANS

### At-Plant

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid Linear ft. <sup>1</sup>	Pound bifenthrin/acre	
Wireworms, Grape colaspis, Grubs, Root maggot, Seedcorn maggot, Army cutworm, Cutworm species, True armyworm, Armyworm species, Seedcorn beetle  <sup>2</sup> Suppression of: "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , or <i>Phytophthora</i>	0.20 – 0.49	0.04 – 0.10	Apply as a 5 to 7-inch band over the row on the soil surface, a 5 to 7-inch band over the open furrow (T-band), or in-furrow with the seed. Apply broadcast over the soil surface for control of Army cutworm, Cutworm species, True armyworm, or Armyworm species For control/suppression of diseases, F4092-3 must be applied in-furrow with the seed.  <sup>1</sup> See rate table on page [4] for amount of F4092-3 required per acre for various crop row spacings.

<sup>2</sup> Use highest application rate for better disease management; however, under moderate to severe disease pressure, or when environmental conditions are conducive to rapid disease development, mix F4092-3 with other fungicides labeled for this use for improved performance.

### Pre-Plant Incorporated (PPI) & Pre-Emergence (PRE)

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound bifenthrin/acre	
Armyworm species, Black cutworm, Dingy cutworm, Variegated cutworm	<b>PRE</b> 3.4	<b>PRE</b> 0.04	F4092-3 can be tank mixed and applied with PRE herbicides.
Black cutworm, Dingy cutworm, Variegated cutworm, Armyworm species, Stalk borer, Seedcorn beetle	<b>PPI</b> 4.0 – 5.3	<b>PPI</b> 0.047 – 0.062	F4092-3 can be tank mixed and applied with PPI herbicides. Incorporation of F4092-3 should not be any deeper than the in- tended planting depth and no deeper than 3 inches. Incorpora- tion depth should be close to the intended seed planting depth.

#### Soybean Restrictions:

- Do not apply more than 0.3 pound bifenthrin per acre per year including PPI, at-plant, PRE, and foliar applications of this and other bifenthrin containing products.

## SUCCULENT PEAS AND BEANS - Crop Subgroups 6A and 6B

Bean (*Phaseolus* spp.) (includes lima bean (green), broad bean (succulent), runner bean, snap bean, wax bean); bean (*Vigna* spp.) (includes asparagus bean, blackeyed pea, Chinese longbean, cowpea, moth bean, southern pea, yardlong bean); jackbean; pea (*Pisum* spp.) (includes dwarf pea, edible-podded pea, English pea, garden pea, green pea, snow pea, sugar snap pea); pigeon pea; soybean (immature seed); sword bean

### At-Plant

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid Linear ft. <sup>1</sup>	Pound bifenthrin/acre	
Army cutworm, Armyworm species, Cutworm species, Grape colaspis, Grubs, Root maggot, Seedcorn maggot, True armyworm, Wireworm  <sup>2</sup> Suppression of: "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , or <i>Phytophthora</i>	0.20 – 0.49	0.04 – 0.10	Apply as a 5 to 7-inch band over the row on the soil surface, a 5 to 7-inch band over the open furrow (T-band), or in-furrow with the seed. Apply broadcast over the soil surface for control of Army cutworm, Cutworm species, True armyworm, or Armyworm species. For control/suppression of diseases, F4092-3 must be applied in-furrow with the seed.  <sup>1</sup> See rate table on page [4] for amount of F4092-3 required per acre for various crop row spacings.

<sup>2</sup> Use highest application rate for better disease management; however, under moderate to severe disease pressure, or when environmental conditions are conducive to rapid disease development, mix F4092-3 with other fungicides labeled for this use for improved performance.

### Succulent Peas and Beans – Crop Subgroups 6A and 6B Restrictions:

- Do not apply more than 0.2 pound bifenthrin per acre per year including PPI, at-plant, PRE, and foliar applications of this and other bifenthrin containing products.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

### Pesticide Storage

Store in a cool, dry place and avoid excess heat. Do not freeze. Shake or roll the container to re-dissolve solids after storage and before use. Keep out of reach of children and animals. Store in original containers only. Carefully open containers. After partial use, replace lid of container and close tightly. Do not put concentrate or dilute material into food or drink containers. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call CHEMTREC (Transportation and Spills) at 1-800-424-9300. To confine spill, dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. If a leaky container must be contained within another, mark the outer container to identify the contents.

### Pesticide Disposal

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

### Container Handling

**U-Turn® Container:** Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

**Non-refillable Plastic Containers (in sizes 5 gallons or less):** Non-refillable container. Do not reuse or refill this container.

Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds, pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer container for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, by incineration, or by other procedures approved by State and local authorities.

**Non-refillable container (in sizes greater than 5 gallons):** Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer container for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, by incineration, or by other procedures approved by State and local authorities.

OR

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer container for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, by incineration, or by other procedures approved by State and local authorities.

**Returnable/Refillable Containers (if other than U-Turn Container):** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## Conditions of Sale and Limitation of Warranty and Liability:

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. To the extent consistent with applicable law, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and Buyer assumes the risk of any such use.**

To the extent consistent with applicable law, FMC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC Trademark of FMC Corporation.

© 2021 FMC Corporation. All rights reserved.

**SL-4081 091421 07-27-21**