



WILBUR-ELLIS®
Ideas to Grow With™

GUARANTEED ANALYSIS

Total Nitrogen (N).....	2.00%
0.33% Ammoniacal Nitrogen	
1.67% Urea Nitrogen	
Available Phosphate (P ₂ O ₅).....	17.00%
Soluble Potash (K ₂ O).....	17.00%
Iron (Fe).....	0.10%
0.10% Chelated Iron	
Manganese (Mn).....	0.08%
0.08% Chelated Manganese	
Molybdenum (Mo).....	0.0005%
Zinc (Zn).....	0.15%
0.15% Chelated Zinc	

DERIVED FROM: Ammonium Hydroxide, Urea, Phosphoric Acid, Potassium Hydroxide, Iron EDTA, Iron IDS, Manganese EDTA, Manganese IDS, Zinc EDTA, Zinc IDS, and Sodium Molybdate.

CAUTION: This product is to be used only on crops which respond to molybdenum. Crops high in molybdenum are toxic to grazing animals (ruminants).

KEEP OUT OF REACH OF CHILDREN PRECAUTION

Do not ingest. Avoid contact with skin, eyes or clothing.
Avoid breathing dust, vapor or mist.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfc.org/metals.html>

FIRST AID

In all cases, call a poison control center or doctor for further treatment advice.

IF SWALLOWED, call a poison control center or doctor immediately. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. **IF ON SKIN**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF INHALED**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration. **IF IN EYES**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.

FIELD CROPS: Apply 1-8 quarts per acre during peak vegetative growth and repeat during reproductive growth.

GREENHOUSE CROPS: Apply 1-2 pints in not less than 100 gallons of water as needed.

GRASSES GROWN FOR HAY OR SEED (Including Bluegrass, Timothy, Bromes, and Fescues): Apply 2-8 quarts per acre in early spring during vegetative growth, then repeat just prior to head emergence. Use of a high quality liquid nitrogen source such as the NDemand® line of products or other suitable sources may enhance uptake.

LEGUMES: Apply 1-3 quarts per acre during peak vegetative growth and repeat at first bloom.

SMALL GRAINS: Apply 1-8 quarts per acre during peak vegetative growth and repeat at panicle initiation.

VEGETABLES AND CUCURBITS: Apply 1-8 pints per acre at 7 to 10-day intervals, especially during the main growth period.

POTATOES: Apply 1-2 gallons per acre at 7 to 10-day intervals, especially during the main growth period.

For chemical spills, leaks, fire or exposure, call CHEMTREC: (800) 424-9300.

SOIL APPLICATIONS

FOLI-GRO ACTION is a premium quality, low salt index fertilizer that can be applied to seeded crops at 1-5 gallons per acre. Apply to seed zone at time of planting.

Bluegrass, canola, rape, mustard, soybeans, and other small seed crops should have **FOLI-GRO ACTION** placed no closer than 2 inches from the seed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Store in a safe manner. Store in original container only and keep tightly sealed when not in use. Dispose of unused product and empty containers in accordance with federal, state and local regulations.

COMPATIBILITY

FOLI-GRO ACTION is compatible with many co-applied nutrient sources as well as plant protection products. Always jar test new combinations for compatibility prior to field mixing. Do not use **FOLI-GRO ACTION** with oil and lime sulfur. For specific crop recommendations, consult with the Wilbur-Ellis representative in your area.

GENERAL INFORMATION

FOLI-GRO ACTION® is formulated primarily for foliar applications to prevent or correct micronutrient deficiencies in a wide range of agronomic and ornamental plants. Its use is suggested as a supplement to a regular, balanced fertilizer program to enhance yields and improve quality. Application of **FOLI-GRO ACTION** is a means of obtaining a quick response to needed elements. Foliar applications of **FOLI-GRO ACTION** may be particularly beneficial during periods of peak nutrient demand, for crops grown on soils having poor nutrient availability or to crops suffering from a weakened root system.

Avoid foliar nutrient application to crops during periods of moisture, salinity or heat stress. Applications of foliar nutrients to sensitive crops such as leafy vegetables or soft-skinned fruit during periods of excessive temperatures may cause cosmetic injury. **FOLI-GRO ACTION** should be used as part of a comprehensive Total Nutrition System® for optimizing plant growth, development, yield and quality.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of the product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS;" AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Buyer and User accept all risks arising from any use of this product, including without limitation uses contrary to label instructions, under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

To the extent permitted by law, neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS logo, Ideas to Grow With, ACTION, FOLI-GRO, NDemand, R-11, SYL-COAT, and Total Nutrition System are registered trademarks, and FOLI-GRO logo is a trademark of Wilbur-Ellis Company LLC. K-083018

DIRECTIONS FOR USE

FOLI-GRO ACTION may be applied as a foliar spray by air or by ground. Maintain agitation to provide uniform distribution within the spray tank. Thorough and even spray coverage is essential for maximum nutrient uptake and crop safety. Always include a high quality wetting agent, such as R-11® or SYL-COAT®, at recommended rates. Optimum rate of application will vary based on a variety of factors including degree of need or deficiency, growth stage, yield potential, environmental conditions, existing soil fertility levels, etc.

CONVENTIONAL SPRAYER: Apply the recommended amount of **FOLI-GRO ACTION** with a minimum of 20 gallons of water per acre.

CONCENTRATE OR AERIAL: Usually 5 gallons of water per acre is sufficient to apply the recommended amount of **FOLI-GRO ACTION**. If less water is used, a slight burning of the foliage may occur. Aerial applications should not exceed 1 pint per gallon of water.

FOLIAR APPLICATIONS

CANOLA, MUSTARD, RAPE: Apply 1-4 quarts per acre during vegetative growth. Repeat at first bloom.

CITRUS: Apply 1-3 gallons per acre at 7 to 10-day intervals.

DECIDUOUS TREE CROPS AND VINES: Apply 2-8 quarts as needed from bud swell through post-harvest.

Manufactured by: **WILBUR-ELLIS COMPANY LLC**
7 E. Washington Ave., Yakima, WA 98903

LOT#

NET CONTENTS: 2.5 GALLONS (9.46 Liters)
WEIGHT PER GALLON: 11.63 POUNDS at 68°F