

Specimen Label

MANCOZEB	GROUP	M3	FUNGICIDE
----------	-------	----	-----------



Fore[®] 80WP Rainshield[®]

SPECIALTY FUNGICIDE

™®Trademarks of Corteva Agriscience and its affiliated companies

Product in Water-Soluble Packets

Active Ingredients

Mancozeb: A coordination product of zinc ion and manganese ethylene bisdithiocarbamate	80.0%
in which the ingredients are:	
Manganese ⁺⁺	16.00%
Zinc ⁺⁺	2.00%
Ethylene bisdithiocarbamate ion (C ₄ H ₆ N ₂ S ₄)	62.00%
Other Ingredients	20.0%
Total	100.0%

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Precautionary Statements

Hazards to Humans and Domestic Animals

EPA Reg. No. 62719-388

Keep Out of Reach of Children

CAUTION

Harmful If Absorbed Through The Skin • Causes Moderate Eye Irritation • Avoid contact with skin, eyes or clothing • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are butyl rubber and neoprene rubber.

Mixers, loaders, applicators and all other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes and socks
- Chemical resistant gloves (except for pilots)

In addition, all handlers (except pilots, groundboom applicators, and airblast applicators) must wear:

- Wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N¹, R, or P filter; OR a NIOSH-approved powered air-purifying respirator with a HE filter.

See engineering controls for additional requirements.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls

Water soluble packets, when used correctly, qualify as a closed mixing/loading system under the Worker Protection Standard [40 CFR 170.607(d)]. Mixers and loaders handling this product while it is enclosed in intact water soluble packets may elect to wear reduced PPE of long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves. When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes and socks
- Chemical-resistant gloves

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Applications on golf courses or around municipal or industrial (office park) buildings and to ornamentals not grown for resale are not within the scope of the Worker Protection Standard.

- Keep unprotected persons out of treated area until sprays have dried.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage: Keep away from fire and sparks. Do not allow this product to become wet or overheated in storage: decomposition, impaired activity, or fire may result. Keep container closed when not in use.

Decomposition produces a foul odor; if observed, check for hot containers and immediately remove to open areas for disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 50 lb or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable nonrigid containers:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable rigid containers larger than 50 lb:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable rigid containers larger than 50 lb:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Product Information

Fore® 80WP Rainshield® specialty fungicide is a unique and different broad-spectrum protectant fungicide that, when used according to label directions, provides high levels of disease control.

Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program. The addition of a spreader-sticker will improve fungicide performance by providing a more uniform spray deposit, increased foliar redistribution, and improved fungicide retention during periods of wet weather.

Resistance Management

Repeated use of products for control of specific plant pathogens may lead to selection of resistant strains of fungi and result in a reduction of disease control. Mancozeb, the active ingredient in Fore 80WP Rainshield is one of EPA's Target Site of Action Group M3 fungicides and due to the multi-site mode of action of mancozeb, the risk of resistance development is highly unlikely and to date no reports of resistance development have been reported. For guidance on a particular crop and disease control situation, consult your state extension specialist for official state recommendations.

As part of a resistance management strategy:

- When tank mixing or alternating, use an effective partner – one that provides satisfactory disease control when used alone at the mixture rate.
- Apply early to keep fungal populations low.
- Incorporate integrated pest management (IPM) practices into your program which can help reduce disease development and spread.

Restrictions

- **Chemigation:** Do not apply this product through any type of irrigation system.
- Aerial application by fixed or rotary wing aircraft on ornamentals is prohibited.

Spray Drift Management

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed

Do not apply at wind speeds greater than 15 mph.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of mancozeb. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Use Rate Determination

Carefully read, understand, and follow label use rates and restrictions. Under low disease conditions, minimum label rates per application can be used while maximum label rates and the minimum interval may be used for severe or threatening disease conditions.

For proper application to turfgrass, determine the square footage to be treated, divide the footage by 1000, multiply by the required fungicide use rate per 1000 sq ft, and then determine the amount of water required to provide adequate coverage. When treating ornamentals, determine the required fungicide use rate and the spray gallonage required to provide a thorough coverage. Prepare only the amount of spray solution required to treat the desired area.

Handling of Water Soluble Packets

Water Soluble Packages (WSPs) are designed to dissolve in water. Agitation may be used, if necessary, to help dissolve the WSP. Failure to follow handling and mixing instructions can increase your exposure to the pesticide products in WSPs. WSPs, when used properly, qualify as a closed mixing/loading system under the Agricultural Worker Protection Standard [40 CFR 170.607(d)].

Handling Instructions

1. Mix in spray tank only.
2. Handle WSP(s) in a manner that protects package from breakage and/or unintended release of contents. If package is broken, put on PPE required for clean-up and then continue with mixing instructions.
3. Keep the WSP(s) in out packaging until just before use.
4. Keep the WSP dry prior to adding to the spray tank.
5. Handle with dry gloves and according to the label instructions for PPE.
6. Keep WSP intact. Do not cut or puncture WSP.
7. Reseal the WSP outer packaging to protect any unused WSP(s).

Mixing Directions

Follow the steps below when mixing this product, including if tank mixed with other pesticide products. If being tank mixed, the mixing directions 1 through 9 below take precedence over the mixing directions of the other tank mix products. WSPs may, in some cases, be mixed with other pesticide products so long as the directions for use and all mixed products do not conflict. Do not tank mix this product with products that prohibit tank mixing or have conflicting mixing directions.

1. If a basket or strainer is present in the tank hatch, remove prior to adding the WSP to the tank.
2. Fill tank with water to approximately one-third to one-half of the desired final volume of spray.
3. Stop adding water and stop any agitation.
4. Place intact/unopened WSP(s) into the tank.
5. Do not spray water from a hose or fill pipe to break or dissolve the WSP(s).
6. Start mechanical and recirculation agitation from the bottom of tank without using any overhead recirculation, if possible. If overhead recirculation cannot be turned off, close the hatch before starting agitation.
7. Dissolving the WSP(s) may take up to 5 minutes or longer, depending on water temperature, water hardness and intensity of agitation.
8. Stop agitation before tank lid is opened.
9. Open the lid to the tank, exercising caution to avoid contact with dusts or spray mix, to verify that the WSPs have fully dissolved and the contents have been thoroughly mixed into the solution.
10. Do not add other allowed products or complete filling the tank until the bags have fully dissolved and pesticide is thoroughly mixed.
11. Once the WSP have fully dissolved and any other products have been added to the tank, resume filling the tank with water to the desired level, close the tank lid, and resume agitation.
12. Use the spray solution when mixing is complete.
13. Maintain agitation of the diluted pesticide mix during transport and application.
14. It is unlawful to use any registered pesticide, including WSPs, in a manner inconsistent with its label.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Compatibility

Fore 80WP Rainshield is compatible with boron and spray oils; however, the water soluble packets must be completely dissolved before adding spray oils or products containing boron to spray mixtures.

Fore 80WP Rainshield is compatible with most commonly used agricultural fungicides, insecticides and growth regulators. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use. Do not tank mix with any product which contains a prohibition on tank mixing. When two pesticides are tank mixed, the more restrictive label conditions apply.

The physical spray compatibility of specific tank mixes can be checked by preparing a sample suspension in a glass container and observing the suspension for any adverse effects, which might interfere with proper spray application.

Spray Adjuvants

The addition of a surfactant to sprays containing Fore 80WP Rainshield will improve initial spray deposits, fungicide redistribution and weatherability. Place Fore 80WP Rainshield into suspension prior to

adding an adjuvant to the spray mixture. Read and carefully observe the precautionary statements and all other information appearing on both product labels prior to spray preparation. When an adjuvant is to be used with this product, Corteva Agriscience recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Application Directions

Fore 80WP Rainshield is a protectant fungicide. Optimum disease control is achieved with thorough coverage sprays applied on a regular schedule.

Disease Scouting

Fore 80WP Rainshield is a broad-spectrum protectant fungicide. If not applied on a routine protectant spray schedule, frequently observe turfgrass and ornamental plants for disease signs or symptoms. Apply fungicide at the required label use rate and spray schedule, at the first sign of disease, report of disease in the area, or during environmental conditions favorable for disease development.

Turfgrass

For use on golf courses, and industrial or municipal turf areas. Not for use on sod farms, athletic fields and residential turf.

Restrictions:

Golf Courses:

- For cool season turfgrasses; greens, tees and aprons - limit to a maximum of five applications per year at a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application.
- For cool season turfgrasses; fairways - limit to a maximum of four applications per year at a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application.
- For warm season turfgrasses; fairways - limit to a maximum of three applications per year at a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application.
- For warm season turfgrasses; greens, tees and aprons - limit to a maximum of four applications per year at a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application.
- Require a minimum of a 10-day interval between applications.

All Other Turfgrass:

- Limit to a maximum of four applications per year and a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application
- Require a minimum of a 10-day interval between applications

Each 24 oz water soluble packet treats 6000 sq ft at the 4 oz per 1000 sq ft rate, 4000 sq ft at the 6 oz per 1000 sq ft rate, and 3000 sq ft at the 8 oz per 1000 sq ft rate. Note that the 8 oz per 1000 sq ft rate is the same as the maximum rate of 21.75 lb per acre.

Start application when grass greens up in spring or when disease first appears and repeat at 10- to 14-day intervals or until disease threat is passed. When conditions are especially favorable for disease development, apply maximum fungicide use rate on a 10-day spray schedule. Apply in sufficient water to provide adequate coverage.

Turfgrass Tolerance

Maintain treated turfgrass in a vigorous growing condition. Turfgrass under stress will not respond as well to fungicide treatments as well-maintained turfgrass. Turfgrass tolerance to this product has been found to be acceptable; however, this product and tank mixtures with other products have not been tested on all varieties of every turfgrass species or under all possible growing conditions. If user is unfamiliar with the performance of Fore 80WP Rainshield or tank mixtures, under user growing conditions, treat a limited area of turfgrass prior to initiating large-scale applications.

Crop	Diseases Controlled	Fore 80WP Rainshield (oz/1000 sq ft)	Use Directions	Restrictions
assorted grasses	helminthosporium melting-out rust (leaf, stem, stripe)	4		<ul style="list-style-type: none"> • Do not graze treated areas. • Do not use on turfgrasses intended for grazing, such as range or pasture grasses. • Do not feed grass clippings to livestock. • Do not use on grasses grown for seed.
	copper spot fusarium blight red thread slime mold	4 - 8		
	algae	6		
	dollar spot	6 - 8		
	rhizoctonia brown patch	4	Apply on a 10-day spray schedule.	
	pythium blight	8	Apply at 10-day intervals	

Crop (Cont.)	Diseases Controlled	Fore 80WP Rainshield (oz/1000 sq ft)	Use Directions	Restrictions
assorted grasses	rapid blight (a putative chytridiomycete fungus)	8	Apply at a 14-day interval if conditions are favorable for disease development.	<ul style="list-style-type: none"> Do not graze treated areas. Do not use on turfgrasses intended for grazing, such as range or pasture grasses. Do not feed grass clippings to livestock. Do not use on grasses grown for seed.
	fusarium snow mold	6 - 8	Apply at 2- to 6-week intervals during winter.	
	gray leaf spot	8	Apply at 14-day intervals if conditions are favorable for disease development.	

Ornamentals

Restrictions:

- Cut flowers and greenhouse grown ornamentals: Limit to 20 applications per year.
- Do not use any part of treated plants, including fruit, nuts, bark, or sap, for food or feed purposes.
- Do not use on pachysandra.
- Do not apply more than a total of 8.4 lb of Fore 80WP Rainshield per acre (6.72 lb ai per acre) per year.
- Not for use in residential greenhouses.

Neither the manufacturer nor the seller has determined the effects of using Fore 80WP Rainshield on ornamentals not specified on this label. Prior

to any large-scale applications on such plants, the user should determine the effects of Fore 80WP Rainshield by testing a small section of the type of plants treated. User assumes all risk arising out of application to unlabeled plants. The Conditions of Sale and Warranty apply to all uses.

For outdoor or greenhouse use, apply the equivalent of 1 1/2 lb of Fore 80WP Rainshield per 100 gallons of dilute spray (1 1/2 lb of Fore 80WP Rainshield per acre). The addition of a surfactant to spray solutions will improve performance.

Begin spraying when plants are well leafed out or at first sign of disease in a full coverage spray at 7- to 10-day intervals throughout season or follow State Extension Service recommended retreatment intervals for disease control on the following ornamental plants:

Crop	Diseases Controlled	Crop Specific Use Directions
African violet	botrytis blight	
ageratum	botrytis blight rust	
almond (ornamental)	leaf spot	
alyssum		
anthurium	anthracnose, spadix rot	
apple (ornamental)	fabraea leaf spot rust scab	
arborvitae	cercospora blight	
areca palm	leaf spot	
ash, mountain	entomosporium leaf spot guignardia leaf blotch	
ash, white	anthracnose cylindrosporium leaf spot	
aster	leaf spot	
aster, perennial	puccinia rusts	
aucuba, japonica	alternaria leaf spot anthracnose	
azalea	cylindrocladium rot petal blight phytophthora twig and bud blight	Apply in a full coverage spray, 2 to 3 times a week, while flowers are opening. Direct spray into flowers and thoroughly spray ground under bushes.
begonia	botrytis blight	
bougainvillea	leaf spot	
boxwood		
buffaloberry	cylindrosporium leaf spot	
camellias	petal blight	Apply in a full coverage spray, 2 to 3 times a week, while flowers are opening. Direct spray into flowers and thoroughly spray ground under bushes.
carnation	rust septoria leaf spot	
cedar, red (juniper)	cercospora blight phomopsis blight	
chrysanthemum	ascochyta blight botrytis petal spot rust	Apply twice weekly during blooming period.
cockscomb (<i>Celosia</i>)	alternaria leaf spot	
conifers (Christmas trees)	lophodermium needle cast pine gall rust scirrhia brown spot	Begin application in spring or early summer before infection occurs. Repeat after heavy rains and at two- week intervals as long as needed.
cordyline	cercospora leaf spot	

Crop (Cont.)	Diseases Controlled	Crop Specific Use Directions
crabapple (ornamental)	cedar-apple rust scab sphaeropsis leaf spot	
cypress, Arizona (<i>Cupressus</i> sp.)	cercospora blight monochaetia canker	
dahlia	botrytis blight	
delphinium		
dieffenbachia	leptosphaeria brown spot	
dogwood, flowering	anthracnose elsinoe leaf spot septoria leaf spot	Apply when buds begin to open, when bracts have fallen, 4 weeks later and again in late summer after flower buds for next season have formed.
dracaena	fusarium leaf spot	
elm	black leaf spot	
euonymus	anthracnose	
fatsia		
fern	rhizoctonia blight	
figus	cercospora leaf spot	
fig	cylindrocladium leaf spot	
firethorn (<i>pyracantha</i>)	fusicladium scab	
fir, Douglas	Swiss needle cast	
fir, fraser		
fuchsia	botrytis blight rust	
geranium	rust	
gladiolus	botrytis blossom blight curvularia leaf spot	Make regular weekly applications starting before diseases appear and increase to 2 or 3 applications per week during periods of heavy disease and during rainy weather. On flower spikes, reduce spray concentration to 3/4 lb per 100 gallons.
gloxinia	botrytis blight	
gypsophila		
hawthorn	cedar-apple rust fabraea leaf spot frog-eye leaf spot hawthorn rust scab	
hickory	gnomonium leaf spot	
holly	purple spot	
hollyhock	anthracnose cercospora leaf spot puccinia rust	
honeysuckle	herpobasidium blight	
horsechestnut, buckeye	alternaria leaf spot guignardia leaf blotch	
hydrangea	botrytis blight cercospora leaf spot	
impatiens	botrytis blight	
iris	didymellina leaf spot mycosphaerella leaf spot mystrosporium ink spot	(formerly didymellina)
juniper	phomopsis blight	
larkspur	rust	
laurel, mountain	cercospora leaf spot petal blight	Apply in a full coverage spray, 2 to 3 times a week, while flowers are opening. Direct spray into flowers and thoroughly spray ground under bushes.
ligustrum	cercospora leaf spot	
lily	botrytis blight	
magnolia	gloeosporium leaf spot	
maple	alternaria leaf spot phyllosticta leaf spot	
marigold	botrytis blossom blight	Restriction: Do not use on French dwarf double or Signet type marigold seedlings.
narcissus	botrytis blight (fire) smoulder	
oak	actinopelte leaf spot taphrina leaf blister	

Crop (Cont.)	Diseases Controlled	Crop Specific Use Directions
orchid (<i>Dendrobium</i>)	botrytis blossom blight	
oxalis	rust	
pansy	anthracnose	
pears (ornamental)	fabraea leaf spot rust scab	
peony	botrytis blossom blight phytophthora blight	Apply in early spring and early fall, drenching soil around plants as well as the foliage. Promptly destroy all infected plant parts.
peperomia	cercospora leaf spot	
petunia	botrytis blight	
philodendron	dactylaria leaf spot phytophthora leaf spot	
phlox	leaf spot	
photinia	entomosporium leaf spot	
pine, Australia	cyclaneusma needle cast	
pine, Scotch	cyclaneusma needle cast gall rust	
pittosporum	alternaria leaf spot	
pleomele	fusarium leaf spot	
poinsettia	sphaceloma scab	
poplar	rust	
primrose	botrytis blight	
protea		
quince (ornamental)	fabraea leaf spot rust scab	
rhododendron	cercospora leaf spot discosia leaf spot petal blight	Apply in a full coverage spray, 2 to 3 times a week, while flowers are opening. Direct spray into flowers and thoroughly spray ground under bushes.
rose	black spot cercospora leaf spot rust	
rosemary	rhizoctonia aerial blight	
schefflera	alternaria blight	
Scotts pine	needle cast	
skunkbush, sumac	cylindrosporium leaf spot	
snapdragon	rust	
spathiphyllum	myrothecium leaf spot	
statice	cercospora frog-eye	
strawflower	rust	
syngonium	cephalosporium leaf spot	
thorn apple	rust	
tulip	botrytis blight (fire)	
venus, flytrap	anthracnose	
viburnum	downy mildew ramularia leaf spot	
walnut	anthracnose	
zinnia	alternaria leaf blight	

ATTENTION: This product contains mancozeb and ETU, chemicals known to the State of California to cause cancer. ETU is also known to the State of California to cause birth defects or other reproductive harm.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Corteva Agriscience be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

TMTrademarks of Corteva Agriscience and its affiliated companies

Produced for
Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268

Label Code: CD02-839-020
Replaced Label: D02-839-003

EPA accepted 06/22/18

Revisions:

- 1) Updated company name and trademark statement to Corteva Agriscience (from Dow AgroSciences).