

# Freehand® CA

## 1.75G Herbicide

**For use as a preemergence weed control herbicide in ornamental production, landscape and grounds maintenance, and other specified noncrop areas**

**Active Ingredients:**

dimethenamid-P: (S)-2-chloro-N-[(1-methyl-2-methoxy)ethyl]-N-(2,4-dimethyl-thien-3-yl)-acetamide. . . . .	0.75%
pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine. . . . .	1.00%

<b>Other Ingredients:</b> . . . . .	98.25%
-------------------------------------	--------

<b>Total:</b> . . . . .	100.00%
-------------------------	---------

50 lbs contains 0.375 lb of dimethenamid-P and 0.5 lb of pendimethalin.

**EPA Reg. No. 7969-358**

**EPA Est. No.**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific and/or use site restrictions.

**In case of emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).**

**Net Contents:**

FIRST AID	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give any liquid to the person.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eyes.</li> <li>• Call a poison control center for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Agricultural Solutions US LLC (hereafter "BASF") for emergency medical treatment information: 1-800-832-HELP (4357).</p>	

## Precautionary Statements

### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust.

### Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to

**Category A** on an EPA chemical-resistance category-selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof materials, such as barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, polyvinyl chloride  $\geq 14$  mils, or viton  $\geq 14$  mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This product is toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

Dimethenamid-P has properties that may result in ground-water contamination. Application in areas where soils are permeable or coarse and groundwater is near the surface could result in groundwater contamination.

Dimethenamid-P has properties that may result in surface water contamination via dissolved runoff and runoff erosion. Practices should be followed to minimize the potential for dissolved runoff and/or runoff erosion.

### Endangered Species Protection

This product may have effects on federally listed threatened or endangered plant species or their critical habitat. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the

pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

To avoid adverse effects on endangered plant species, applicators in ornamentals production must comply with the following mitigation measures where and when endangered plant species are known to occur in proximity of the application site:

If applied by ground, leave untreated buffer zone of 200 feet between treatment area and known endangered plant populations.

### Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application. The use of **FreeHand® CA 1.75G herbicide** not consistent with this label can result in injury to crops, animals, or persons.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

BASF does not authorize the use of this product in manufacturing, processing or preparing custom blends with other products for application to ornamentals.

**DO NOT** contaminate irrigation ditches or water used for domestic purposes.

**FreeHand CA 1.75G** is intended for use only by certified applicators or persons under their direct supervision.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL OR CROP INJURY.**

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof materials, such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

### NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**DO NOT** enter or allow others to enter the treated area until dusts have settled. Only protected applicator shall be in the treatment area during application.

### STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

#### Pesticide Storage

**DO NOT** store above 120° F. Store in original containers and keep closed. Store in a cool, dry place.

#### Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

(continued)

## STORAGE AND DISPOSAL *(continued)*

### Container Handling

#### (for plastic containers)

**Nonrefillable Container. DO NOT reuse or refill this container.** After completely emptying container into application equipment, offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

#### (for paper or plastic bags)

**Nonrefillable Container. DO NOT reuse or refill this container.** After completely emptying container into application equipment, dispose of empty bag in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

## In Case Of Emergency

In case of large-scale spillage regarding this product, call:

- CHEMTREC 1-800-424-9300
- BASF 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF 1-800-832-HELP (4357)

## Use Information

### Mode of Action

**FreeHand® CA 1.75G herbicide** combines two active ingredients: dimethenamid-P, a chloroacetamide, and pendimethalin, a dinitroaniline. Combined, they inhibit both plant cell microtubule assembly (cells do not divide and multiply) and cell growth, inhibiting the germinating shoot portion of the susceptible weed seedlings. A small amount of uptake may also occur through the emerging root system of the weed seedling. **FreeHand CA 1.75G** is **Chemical Group 3** and **Chemical Group 15** mode-of-action classifications for broader weed control spectrum.

### Use Sites

**FreeHand CA 1.75G** is a selective preemergence herbicide for the control of certain annual grasses, annual broadleaf weeds and sedges as they germinate in:

- **Commercial ornamental production**
- **Landscape and grounds maintenance**
- **Tree plantations including Christmas trees**

**FreeHand CA 1.75G** may be applied as a soft-residual bareground treatment in the use sites listed above.

**FreeHand CA 1.75G** controls weeds as they germinate but will not control emerged and established weeds. Therefore, areas to be treated should be free of established weeds. If weeds develop prior to activation of the herbicide, shallow cultivate to destroy existing weeds or, where practical, remove by hand.

For the control of established weeds,

**FreeHand CA 1.75G** may be used in conjunction with herbicides registered for postemergence use.

**FreeHand CA 1.75G** may be used before or after applications of herbicides registered for postemergence use (i.e. **Basagran® T&O herbicide**, **Finale® herbicide**, **Image® 70 DG herbicide**, **Roundup® herbicide**, **Segment® herbicide**) for the control of established weeds. **DO NOT** apply sprays containing **Finale** or **Roundup** over the top of desirable plants. Consult the labels of those herbicides for use rates, timings, and precautions or restrictions.

A **FreeHand CA 1.75G** treatment may be followed by any registered herbicide to control weeds not listed on the **FreeHand CA 1.75G** label.

## Application Information

**FreeHand CA 1.75G** will provide the most effective weed control when applied by ground spreader equipment and subsequently incorporated into soil by 1/2 inch of rainfall in sprinkler irrigation.

### Ground Application

Apply **FreeHand CA 1.75G** with properly calibrated spreader equipment to provide uniform distribution. Check application routinely to determine proper calibration. Avoid overlaps that will increase use rates above those stated on this label. Avoid application when winds may cause drift. Avoid unintentional contact of granules with driveways, stone, wood or other porous surfaces.

**FreeHand CA 1.75G** may cause temporary discoloration of treated surfaces. Sweep surfaces clean immediately to avoid staining. If granules are crushed into surface, rinse thoroughly.

### Application Use Rates

For preemergence control of the listed weed species (see **Table 3** in **Weed Species Controlled** section) using broadcast spreader equipment, apply

**FreeHand CA 1.75G** at the following rates. Uneven application can decrease weed control or cause plant injury. Overapplication can result in stand loss, plant injury, or soil residues.

**Table 1. Application Rates for FreeHand CA 1.75G**

Application Rate (lbs product per acre)	Amount (lbs product per 1000 sq ft)	Amount (lb product per 100 sq ft)
100	2.3	0.23
150	3.4	0.34
200	4.6	0.46

All granular herbicide products, including **FreeHand® CA 1.75G herbicide**, have different prill sizes. Therefore, to deliver the proper rate, the applicator must calibrate prior to application of product. The **FreeHand CA 1.75G** calibration tray may be used to calibrate the correct application rate. Rate settings for a specific type of spreader device must be determined by the applicator. A guide to the use of various spreaders is provided in the following table.

## Spreader Use Guide

Spreader Type	Spreader Operation	Setting	Swath of Throw (feet)	Rate Range (lbs product/A)
EarthWay <sup>1</sup>	1-sided	7 to 10	3	200
EarthWay <sup>2</sup>	2-sided	11 to 12	3 to 5	150 to 200
John Deere-Lesco Rotary	Not applicable	D to F	Not given	100 to 150
Scott Rotaries R-7, R-7x, R-8, R-8A, SR-1 (cone 9)	Not applicable	D to E1/2	8	100 to 150
Solo Spreader 421S <sup>3</sup>	1-sided	1 to 3	3 to 4	100
Solo Spreader 421S <sup>4</sup>	2-sided	2 to 6	3 to 4	200
Spyker #75 <sup>a</sup>	1-sided	3 to 5	3 to 5	100
Warren T7II <sup>b</sup>	1-sided	3 to 5	3 to 5	100
Warren T7II <sup>c</sup>	2-sided	3 to 6	3 to 6	100

### Spreader Efficiency Tips

- <sup>1</sup> **For EarthWay 1-sided:** Close right slide on spreader to throw to the right.
- <sup>2</sup> **For EarthWay 2-sided:** Close left slide about 1/4 to achieve better left center to right distribution for 2-sided throw.
- <sup>3</sup> **For Solo 1-sided:** Position lever to the right (this gives a pattern all to the left and avoids depositing product into operator's hands or body).
- <sup>4</sup> **For Solo 2-sided:** Position lever halfway between center and right position.
- <sup>a</sup> **For Spyker:** Rotate spreader to right so first mold pen mark is on top at 24° to give best distribution. NOT recommended for 2-sided use.
- <sup>b</sup> **For Warren 1-sided:** Distribution place control on front all the way to the left so pattern spreads to left and not into the applicator's arms.
- <sup>c</sup> **For Warren 2-sided:** Position lever in center for best distribution.

**FreeHand CA 1.75G** may be applied in a single application or in sequential applications.

For extended weed control or where heavy weed infestations are expected, sequential (repeat) applications of **FreeHand CA 1.75G** can be made 5 to 8 weeks between applications.

## Application Restrictions and Limitations

- **Maximum annual use rate - DO NOT** apply more than a total of 400 lbs/A **FreeHand CA 1.75G** per crop in a single growing season.
- **DO NOT** apply more than 200 lbs/A **FreeHand CA 1.75G** in a single application.
- **DO NOT** apply **FreeHand CA 1.75G** to ornamental plants when springtime soil temperature is less than 55° F and under wet soil conditions, or plant injury may occur.
- **DO NOT** apply **FreeHand CA 1.75G** as an aerial application.
- **DO NOT** apply **FreeHand CA 1.75G** in greenhouses, polyhouses or other fully enclosed greenhouse-type structures.
- **DO NOT** treat plants grown for food or feed.
- **DO NOT** use treated plants for food or feed.
- To avoid the possibility of plant damage, **DO NOT** apply **FreeHand CA 1.75G** to ornamental plants growing under stress because of lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures. Diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken plants and increase the possibility of damage from **FreeHand CA 1.75G**.
- If **FreeHand CA 1.75G** is not activated by rainfall or irrigation within 30 days, erratic weed control may result.
- Unusually cold, excessively wet, or hot and dry conditions that delay or extend weed seed germination can reduce weed control.

## Specific Information for Use Sites

### Commercial Ornamental Production

**FreeHand CA 1.75G** can be used in container and field nurseries.

Applications can be made to ornamental plant species listed on this label, such as trees, shrubs, ground covers, herbaceous perennials, and bedding plants. Applications can also be made to conifer and hardwood seedling nurseries, and tree plantations (including Christmas trees).

**NOTE: FreeHand® CA 1.75G herbicide can only be used on established liner beds with well-rooted plants and/or rootstocks.**

## **Plant Tolerance**

**FreeHand CA 1.75G** can be used around and over the top of the established ornamental plants listed in **Table 2** of this label. However, not all varieties or strains of the ornamental plants listed have been tested.

**FreeHand CA 1.75G** may be used on ornamental plant species not listed on this label when grower assumes responsibility. Determine the suitability for such uses by treating a small number of such plants at the specified rate. Evaluate treated plants for at least 2 months following treatment for possible injury. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE USER ASSUMES RESPONSIBILITY FOR ANY CROP DAMAGE OR OTHER LIABILITY ASSOCIATED WITH APPLICATIONS TO ORNAMENTAL PLANTS NOT LISTED ON THIS LABEL.**

Refer to the **Application Information** section in this label before any application of **FreeHand CA 1.75G**. Unintentional consequences, such as ornamental injury, may result because of certain environmental or growing conditions, manner of use or application.

Refer to **Table 1** for application rates, **Table 2** for list of ornamental species, and **Table 3** for weeds controlled.

## Specific Ornamentals<sup>1</sup> Use Site Instructions

Site	Application Instructions and Restrictions
Newly transplanted field-grown nursery plants	<ul style="list-style-type: none"> <li>• <b>Restriction: DO NOT</b> apply until transplants have been watered and soil has been thoroughly packed and settled around transplants. Care must be taken to ensure there are no cracks in the soil where <b>FreeHand® CA 1.75G herbicide</b> could come into contact with the roots.</li> <li>• <b>Restriction: DO NOT</b> apply during bud swell, bud break or at time of first flush of new growth.</li> </ul>
Newly transplanted container-grown nursery plants	<ul style="list-style-type: none"> <li>• <b>Restriction: DO NOT</b> apply until transplants have been watered and soil has been thoroughly packed and settled around transplants. Care must be taken to ensure there are no cracks in the soil where <b>FreeHand CA 1.75G</b> could come into contact with the roots.</li> <li>• For container-grown ornamentals, delay first application of the product to bareroot liners for 2 weeks after transplanting to allow for root establishment.</li> <li>• <b>Restriction: DO NOT</b> apply during bud swell, bud break or at time of first flush of new growth.</li> <li>• <b>Restriction: DO NOT</b> apply to any size plug or liner tray.</li> <li>• <b>Restriction: DO NOT</b> apply to plugs or unrooted liners or cuttings being transplanted into pots.</li> </ul>
Established container or field-grown nursery plants	<ul style="list-style-type: none"> <li>• <b>Restriction: DO NOT</b> apply during bud swell, bud break or at time of first flush of new growth.</li> <li>• Care must be taken to ensure there are no cracks in the soil where <b>FreeHand CA 1.75G</b> could come into contact with the roots.</li> </ul>
Field-grown or container-grown production bulbs	<ul style="list-style-type: none"> <li>• For use in ornamental bulb-like crops such as caladium, calla lily, and lilies.</li> <li>• In field production, apply <b>FreeHand CA 1.75G</b> to the soil surface only after the crop has been planted and the soil has been settled by several irrigations but prior to weed seed germination.</li> <li>• In fall-planted daffodil, iris or lilies, make an initial application of <b>FreeHand CA 1.75G</b> following plant establishment; then make a sequential application of <b>FreeHand CA 1.75G</b> in late winter or early spring prior to weed seed germination.</li> <li>• In container production, apply <b>FreeHand CA 1.75G</b> to a weed-free surface either before bulb emergence or after leaf emergence from an established plant crown.</li> <li>• Apply <b>FreeHand CA 1.75G</b> before, during or after bulb emergence.</li> </ul>
<p>Bareground pad for container placement</p> <p>Gravel or ground floor of open-sided lathhouses (shadehouses) or other polyhouse structures that allow polycovers to be removed on a seasonal basis</p>	Apply to bareground pad (such as soil, mulch, gravel, wood chips, or other permeable base); then water in; then replace containerized ornamentals onto the pad.
Greenhouses, polyhouses or other enclosed structures	<b>Restriction: DO NOT</b> apply in greenhouses, polyhouses or other fully enclosed greenhouse-type structures.
Landscape uses	Care must be taken that soil or planting mixes have settled firmly following transplanting, and there are no cracks that would allow direct contact of <b>FreeHand CA 1.75G</b> with roots. Applications where soil and media surfaces are uniformly covered will result in best weed control and ornamental tolerance.

<sup>1</sup> Plant only those desirable plant species listed on this label into soil treated the previous season with **FreeHand CA 1.75G** or injury may occur.

## Specific Ornamentals<sup>1</sup> Use Site Instructions *(continued)*

Site	Application Instructions and Restrictions		
SENSITIVE Ornamental Species - Special Precautions			
Ornamental grasses	Restriction: DO NOT apply FreeHand® CA 1.75G herbicide to ornamental grasses because unacceptable phytotoxicity may occur.		
Herbaceous perennials	Restriction: DO NOT apply FreeHand CA 1.75G to herbaceous perennials because unacceptable phytotoxicity may occur.		
	Black-eyed Susan	Rudbeckia fulgida	
	Blue star	Amsonia spp.	
	Butterfly flower	Asclepias tuberosa	
	Butterfly weed	Asclepias tuberosa	
	Cape leadwort	Plumbago spp.	
	Columbine	Aquilegia spp.	
	Dead nettle	Lamium spp.	
	Periwinkle	Vinca minor	
	Phlox	Phlox paniculata	
	Pincushion flower	Scabiosa spp.	
	Speedwell, spiked	Veronica spicata	
	other Veronica spp.		
Tickseed	Coreopsis auriculata		
Annual bedding plants	Restriction: DO NOT apply FreeHand CA 1.75G to begonia or severe injury will occur. FreeHand CA 1.75G application to gomphrena and New Guinea impatiens may cause injury in some cultivars.		
Ferns	Applications of FreeHand CA 1.75G to immature ferns during periods of new growth of fronds may result in some injury.		
Hydrangea	Not all hydrangea cultivars may respond to the herbicide application with the same tolerance. Before treating an entire block of plants, apply only to a small number of plants and evaluate for 2 months for tolerance.		
Restriction: DO NOT apply FreeHand CA 1.75G sequentially to the following listed species. During the growing season, a second application of FreeHand CA 1.75G can be made if a herbicide of a different mode of action is applied between FreeHand CA 1.75G applications. FreeHand CA 1.75G applications must be separated by at least 16 weeks.			
Coneflower	Echinacea spp.	Spirea	Spiraea spp.
Goldenrod	Solidago spp.	Thrift	Armeria spp.
Hydrangea	Hydrangea spp.	Turtlehead/Snakehead	Chelone spp.
Loropetalum	Loropetalum chinense	Viburnum	Viburnum spp.
Magnolia	Magnolia spp.	Western hemlock	Tsuga heterophylla
Sage	Salvia spp.		

<sup>1</sup> Plant only those desirable plant species listed on this label into soil treated the previous season with **FreeHand CA 1.75G** or injury may occur.

**Table 2. Ornamental Species**

Common Name	Scientific Name
<b>Trees</b>	
Arborvitae	<i>Thuja</i> spp.
Crape myrtle	<i>Lagerstroemia indica</i>
Cypress, Leyland	<i>Cupressocyparis leylandii</i>
Fir, Douglas	<i>Pseudotsuga menziesii</i>
Fir, Fraser	<i>Abies fraseri</i>
Hemlock	<i>Tsuga</i> spp.
Hemlock, Western	<i>Tsuga heterophylla</i>
Lilac	<i>Syringa</i> spp.
Magnolia	<i>Magnolia</i> spp.
Oak	<i>Quercus</i> spp.
Pine	<i>Pinus</i> spp.
Serviceberry	<i>Amelanchier</i> spp.
<b>Shrubs</b>	
Abelia, glossy	<i>Abelia grandiflora</i>
Azalea	<i>Rhododendron</i> spp.
Bamboo, heavenly	<i>Nandina domestica</i>
Barberry	<i>Berberis</i> spp. <i>Berberis thunbergii</i>
Boxwood, Japanese	<i>Buxus microphylla</i>
Butterfly bush	<i>Buddleia davidii</i>
Cotoneaster	<i>Cotoneaster</i> spp.
Euonymus	<i>Euonymus</i> spp.
Forsythia, border	<i>Forsythia intermedia</i>
Gardenia	<i>Gardenia jasminoides</i>
Hawthorne, Indian	<i>Raphiolepis indica</i>
Hibiscus	<i>Hibiscus</i> spp.
Holly	<i>Ilex</i> spp.
Holly, Chinese	<i>Ilex cornuta</i>
Holly, Japanese	<i>Ilex crenata</i>
Hydrangea	<i>Hydrangea macrophylla</i>
Juniper	<i>Juniperus</i> spp.
Lavender, English	<i>Lavandula angustifolia</i>
Lilac	<i>Syringa</i> spp.
Lily of the Nile	<i>Agapanthus africanus</i>
Loropetalum	<i>Loropetalum chinense</i>
Mahonia	<i>Mahonia</i> spp.
Mock orange	<i>Pittosporum tobira</i>
Nandina	<i>Nandina domestica</i>
Oregon grape	<i>Mahonia</i> spp.
Photinia, Fraser	<i>Photinia x fraseri</i>
Pine, Mugo	<i>Pinus mugo</i>
Privet	<i>Ligustrum</i> spp.
Quince	<i>Chaenomeles</i> spp.
Redroot	<i>Ceanothus</i> spp.
Red-tip	<i>Photinia x fraseri</i>

(continued)

**Table 2. Ornamental Species** (continued)

Common Name	Scientific Name
<b>Shrubs</b> (continued)	
Rhododendron	<i>Rhododendron</i> spp.
Rose	<i>Rosa</i> spp.
Rose-of-Sharon	<i>Hibiscus</i> spp.
Spiraea	<i>Spiraea</i> spp.
Viburnum	<i>Viburnum</i> spp.
Weigela	<i>Weigela florida</i>
Wild lilac	<i>Ceanothus</i> spp.
<b>Ground Covers</b>	
Cinquefoil, spring	<i>Potentilla</i> spp. <i>Potentilla fruticosa</i>
Gazania	<i>Gazania splendens</i>
Lilyturf, big blue or giant	<i>Liriope muscari</i>
Pachysandra	<i>Pachysandra terminalis</i>
Sedum/Stonecrop	<i>Sedum</i> spp.
Verbena	<i>Glandularia</i> spp.
<b>Herbaceous Perennials</b>	
Alumroot	<i>Heuchera sanguinea</i>
Anemone	<i>Anemone hupehensis</i>
Astilbe (False spirea)	<i>Astilbe</i> spp.
Clematis	<i>Clematis</i> spp.
Coneflower	<i>Echinacea</i> spp.
Coral bells	<i>Heuchera sanguinea</i>
Daylily	<i>Hemerocallis</i> spp.
Flag	<i>Iris</i> spp.
Gaura	<i>Gaura lindheimeri</i>
Goldenrod	<i>Solidago</i> spp.
Hosta	<i>Hosta</i> spp.
Hyssop	<i>Agastache</i> spp.
Iris	<i>Iris</i> spp.
Jasmine, confederate	<i>Trachelospermum jasminoides</i>
Lantana	<i>Lantana</i> spp.
Lavender, English	<i>Lavandula angustifolia</i>
Lily of the Nile	<i>Agapanthus africanus</i>
Liriope	<i>Liriope</i> spp.
Lobelia	<i>Lobelia</i> hybrids
Purple coneflower	<i>Echinacea</i> spp.
Sage	<i>Salvia</i> spp. <i>Salvia x superba</i>
Sage, Ramona	<i>Salvia sylvestris</i>
Shasta daisy	<i>Chrysanthemum x superbum</i>
Stonecrop	<i>Sedum</i> spp.
Thrift sea pink	<i>Armeria maritima</i>
Turtlehead/Snakehead	<i>Chelone</i> spp.
Yarrow	<i>Achillea millefolium</i>

(continued)

**Table 2. Ornamental Species** *(continued)*

Common Name	Scientific Name
<b>Bedding Plants</b>	
Alyssum, sweet	<i>Lobularia maritima</i>
Gazania	<i>Gazania</i> spp.
Periwinkle, rose	<i>Catharanthus roseus</i>
Petunia	<i>Petunia</i> spp.
Sweet potato, ornamental	<i>Ipomoea batatas</i>
Verbena	<i>Glandularia x hybrida</i>

### Landscape and Grounds Maintenance

**FreeHand® CA 1.75G herbicide** can be used in landscape and grounds maintenance programs to provide extended preemergence weed control. Areas to be treated include those areas where gravel, pebble, stone, rock and other porous surfaces are used in landscaping instead of organic mulches. **FreeHand CA 1.75G** can be used in established ornamental plantings in nonagricultural areas defined as follows:

- **Landscaped ornamental maintenance areas** in residential and commercial establishments, multifamily dwellings, military and other institutions, university or college campuses, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses, and prairie grass areas, common areas in residential developments, and in nonproduction areas in commercial nurseries, such as storage areas, vegetation filter strips, windbreaks, shelterbelts or cart paths.
- **General grounds maintenance in the following specified noncrop areas** – Parking lots, driveways and roadsides, highway rights-of-way, alleyways, bike and jogging paths, vacant lots, buildings, stone gardens and gravel yards, around statuary or monuments, utility substations, markers/borders and fence lines, and mulch beds. **FreeHand CA 1.75G** may be used under asphalt or concrete treatments as part of a site preparation program.

Refer to **Table 1** for application rates, **Table 2** for list of ornamental species, and **Table 3** for weeds controlled.

### Tree Plantations

Use **FreeHand CA 1.75G** for preemergence weed control during site preparation, establishment and/or maintenance of tree plantations (including Christmas trees). Also use **FreeHand CA 1.75G** for hardwood and conifer regeneration on Conservation Reserve Program (CRP) land or similar areas.

**FreeHand CA 1.75G** may be applied at planting or to established trees. When making an application at planting, it is important that slit closure be achieved to prevent **FreeHand CA 1.75G** from directly contacting the tree roots or being washed into the root zone via the open slit, or root stunting may occur. **DO NOT** apply to newly

transplanted seedlings until plants have been watered and soil has been thoroughly packed and settled around roots.

Refer to **Table 1** for application rates, **Table 2** for list of ornamental (tree) species, and **Table 3** for weeds controlled.

### Weed Species Controlled

**Table 3. Weeds Controlled**

Use **FreeHand CA 1.75G** for preemergence control of the following weed species listed according to use rate.

100 lbs/A	Weeds controlled when <b>FreeHand CA 1.75G</b> is applied at 100 lbs/acre (2.3 lbs <b>FreeHand CA 1.75G</b> /1000 sq ft)	
	Common Name	Scientific Name
	Amaranth, Palmer	<i>Amaranthus palmeri</i>
	Barnyardgrass	<i>Echinochloa crus-galli</i>
	Carpetweed	<i>Mollugo verticillata</i>
	Foxtail, giant	<i>Setaria faberi</i>
	Foxtail, green	<i>Setaria viridis</i>
	Foxtail, yellow	<i>Setaria glauca</i>
	Henbit	<i>Lamium amplexicaule</i>
	Lambsquarters, common	<i>Chenopodium album</i>
	Pigweed, prostrate	<i>Amaranthus blitoides</i>
	Pigweed, redroot	<i>Amaranthus retroflexus</i>
	Pigweed, smooth	<i>Amaranthus hybridus</i>
	Pigweed, tumble	<i>Amaranthus albus</i>
	Pusley, Florida	<i>Richardia scabra</i>

**Table 3. Weeds Controlled** (continued)

<p><b>150 lbs/A</b></p> <p>In addition to the weeds controlled at 100 lbs <b>FreeHand® CA 1.75G herbicide/</b> acre application rate, the following weeds will be controlled when <b>FreeHand CA 1.75G</b> is applied at 150 lbs/acre (3.4 lbs <b>FreeHand CA 1.75G</b>/1000 sq ft)</p>	
Common Name	Scientific Name
Amaranth, Powell	<i>Amaranthus powellii</i>
Bittercress	<i>Cardamine</i> spp.
Bluegrass, annual	<i>Poa annua</i>
Bluegrass, roughstalk	<i>Poa trivialis</i>
Brome, California	<i>Bromus carinatus</i>
Brome, downy	<i>Bromus tectorum</i>
Chamomile, mayweed	<i>Anthemis cotula</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Cudweed	<i>Gnaphalium</i> spp.
Cupgrass, Southwestern	<i>Eriochloa gracilis</i>
Doveweed	<i>Murdannia nudiflora</i>
Evening primrose	<i>Oenothera biennis</i>
Flatsedge, rice	<i>Cyperus iria</i>
Goosegrass	<i>Eleusine indica</i>
Groundsel, common	<i>Senecio vulgaris</i>
Johnsongrass (from seed)	<i>Sorghum halepense</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, cutleaf	<i>Solanum triflorum</i>
Nightshade, Eastern black	<i>Solanum ptycanthum</i>
Nightshade, hairy	<i>Solanum sarrachoides</i>
Purslane, common	<i>Portulaca oleracea</i>
Red rice	<i>Oryza sativa</i>
Ryegrass, Italian	<i>Lolium multiflorum</i>
Sandbur, field	<i>Cenchrus incertus</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Speedwell, corn	<i>Veronica arvensis</i>
Spurge, annual	<i>Chamaesyce</i> spp. <i>Euphorbia</i> spp.
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatus</i>
Witchgrass	<i>Panicum capillare</i>

**Table 3. Weeds Controlled** (continued)

<p><b>200 lbs/A</b></p> <p>In addition to the weeds controlled at 100 lbs and 150 lbs <b>FreeHand CA 1.75G/</b> acre application rates, the following weeds will be controlled when <b>FreeHand CA 1.75G</b> is applied at 200 lbs/acre (4.6 lbs <b>FreeHand CA 1.75G</b>/1000 sq ft)</p>	
Common Name	Scientific Name
Burweed, lawn	<i>Soliva pterosperma</i>
Clover, hop	<i>Trifolium procumbens</i>
Crowfootgrass	<i>Dactyloctenium aegyptium</i>
Cupgrass, woolly	<i>Eriochloa villosa</i>
Eclipta	<i>Eclipta alba</i> <i>E. prostrata</i>
Fiddleneck	<i>Amsinckia intermedia</i>
Itchgrass	<i>Rottboellia exaltata</i>
Junglerice	<i>Echinochloa colona</i>
Kochia	<i>Kochia scoparia</i>
Lovegrass (from seed)	<i>Eragrostis</i> spp.
Nutsedge, yellow	<i>Cyperus esculentus</i>
Panicum, browntop	<i>Panicum fasciculatum</i>
Panicum, fall	<i>Panicum dichotomiflorum</i>
Panicum, Texas	<i>Panicum texanum</i>
Puncturevine	<i>Tribulus terrestris</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Rocket, London	<i>Sisymbrium irio</i>
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Sowthistle	<i>Sonchus arvensis</i>
Sprangletop, Mexican	<i>Leptochloa uninervia</i>
Sprangletop, red	<i>Leptochloa filiformis</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Willowherb	<i>Epilobium</i> spp.
Woodsorrel, yellow	<i>Oxalis</i> spp. <i>O. corniculata</i> <i>O. stricta</i>

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Agricultural Solutions US LLC ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108P

***Basagran**, **FreeHand**, and **Segment** are registered trademarks of BASF.*

***Finale** is a registered trademark of Bayer.*

***Image** is a registered trademark of Central Garden & Pet Company.*

***Roundup** is a registered trademark of Monsanto Technology, LLC.*

© 2024 BASF Agricultural Solutions US LLC  
All rights reserved.

007969-00358.20241014.**NVA 2024-04-0408-0290**

Based on: NVA 2013-04-408-0029

Supersedes: NVA 2013-04-408-0099

BASF Agricultural Solutions US LLC  
2 TW Alexander Drive  
Research Triangle Park, NC 27713

  
We create chemistry