

**Product Name:
Freestyle™**

Freestyle™ is a combination package of products:

Classic® Grande PCP # 29416

DuPont Imazethapyr 240 SL PCP # 31157

Labels are as attached.

Manufacturer/Distributor:

E.I. du Pont Canada Company
1919 Minnesota Court
Mississauga, Ontario, L5N 0C9

Telephone Numbers :

Product Information : 1-800-667-3925

Medical Emergency : 1-800-441-3637 (24 hours)

Freestyle™ and Classic® Grande are registered trademarks or trademarks of E. I. du Pont de Nemours and Company. E.I. du Pont Canada Company is a licensee. Member of CropLife Canada.

*All other products mentioned are trademarks of their respective companies.

PMRA approved revised label, JPR, 2015-06-15
2015-0901

[[BARRIER BAG LABEL]]

GROUP

2

HERBICIDE

CLASSIC[®] GRANDE HERBICIDE

Wettable granules

FOR SALE FOR USE ON SOYBEANS IN EASTERN CANADA ONLY

COMMERCIAL

REGISTRATION NO. 29416 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE: Chlorimuron-ethyl 25.0%

NET CONTENTS: 14-14,400 grams

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

Avoid breathing spray mist.

Avoid contact with skin, eyes and clothing.

Keep the container or bag away from water.

Do not contaminate any body of water.

Do not enter or allow others to enter treated areas during the restricted entry interval of 12 hours after application.

FIRST AID:

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE:

Store product in original container in a secure, dry area away from food or feed. Not for use or storage in or around the home. Keep container tightly closed. Keep product in original container or bag away from moisture.

DISPOSAL:

1. **Triple-or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.**
2. **Follow provincial instruction for any required additional cleaning of the container prior to its disposal.**
3. Make the empty container unsuitable for further use.
4. Dispose of the empty container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

[[DIRECTIONS FOR USE LABEL]]

GROUP

2

HERBICIDE

CLASSIC® GRANDE HERBICIDE

Wettable granules

FOR SALE FOR USE ON SOYBEANS IN EASTERN CANADA ONLY

COMMERCIAL

REGISTRATION NO. 29416 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND THIS BOOKLET BEFORE USING

GUARANTEE: Chlorimuron-ethyl 25.0%

NET CONTENTS: 14-14,400 grams

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

Avoid breathing spray mist.

Avoid contact with skin, eyes and clothing.

Keep the container or bag away from water.

Do not contaminate any body of water.

Do not enter or allow others to enter treated areas during the restricted entry interval of 12 hours after application.

IMPORTANT:

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on other crops. Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than soybeans.

DO NOT APPLY BY AIR.

APPLY CLASSIC® GRANDE HERBICIDE WITH A RECOMMENDED SURFACTANT.

Do not handle pesticides with bare hands. Chemical-resistant gloves significantly reduce hand exposure. Wear gloves for mixing/loading/cleanup operations and when making sprayer and nozzle repairs and adjustments. Do not use leather or cloth gloves. Wear coveralls, or long-sleeved shirt and long pants, for mixing, loading, and applying, and during clean-up and repair activities.

FIRST AID:

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically

STORAGE:

Store product in original container in a secure, dry area away from food or feed. Not for use or storage in or around the home. Keep container tightly closed. Keep product in original container or bag away from moisture.

DISPOSAL:

1. **Triple-or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.**
2. **Follow provincial instruction for any required additional cleaning of the container prior to its disposal.**
3. Make the empty container unsuitable for further use.
4. Dispose of the empty container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

GENERAL INFORMATION:

CLASSIC® GRANDE HERBICIDE is recommended for selective postemergence control of broadleaf weeds in soybeans. CLASSIC® GRANDE HERBICIDE can also be applied pre plant or post plant preemergence to soybeans, either conventional or glyphosate tolerant. CLASSIC® GRANDE HERBICIDE is formulated as a dry flowable granule to be mixed in water and applied as a uniform broadcast spray. It is noncorrosive, nonflammable, nonvolatile, and does not freeze. Keep product away from excess moisture or physical damage during storage and prior to mixing in the spray tank. Do not reuse container or bag.

CLASSIC® GRANDE HERBICIDE can be applied with a recommended surfactant either as a pre plant or post plant preemergence application to soybeans, either conventional or glyphosate tolerant or early postemergence to the main flush of actively growing weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of CLASSIC® GRANDE HERBICIDE by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress may not be adequately controlled and regrowth may occur. CLASSIC® GRANDE HERBICIDE may only be applied using ground equipment. For best results, ensure thorough spray coverage of target weeds.

CLASSIC® GRANDE HERBICIDE rapidly stops growth of susceptible weeds. However, typical symptoms (discoloration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Favourable growing conditions following treatment promote the activity of CLASSIC® GRANDE HERBICIDE while cold, dry conditions delay the activity.

Degree of control and duration of effect depend on weed sensitivity, weed size, crop competition, growing conditions and spray coverage.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY:

The following precautions are recommended for application of CLASSIC® GRANDE HERBICIDE on soybean.

CLASSIC® GRANDE HERBICIDE applications made during or immediately following a large and rapid fluctuation in temperature (20°C or greater difference between night time lows and daytime highs) which stresses the crop may increase the potential for crop injury.

Apply CLASSIC® GRANDE HERBICIDE when temperature is expected to be below 28 degrees C within 12 hours of application.

Apply when the minimum temperature in the 24 hours before and after application is above 5 degrees C. If soybeans have been injured by frost, wait at least 48 hours before applying the herbicide(s).

DIRECTIONS FOR USE:

RATES AND TIMING:

PRE EMERGENT:

For the season long control of dandelion (up to bolting) apply 36 g/ha of CLASSIC® GRANDE HERBICIDE, plus Agral 90 at 0.2%v/v, pre plant or post plant pre-emergence to soybeans, either in conventional or glyphosate tolerant.

POSTEMERGENT APPLICATION:

Optimum timing of CLASSIC® GRANDE HERBICIDE for annual broadleaf weed control is when soybeans are in the 1 to 3 trifoliolate stage of growth. Do not apply CLASSIC® GRANDE HERBICIDE before the first trifoliolate has fully expanded or after the initiation of flowering. CLASSIC® GRANDE HERBICIDE can be applied up to 60 days before harvest.

Use 36 grams of CLASSIC® GRANDE HERBICIDE per hectare. A non-ionic surfactant such as AGRAL* 90, CITOWETT* PLUS, or AG-SURF* must also be used at 0.2% v/v. OPTIONAL: The addition of liquid fertilizer containing 28% nitrogen to the spray mixture at the rate of 2 litres per hectare may improve control of velvetleaf and should be added after the surfactant.

For best results, apply to young actively growing weeds before the soybean canopy closes. Weeds that emerge after treatment may not be controlled.

NOTE: Do not apply to soybeans that have been stressed by severe weather conditions such as frost or hot, humid weather as crop injury may result.

If rain occurs after application, control may be reduced. Two - four hours of dry weather are needed to allow CLASSIC® GRANDE HERBICIDE to be absorbed by weed foliage.

WEEDS CONTROLLED BY CLASSIC® GRANDE HERBICIDE ALONE:

Weeds	Leaf Stage	Height (cm)
Common Ragweed	2-6	10 cm tall or across
Redroot Pigweed	2-8	"
Velvetleaf	2-4	"
Yellow Nutsedge	Up to 8	5-15 cm tall or across
Dandelion*	pre-bloom	15 cm tall or across

*Top Growth Control (4 to 6 weeks)

CLASSIC® GRANDE HERBICIDE TANK MIXES:

Apply CLASSIC® Grande HERBICIDE at 36 grams per hectare, tank mixed with glyphosate (present as potassium salt, isopropylamine salt, or di-ammonium salt) and adjuvant (where required).

Companion	Application	Application	For Use On	Weeds	Application
-----------	-------------	-------------	------------	-------	-------------

Herbicide	Rate (grams acid equivalent per hectare)	Timing		Controlled	Leaf Stage or Application Height (cm)
glyphosate (present as potassium salt, isopropylamine salt, or di-ammonium salt)	900 g ae/ha	Pre-plant burndown	Conventional & glyphosate tolerant soybeans	Dandelion (season long control)	Up to bolting
				Annual Sow-thistle	2-5 leaf
				Common Lamb's Quarters	Up to 6 leaf
				Prickly Lettuce	1-7 leaf
				Yellow Nutsedge (suppression only)	Up to 3 leaf
glyphosate (present as potassium salt, isopropylamine salt, or di-ammonium salt) plus non ionic surfactant (Agral 90, Citowett Plus or Ag-Surf at 0.2%v/v)	900 g ae/ha	1-3 Trifoliolate	glyphosate tolerant soybeans	Annual Sow-thistle (Season Long Control)	Less than 15 cm tall and across
				Common Lamb's Quarters	Up to 6 leaf
				Dandelion (Season Long Control)	Less than 15 cm tall and across
				Prickly Lettuce	1-7 leaf
				Yellow Nutsedge (Season long control)	Up to 8 leaf

Classic® Grande Herbicide + metribuzin:

Apply CLASSIC® Grande HERBICIDE at 36 grams per hectare, tank mixed with a metribuzin-containing end use product listed below at 550 grams per hectare for residual season long control of Canada fleabane including glyphosate resistant biotypes in soybeans. This tankmix is to be applied pre-plant or post-plant preemergence. To ensure control of emerged Canada fleabane seedlings, tank-mix pre-plant applications with a burndown partner such as 2,4-D Ester or Eragon. Pre-plant applications tank mixed with 2,4-D Ester must be applied a minimum 7 days before planting soybeans. Classic® Grande Herbicide plus metribuzin can be applied with glyphosate if required.

Tank mix partners:
TriCor MX Herbicide
TriCor 75 DF Herbicide
Lexone DF Herbicide
Sencor 75 DF Herbicide

Consult the respective glyphosate herbicide labels for specific instructions for 'Directions for Use', Precautions, and for other weeds controlled by glyphosate herbicide.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact E.I du Pont Canada Company at 1-800-667-3925 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by E.I du Pont Canada Company.

If rain occurs after application, control may be reduced. Two - four hours of dry weather are needed to allow CLASSIC® Grande Herbicide to be absorbed by weed foliage.

WEEDS CONTROLLED BY CLASSIC® GRANDE HERBICIDE + ASSURE® II Herbicide:

For the control of Annual Grasses, Common Ragweed, Redroot Pigweed and Velvetleaf, apply a tank mixture of CLASSIC® GRANDE HERBICIDE + ASSURE® II Herbicide. Use CLASSIC® GRANDE HERBICIDE at 36 grams per hectare plus ASSURE II® Herbicide at 0.38 to 0.5 litres per hectare. Add CANPLUS* 411 at 1.0% v/v (1 litre per 100 litres of spray solution) or Sure-Mix* at 0.5% v/v (0.5 litre per 100 litres of spray solution).

Use the 0.5 litre per hectare rate of ASSURE® II Herbicide when quackgrass and yellow foxtail are present in the field. The tank mix of Classic® Grande Herbicide and Assure II Herbicide may result in reduced control of these two grassy weeds

Consult the ASSURE® II Herbicide label for specified rates and optimum leaf stage timing for grassy weeds. If leaf stages of the grass and broadleaf weeds do not coincide, a sequential application of the grass and broadleaf herbicides is required to ensure satisfactory control.

Consult the ASSURE® II label for general information and use precautions.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT

FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than E. I. du Pont Canada Company and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. E. I. du Pont Canada Company, itself makes no representation or warranty with respect to product performance (efficacy) and crop tolerance (phytotoxicity) claims for this product when used on the crops listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold E.I. du Pont Canada Company harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.

SOYBEANS (Eastern Canada):

POSTEMERGENCE: Apply 36 g/ha CLASSIC® 25 DF Herbicide + 0.2% v/v of a non-ionic surfactant + 2 L/ha liquid nitrogen fertilizer at 28% UAN for the suppression of wild carrot (*Daucus carota*) and volunteer adzuki beans (*Vigna angularis* Ohwi & Ohashi). Make only one application per year. Apply to soybeans in the 1-3 trifoliolate stage of growth; apply to wild carrot when in the 2-20 leaf stage of growth and apply to volunteer adzuki beans when in the 1-3 trifoliolate stage of growth.

Apply in a spray volume of 140-190 L water/ha.

Do not apply within 60 days of harvest.

Refer to other sections of this label for additional application instructions, soil limitations and/or use precautions. The addition of liquid fertilizer containing 28% nitrogen to the spray mixture at the rate of 2 L/ha is required for the suppression of volunteer adzuki beans and wild carrots and it may also improve control of velvetleaf.

MIXING INSTRUCTIONS:

1. Fill clean tank about 1/4 full with fresh water.
2. Turn on full agitation.
3. Add the proper amount of CLASSIC® GRANDE HERBICIDE to the water in the spray tank with the agitator running. Maintain full agitation until product is fully dispersed. Continuous agitation is required to keep the product in suspension.
4. After CLASSIC® GRANDE HERBICIDE has been well mixed and is in suspension, add the tank mix partner, if applicable.
5. Add a recommended amount of spray adjuvant (if required).
6. Fill the remainder of the spray tank with clean water.

On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of the volume about to be mixed.

Agitation is required for uniform mixing and application. The optimum water volume for CLASSIC® GRANDE HERBICIDE application is 140-190 litres of water per hectare (minimum of 100 litres of water per hectare). Flat fan nozzles are recommended. Use 50 mesh filter screens or larger. Use spray preparation of CLASSIC® GRANDE HERBICIDE within 24 hours or product degradation may occur resulting in a loss of weed control. Use vigorous agitation to thoroughly disperse spray mixtures that have been allowed to stand in the tank.

NOTE: CLASSIC® GRANDE HERBICIDE will degrade in acidic or highly alkaline water. Mix no more than can be used in one day. If spraying is interrupted, thoroughly re-agitate the spray mixture before resuming spraying.

Do not mix or load where there is potential to contaminate wells or aquatic systems.

SPRAYER CLEANUP:

To avoid subsequent injury to crops other than soybeans immediately after spraying and prior to spraying other crops, thoroughly remove all traces of CLASSIC® GRANDE HERBICIDE from mixing and spray equipment as follows:

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes. Visually inspect tank to assure removal of all visible residues of CLASSIC® Grande Herbicide. If necessary, repeat step 1. **DO NOT CLEAN SPRAY EQUIPMENT NEAR WELLS, AQUATIC SYSTEMS, OR OTHER WATER SOURCES, OR NEAR DESIRABLE VEGETATION.**
2. Fill tank with clean water while adding 1 litre household ammonia (containing a minimum 3% ammonia) per 100 litres of water. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.

3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

REPLANTING TO OTHER CROPS:

CLASSIC® GRANDE HERBICIDE is degraded by natural soil processes, and field tests have shown that the following crops may safely be planted at the prescribed interval following soybeans that have been treated with CLASSIC® GRANDE Herbicide. The recropping intervals are dependent on the pH of the soil, as well as the sensitivity of the specific rotational crop.

Replanting Interval (Months)

Soil pH	Winter Wheat	Tomatoes	Field Corn, Soybeans	White Bean, Alfalfa	For Southern Ontario only: Cabbage, Garden Pea, Sweet Corn*
<7.0	3	12	10	10	11
>7.0<7.4	3	12	10	10	Do Not Plant
>7.4<7.8	4	12	10	Do Not Plant	
>7.8<8.0	4	12	Do Not Plant		

* Warning: Sweet corn varieties may vary in their sensitivity to chlorimuron ethyl residues. Soil pH may vary significantly within a single field. It may be affected by soil amendments (i.e. lime, fertilizers, etc.) or soil conditions (i.e. eroded knolls, etc.).

For other crops, a field bioassay is recommended before planting. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year.

FIELD BIOASSAY:

Select a representative area or areas of the field previously treated with CLASSIC® GRANDE HERBICIDE to plant your bioassay crop(s). Be sure to consider factors such as size of field, soil texture, drainage and turn-around areas when selecting the site(s) that are most representative of the soil conditions in the field. On large fields, more than one site may be needed in order to obtain reliable results.

Plant the test strips perpendicular to the direction in which the field was sprayed. The strips should be long enough to cross the width of several spray swaths. Large test strip areas are more reliable than small ones.

Use standard tillage and seeding equipment to plant the bioassay. Prepare a seed bed and plant the crops and varieties you want the option of growing the following year. It is important to use the same planting time, conditions, techniques and cultural practices you normally use to plant and grow the bioassay crop(s). Also plant into an adjacent area not treated with CLASSIC® GRANDE HERBICIDE to use as a comparison.

As the crop(s) emerges and grows, examine these key points in CLASSIC® GRANDE HERBICIDE treated and non-treated areas:

- crop stand
- plant colour and vigour
- root development
- yield
- rate of growth

Allow the bioassay crop(s) to grow to maturity while making your observations. Do not overspray the test strips with herbicides that may damage the bioassay crop(s). If the bioassay indicates that CLASSIC®

GRANDE HERBICIDE residues are still present, continue cropping only to those crops listed on the label and do not rotate to other crops until bioassay results indicate that susceptible crops are growing normally.

USE PRECAUTIONS:

Do not apply CLASSIC® GRANDE HERBICIDE during periods of intense rainfall or to soils saturated with water. Do not apply directly to standing or running water. Do not apply in areas where surface water from the treatment site can run off to adjacent cropland, either planted or to be planted, or into bodies of water (such as streams, ponds, rivers, lakes, and irrigation water), wetlands, or wells. Applications should only be made when there is no hazard of spray drift contaminating non-target land areas since very small quantities of the CLASSIC® GRANDE HERBICIDE spray solution may severely injure susceptible crops and other sensitive terrestrial habitats during both growing and dormant periods.

Do not apply to soybeans that have been stressed by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage as crop injury may result.

Under extreme weather conditions, such as hot, dry weather, excessive moisture, or frost, weed control may be reduced.

Rainfall within 4 hours after application of CLASSIC® GRANDE HERBICIDE may reduce weed control.

Because soybean varieties differ in their tolerance to herbicides, limit first use of CLASSIC® GRANDE HERBICIDE to a small area of each variety prior to adoption as a field practice.

Overspray or drift to important aquatic and terrestrial wildlife habitats should be avoided. Leave a 16 metre buffer zone between the last spray swath and bodies of water such as ponds, lakes, rivers, and streams. Leave a 39 metre buffer zone between the last spray swath and wetlands, vegetated areas at the edges of bodies of water, woodlots and shelterbelts, and other terrestrial wildlife habitats.

CLASSIC® GRANDE HERBICIDE must not be applied within 60 days of harvest.

Make only one application of CLASSIC® GRANDE HERBICIDE per season.

Do not graze the treated crop or cut for hay; sufficient data is not available to support such use.

Do not apply CLASSIC® GRANDE HERBICIDE to soybean grown on soil of pH \geq 8.0 due to risk of rotational crop injury.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, CLASSIC® GRANDE HERBICIDE is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to CLASSIC® GRANDE HERBICIDE and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of CLASSIC® GRANDE HERBICIDE or other Group 2 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.

- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local E. I. du Pont Canada Company representative or the E. I. du Pont Canada Company hotline at 1-800-667-3925.

®/™ Registered trademarks/trademark of E. I. du Pont Company. E. I. du Pont Canada Company is a licensee. Member of CropLife.

* All other products mentioned are trademarks of their respective companies

2013-10-09
2013-4471

Container label

GROUP	2	HERBICIDE
-------	---	-----------

DuPont™ Imazethapyr 240 SL

Herbicide Solution

COMMERCIAL (AGRICULTURAL)

GUARANTEE: Imazethapyr 240 g/L

REGISTRATION NO. 31156
PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BROCHURE BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 1-1050 litres

IN CASE OF SPILLS, POISONING OR FIRE TELEPHONE EMERGENCY
RESPONSE NUMBER 1-800-441-3637 (24 HOURS A DAY).

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS

- KEEP OUT OF REACH OF CHILDREN.
- This product may be harmful if swallowed, inhaled or absorbed through the skin.
- Do not contaminate food or feed products.
- Avoid contact with eyes, skin and clothing.
- Avoid breathing vapour or spray mist. Use with adequate ventilation.
- May cause eye damage. May cause skin irritation.
- Wear dust and/or splash-proof goggles or face shield and chemically resistant gloves when mixing, loading and during application, clean-up and repair.
- Wear long-sleeved shirt and long-legged pants when handling.
- Do not eat, drink or smoke when using.
- Wash hands and face before eating, drinking, smoking and using the toilet.
- Wash thoroughly with soap and water after handling.
- Take a shower IMMEDIATELY after work.
- Store and wash all protective clothing separately from normal laundry.
- Clean protective equipment (gloves, goggles, face shield) upon removal with soapy water.
- Clean spray equipment thoroughly after use. Wash in detergent and hot water before reuse.
- Wear freshly laundered clothes daily.
- If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.
- DO NOT APPLY BY AIR.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

1. Store the leftover product in original tightly closed container.
2. Keep product from freezing. DO NOT store below 0°C. If the product is exposed to temperatures below 0°C during shipment or storage, make sure the product has thawed completely, and shake the container vigorously.
3. DO NOT ship or store the product near food, feed, seed and fertilizers.
4. Store the product in cool, dry, locked, well-ventilated areas without floor drain.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

Booklet label

GROUP	2	HERBICIDE
-------	---	-----------

DuPont™ Imazethapyr 240 SL

Herbicide Solution

COMMERCIAL (AGRICULTURAL)

GUARANTEE: Imazethapyr 240 g/L

REGISTRATION NO. 31156
PEST CONTROL PRODUCTS ACT

READ THE LABEL AND THIS BROCHURE BEFORE USING
KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 1-1050 Litres

IN CASE OF SPILLS, POISONING OR FIRE TELEPHONE EMERGENCY
RESPONSE NUMBER 1-800-441-3637 (24 HOURS A DAY).

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

DIRECTIONS FOR USE

GENERAL INFORMATION

DuPont (tm) Imazethapyr 240 SL is a selective herbicide that can be applied as an early pre-plant, pre-plant incorporated, pre-emergent or post-emergent treatment in various crops. The application method depends upon the crop, anticipated weed spectrum and the preference of the applicator. With early pre-plant and pre-emergent treatments, susceptible weeds emerge, are present as stunted plants and then die. When DuPont (tm) Imazethapyr 240 SL is applied post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

Eastern Canada Directions:

Use DuPont (tm) Imazethapyr 240 SL at 312 mL (75 g active) – 420 mL (100 g active) in 400 L of water per hectare. See appropriate rate tables for specific application rates for each crop.

NOTE: A NONIONIC SURFACTANT CONTAINING AT LEAST 80% ACTIVE INGREDIENT (e.g. AGRAL® 90, AG-SURF®) at the rate of 0.25% (v/v) MUST BE ADDED to the spray solution (2.5 L of surfactant per 1000 L of spray solution).

REGISTERED CROPS

Adzuki Beans

Alfalfa Grown for Seed Production

Dry Common Beans (black, Dutch brown, kidney, white, yellow eye and cranberry beans only)

Lima Beans (Ontario only)

Processing Peas

Snap Beans

Snow Peas

Soybeans

MOISTURE REQUIREMENTS

As with most soil-applied herbicides, pre-emergent applications of DuPont (tm) Imazethapyr 240 SL require moisture for activation. Soil-applied DuPont (tm) Imazethapyr 240 SL requires sufficient water within 7 days of application to moisten the soil to a depth of 5 cm for activation. If adequate moisture is not received within 7 to 10 days of application, perform a shallow inter-row cultivation 5-8 cm deep using a roller or S-tine cultivator to control escaped weeds until the field receives adequate moisture. For early pre-plant applications (soybeans only), more than 7-10 days may elapse before the receipt of adequate precipitation to activate the herbicide and reduce the risk of weed escapes. Growers preferring surface applications of herbicides may choose this type of application of DuPont (tm) Imazethapyr 240 SL.

PLANT BACK RESTRICTIONS AND ROTATIONAL CROPS

In cases of crop failure, replant only soybeans, kidney beans, cranberry beans, Dutch brown beans, black beans, yellow eye beans, white beans, lima beans, adzuki beans and processing peas in the year of application. Winter wheat may also be re-planted in cases of crop failure or as a rotational crop 100 days

following a DuPont (tm) Imazethapyr 240 SL application. Soil preparation for re-planting should be no deeper than 10 cm.

Field corn, soybeans, winter wheat, spring wheat, spring barley, kidney beans, cranberry beans, Dutch brown beans, black beans, yellow eye beans, white beans, lima beans, adzuki beans and processing peas may be planted the season following a DuPont (tm) Imazethapyr 240 SL application. Conduct a field bioassay (a test strip grown to maturity) the year BEFORE growing any other crop.

APPLICATION INSTRUCTIONS

CROP: SOYBEANS

Apply DuPont (tm) Imazethapyr 240 SL as an early pre-plant, pre-plant incorporated, pre-emergent or postemergent treatment in soybeans.

For fields that contain weeds other than those listed in the “Weed Control in Soybeans: DuPont (tm) Imazethapyr 240 SL Alone” table, tank mix for broad-spectrum weed control (see “Herbicide Tank Mix Options – Soybeans”). The choice of product for tank mixing will depend on the specific weed(s) to be controlled. Consult the labels of the tank mix products to determine which product will provide control of the specific weeds present in the field.

Weed Control in Soybeans: DuPont (tm) Imazethapyr 240 SL Alone

Weeds	Application Timing					
	Early Pre-plant (prior to weed emergence)	Early Pre-plant (emerged weeds prior to 2 leaf stage)	Pre-plant Incorporated	Pre-Emergent	Early Post-emergent (before weeds reach 2 leaf stage)	Post-emergent (maximum leaf stage)
Broadleaf weeds						
Lamb’s quarters	C	PC	C ²	C	PC	
Redroot pigweed	C	C	C	C	C	C(12)
Smartweed	C			C		
Lady’s thumb	C			C ³		
Wild mustard	C	C	C	C	C	
Velvetleaf	C ¹	C ¹	C ¹	C ¹	C ¹	C(8)
Ragweed, common			PC ²	C ³		
Ragweed		PC			C ⁴	
Eastern black nightshade	PC	C	C	C	C	
Wild buckwheat		PC			C	
Cocklebur		C ¹			C ^{1,4}	
Grasses						
Foxtail, green and yellow	C	C	C	C	C	C(4)
Barnyard grass	C	PC	PC ²	C ³	C ⁴	C(6)
Old witchgrass	C			C ³	C	
Proso millet	PC	PC	PC	PC	PC	
Crabgrass, large					PC	
Perennials						
Yellow nutsedge					PC	

C = Control PC = Partial Control and Reduces Competition

1 Some plants of velvetleaf and/or cocklebur that germinate deeper in the soil and emerge late may escape treatment.

2 Tank mixing is recommended for fields with a history of heavy infestations of this weed species.

3 The higher label rate or tank mixing is recommended for fields with a history of heavy infestations of this weed species.

4 The higher label rate is required for heavy infestations of this weed species.

Herbicide Tank Mix Options - Soybeans

Tank Mix Option	Application Timing			
	Early Pre-plant	Pre-plant Incorporated	Pre-Emergent	Post-emergent
Gramoxone®	X ¹			
Roundup® ⁴ or Glyfos®	X ¹			X ^{1,3}
Roundup + FirstRate®	X ²			
Sencor®/Lexone®		X ^{1,2}	X ¹	
Treflan®/Rival®/Trifluralin® or Edge		X ¹		
Lorox®/Afolan®/Linuron®			X ¹	
Basagran®				X ^{1,2}
Basagran® Forte				X ^{1,2}

1 Refer to the label of the specific tank mix product for information regarding: rates, recommendations, precautions and restrictions.

2 Refer to Tank Mix Options under the appropriate Application Timing in this label.

3 Glyphosate Tolerant Soybeans only (i.e., varieties with the Roundup Ready®). Refer to CROP: GLYPHOSATE TOLERANT gene SOYBEANS section.

4 Roundup Original Liquid Herbicide, Roundup Transorb™ Liquid Herbicide or Roundup WeatherMax™ with Transorb 2 Technology Liquid Herbicide.

Timing	Early Pre-plant Application - Soybeans Apply DuPont (tm) Imazethapyr 240 SL up to 30 days before planting in conventional, reduced tillage or no-till soybeans. DuPont (tm) Imazethapyr 240 SL alone may be applied as a surface application using this technique. Only one additional working of the soil to prepare the seedbed is recommended following the application. This final seedbed preparation should not work the soil deeper than 10 cm. Deeper tillage will result in reduced concentration of herbicide in the weed germination zone and reduction in weed control. DO NOT plow following the application.	
Rate	Early Pre-plant – Prior to Weed Emergence	420 mL/ha
	Early Pre-plant – to Emerged Weeds (before the weeds reach the 2 true leaf stage).	420 mL/ha + non-ionic surfactant. Nonionic surfactant MUST BE ADDED. Liquid fertilizer added to the spray solution will provide quicker burndown of weed
Water Volume	100-400 L/ha	
Surfactant/Adjuvant	For Early Pre-plant – to Emerged Weeds: Non-ionic surfactant – 0.25% v/v (e.g. 2.5 L/1000 L of spray solution) Liquid fertilizer solution - 2 L/ha	
Weeds Controlled	See “Weed Control in Soybeans” table above.	
Remark	Plant only soybeans during the season of application.	

Tank Mix Options	<p>DuPont (tm) Imazethapyr 240 SL may also be tank mixed with the herbicides listed in the “Herbicide Tank Mix Options – Soybeans” table. Always follow the most conservative rates, recommendations, precautions and restrictions on the tank mix labels.</p> <p>For the combined tank mix with ROUNDUP and FIRSTRATE, use the following:</p> <table border="1" style="margin-left: 40px;"> <tr> <td>Rates</td> <td>DuPont (tm) Imazethapyr 240 SL – 312 mL/ha ROUNDUP – 2.5 L/ha (0.900 kg ai/ha) FIRSTRATE – 21 g/ha (0.0175 kg ai/ha)</td> </tr> <tr> <td>Water Volume</td> <td>100-400 L/ha</td> </tr> </table> <p>Comments: This tank mix provides control of emerged weeds found on the ROUNDUP herbicide label and residual control of germinating weeds found on the DuPont (tm) Imazethapyr 240 SL and FIRSTRATE herbicide labels. Temporary crop injury may occur, however, yield will not normally be affected. Avoid sprayer overlap. Severe crop injury will occur.</p>	Rates	DuPont (tm) Imazethapyr 240 SL – 312 mL/ha ROUNDUP – 2.5 L/ha (0.900 kg ai/ha) FIRSTRATE – 21 g/ha (0.0175 kg ai/ha)	Water Volume	100-400 L/ha
Rates	DuPont (tm) Imazethapyr 240 SL – 312 mL/ha ROUNDUP – 2.5 L/ha (0.900 kg ai/ha) FIRSTRATE – 21 g/ha (0.0175 kg ai/ha)				
Water Volume	100-400 L/ha				

Timing	<p>Pre-plant Incorporated Application - Soybeans</p> <p>Incorporate DuPont (tm) Imazethapyr 240 SL evenly throughout the top 5 cm of the soil profile. Incorporation may be achieved with a double pass using discs or cultivator operated at 8 to 12 kph with the second pass at an angle to the first. Cultivators must have 3 or 4 rows of flexible sweeps staggered and spaced less than 15 cm apart followed by a drag or rolling basket to ensure no soil is left unturned.</p>											
Rate	312 mL/ha											
Water Volume	100-400 L/ha											
Weeds Controlled	See “Weed Control in Soybeans” table above.											
Remark	DO NOT apply DuPont (tm) Imazethapyr 240 SL as a pre-plant incorporated application more than one year in sequence. Allow at least 24 months between pre-plant incorporated applications.											
Tank Mix Options	<p>For fields that contain heavy lamb's-quarters, common ragweed and/or barnyard grass infestations or weeds other than those listed under “DuPont (tm) Imazethapyr 240 SL Alone”, tank mixing may be required to provide broad-spectrum weed control (see “Herbicide Tank Mix Options – Soybeans” table). Follow the most conservative rates, recommendations, precautions and restrictions on the tank mix labels. For SENCOR and LEXONE herbicide, refer to the following table for rates:</p> <table border="1" style="margin-left: 40px;"> <thead> <tr> <th rowspan="2">Soil Texture¹</th> <th colspan="2">Application Rate (g/ha)</th> </tr> <tr> <th>SENCOR 75DF</th> <th>LEXONE 75 DF</th> </tr> </thead> <tbody> <tr> <td>Medium (loam, silt loam, silt, sandy clay, sandy clay loam)</td> <td>750</td> <td>540</td> </tr> <tr> <td>Heavy (silty clay, silty clay loam, clay and clay loam)</td> <td>750</td> <td>640</td> </tr> </tbody> </table> <p>¹ Do not use on light (loamy sand, sandy loam) textured soils. Do not use on soil with less than 2% organic matter.</p>	Soil Texture ¹	Application Rate (g/ha)		SENCOR 75DF	LEXONE 75 DF	Medium (loam, silt loam, silt, sandy clay, sandy clay loam)	750	540	Heavy (silty clay, silty clay loam, clay and clay loam)	750	640
Soil Texture ¹	Application Rate (g/ha)											
	SENCOR 75DF	LEXONE 75 DF										
Medium (loam, silt loam, silt, sandy clay, sandy clay loam)	750	540										
Heavy (silty clay, silty clay loam, clay and clay loam)	750	640										

Timing	Pre-emergent Application - Soybeans Pre-emergent applications of DuPont (tm) Imazethapyr 240 SL may be applied before the crop and weeds emerge.	
Rate	312 - 420 mL/ha	
Water Volume	100-400 L/ha	
Weeds Controlled	See “Weed Control in Soybeans” table above.	
Tank Mix Options	For fields that contain heavy infestations of common ragweed, old witchgrass, barnyard grass, lady’s-thumb or weeds other than those listed under “DuPont (tm) Imazethapyr 240 SL Alone”, tank mixing may be required to provide broad-spectrum weed control (see “Herbicide Tank Mix Options – Soybeans” table). Follow the most conservative rates, recommendations, precautions and restrictions on the tank mix labels.	
Timing	Early and Late Post-emergent Application - Soybeans Apply DuPont (tm) Imazethapyr 240 SL early post-emergent after the crop has emerged and before the weeds reach the 2 true leaf stage. DuPont (tm) Imazethapyr 240 SL may also be applied late post-emergent after the crop has emerged for control of certain weed species up to the growth stages indicated in the “Weed Control in Soybeans: DuPont (tm) Imazethapyr 240 SL Alone” table. To minimize weed competition with the crop, application should be made as early as possible after weed emergence.	
Rate	Early Post-emergent – (before the weeds reach the 2 true leaf stage)	312-420 mL/ha + non-ionic surfactant + liquid fertilizer
	Late Post-emergent	420 mL/ha + non-ionic surfactant + liquid Fertilizer.
Water Volume	100-400 L/ha	
Surfactant/ Adjuvant	Non-ionic surfactant – 0.25% v/v (e.g. 2.5 L/1000 L of spray solution) Liquid fertilizer solution (10-34-0, 28-0-0 or 32-0-0) - 2 L/ha	
Weeds Controlled	See “Weed Control in Soybeans” table above.	
Remark	Addition to the spray solution of a non-ionic surfactant plus liquid fertilizer is essential for post-emergent application, improving uptake of the product by weeds resulting in improved herbicidal activity. Avoid applications when weeds and/or crop are under growth stress. Stunting of the crop may result following post-emergence application. This condition is the result of stem inter-node shortening but will not result in yield reduction.	
Tank Mix Options	DuPont (tm) Imazethapyr 240 SL may also be tank mixed with the herbicides listed in the “Herbicide Tank Mix Options – Soybeans” table. Always follow the most conservative rates, recommendations, precautions and restrictions on the tank mix labels. Comments: Use the 312 mL/ha rate of DuPont (tm) Imazethapyr 240 SL when tank mixed with Basagran or Basagran Forte herbicide. For DuPont (tm) Imazethapyr 240 SL plus BASAGRAN herbicide, a non-ionic surfactant at the rate of 0.25% (v/v) and fertilizer solution at the rate of 2 L/ha MUST BE ADDED to the spray solution. For DuPont (tm) Imazethapyr 240 SL plus BASAGRAN FORTE herbicide, only fertilizer solution MUST BE ADDED at the rate of 2 L/ha.	

CROP: GLYPHOSATE TOLERANT SOYBEANS
(i.e., varieties with the Roundup Ready® gene)

Timing	Post-emergent Application For broadleaf and grass weeds other than those listed in “Weed Control in Soybeans: DuPont (tm) Imazethapyr 240 SL Alone” table, DuPont (tm) Imazethapyr 240 SL may be tank mixed with ROUNDUP herbicide and applied to Roundup Ready soybeans. Follow the rates, application timings, recommendations, precautions and restrictions on the ROUNDUP herbicide label. Refer to the ROUNDUP herbicide label for adjuvant recommendations.
Rate	312 mL/ha
Water Volume	100-400 L/ha
Remarks	WARNING Apply DuPont (tm) Imazethapyr 240 SL + ROUNDUP herbicide tank mix ONLY to glyphosate tolerant soybeans, i.e. varieties with the Roundup Ready® gene. SOYBEANS WHICH ARE NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.

CROP: EDIBLE BEANS (including kidney beans, cranberry beans, Dutch brown beans, black beans, lima beans (Ontario only), yellow eye beans, white beans and adzuki beans)

Apply DuPont (tm) Imazethapyr 240 SL as a pre-emergent treatment in kidney beans, cranberry beans, Dutch brown beans, black beans, yellow eye beans, white beans, adzuki beans and lima beans (Ontario only), and as a pre-plant incorporated treatment in white beans, kidney beans, cranberry beans and adzuki beans.

Weed Control in Edible Beans: DuPont (tm) Imazethapyr 240 SL Alone

Weeds	Application Timing	
	Pre-Emergent	Pre-plant Incorporated (kidney beans, cranberry beans, white beans and adzuki beans only)
Broadleaf weeds		
Lamb’s quarters	C	C
Redroot pigweed	C	C
Smartweed	C	C
Lady’s thumb	C	C
Wild mustard	C	C
Velvetleaf	C ¹	C ¹
Ragweed, common	PC	PC
Eastern black nightshade	C	C
Grasses		
Foxtail, green and yellow	C	C
Proso millet	PC	PC

C = Control PC = Partial Control and Reduced Competition

¹ Some plants of velvetleaf that germinate deeper in the soil and emerge late may escape treatment.

Timing	Pre-emergent Application – Edible Beans
Rate	312 mL/ha
Water Volume	100-400 L/ha
Tanks Mix Options	For fields that contain heavy infestations of broadleaf weeds other than those listed under “DuPont (tm) Imazethapyr 240 SL Alone”, tank mixing may be required to provide broad-spectrum weed control. Cranberry and Kidney Beans Tank mix DuPont (tm) Imazethapyr 240 SL with 1.15 – 1.75 L/ha (1.05-1.60 kg ai/ha) of DUAL® MAGNUM herbicide to control labelled broadleaf weeds and grasses in cranberry beans and kidney beans. Refer to the DUAL MAGNUM herbicide label for additional recommendations, precautions and restrictions not specified on this label.

Timing	Pre-plant Incorporated Application (kidney beans, cranberry beans, white beans; adzuki beans in Eastern Canada only)
Rate	312 mL/ha
Water Volume	100-400 L/ha
Tanks Mix Options	White Beans Tank mix DuPont (tm) Imazethapyr 240 SL with 1.25-2.4 L/ha of TREFLAN® herbicide to control a broader spectrum of weeds in white beans only. Follow recommendations, precautions and restrictions on the TREFLAN herbicide label. Consult the tank mix partner label for weeds controlled other than those listed for DuPont (tm) Imazethapyr 240 SL alone. Cranberry and Kidney Beans Tank mix DuPont (tm) Imazethapyr 240 SL with 1.15 – 1.75 L/ha (1.05-1.60 kg ai/ha) of DUAL MAGNUM herbicide to control labelled broadleaf weeds and grasses in cranberry beans and kidney beans. Refer to the DUAL MAGNUM herbicide label for additional recommendations, precautions and restrictions not specified on this label.

CROP: PROCESSING PEAS

Timing	Pre-emergent or Pre-plant Incorporated Application
Rate	312 mL/ha
Water Volume	200 L/ha
Weeds Controlled	See “DuPont (tm) Imazethapyr 240 SL Alone: Weed Control in Soybeans”

CROP: SNOW PEAS

Timing	Pre-emergent or Pre-plant Incorporated Application
Rate	312 mL/ha
Water Volume	100-400 L/ha
Weeds Controlled	See “DuPont (tm) Imazethapyr 240 SL Alone: Weed Control in Soybeans”

CROP: SNAP BEANS

Timing	Pre-emergent or Pre-plant Incorporated Application
Rate	312 mL/ha
Water Volume	200 L/ha
Weeds Controlled	See “DuPont (tm) Imazethapyr 240 SL Alone: Weed Control in Soybeans”

CROP: ALFALFA GROWN FOR SEED PRODUCTION

Timing	Pre-emergent or Pre-plant Incorporated Application
Rate	312-420 mL/ha
Water Volume	200 L/ha
Surfactant/ Adjuvant	A non-ionic surfactant at the rate of 0.25% (v/v) MUST BE ADDED to the spray solution (e.g., 2.5 L of surfactant per 1000 L of spray solution). Fertilizer Solutions MUST BE ADDED to the spray solution at the rate of 2 L/ha to provide quicker burndown of weeds.
Weeds Controlled	See “DuPont (tm) Imazethapyr 240 SL Alone: Weed Control in Soybeans”

MIXING INSTRUCTIONS

1. Ensure the spray tank is clean before use. Follow the clean-out recommendations stated on the label of the product that was previously used.
2. Fill the spray tank one-half full to three-quarters full of water and start agitation.
3. Using a calibrated measuring device, add the required amount of tank mix partner (refer to the tank mixture section of each crop for tank mixtures).
4. Mix thoroughly.
5. Using a separate calibrated measuring device, add the required amount of DuPont (tm) Imazethapyr 240 SL to the tank while agitating the spray solution.
6. While the solution remains agitating, add the required amount of non-ionic surfactant if required.
7. If required, add the required amount of liquid fertilizer (28-0-0, 10-34-0 or 32-0-0) to the spray solution.
8. Continue agitation while filling the remainder of the spray tank with water.
9. Clean the spray tank after use.

ENVIRONMENTAL PRECAUTIONS

DuPont (tm) Imazethapyr 240 SL is toxic to aquatic organisms.

DO NOT apply this product directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches, and wetlands), estuaries, or marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

RESTRICTIONS AND LIMITATIONS

1. DO NOT APPLY DUPONT (TM) IMAZETHAPYR 240 SL BY AIR. APPLY WITH GROUND EQUIPMENT ONLY.
2. DO NOT over apply DuPont (tm) Imazethapyr 240 SL. Over application may result in injury particularly if the crop is under stress.
3. Crop Pre-harvest Interval

Crop	Application to Harvest Interval (days)
Soybeans	100
Dry Beans ¹	100
Lima Beans	90
Processing Peas	50
Snow Peas	60
Snap Beans	40

¹ kidney, adzuki, Dutch brown, black, yellow eye, white and cranberry beans

4. CAUTION: Do not graze treated crops or cut for hay, sufficient data are not available to support such use.

5. DO NOT let spray drift contaminate crops in adjacent fields.

6. ONLY ONE (1) application of DuPont (tm) Imazethapyr 240 SL may be made during the season.

7. DO NOT apply DuPont (tm) Imazethapyr 240 SL as a pre-plant incorporated application in all crops more than one year in sequence. Allow at least 24 months between pre-plant incorporated applications.

8. Emerged weeds in pre-emergent applications which reach the 2 - 3 true leaf stage might be considered as escapes. Shallow cultivation or application of a post-emergent herbicide is recommended. DO NOT cultivate deeply.

9. DO NOT apply DuPont (tm) Imazethapyr 240 SL when crop is under stress conditions because crop injury may result.

10. Post-emergent application of DuPont (tm) Imazethapyr 240 SL to soybeans may cause stunting. This condition is the result of stem inter-node shortening and does not cause yield reductions if DuPont (tm) Imazethapyr 240 SL has been used at label rates and following label recommendations.

11. DuPont (tm) Imazethapyr 240 SL may cause stunting or delayed maturity in white beans and kidney beans. Stunting is the result of stem inter-node shortening and should not cause yield reductions if DuPont (tm) Imazethapyr 240 SL has been used at label rates and following label recommendations.

12. Over-spray or drift to important wildlife habitats such as shelterbelts, water-bodies, wetlands, woodlots, vegetated ditch-banks, hedgerows and other cover on the edge of fields should be avoided. Leave a 15-meter buffer zone between the last spray swath and the edge of any of these habitats.

Western Canada Directions:

DIRECTIONS FOR USE ON FIELD PEAS

DuPont (tm) Imazethapyr 240 SL is a selective herbicide that can be applied as an early post-emergence treatment in field peas. When DuPont (tm) Imazethapyr 240 SL is applied early post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

SOIL ZONES: Black and grey wooded soils

CROPS: Field peas up until the sixth (6th) trifoliate leaf stage

ROTATIONAL CROPS

There is the possibility of residual soil activity from DuPont (tm) Imazethapyr 240 SL the year following application. Research studies have shown the following crops can be safely grown in black and grey wooded soil zones the year following a DuPont (tm) Imazethapyr 240 SL application:

Spring barley, spring wheat, lentils, alfalfa, field peas

Conduct a field bioassay (a test strip grown to maturity) the year BEFORE growing any other crop than those listed above.

RATES

Apply 50 g active (a.e.) in 100 to 400 litres of water per hectare. Equivalent to 210 mL of product per hectare.

NOTE: A NONIONIC SURFACTANT CONTAINING AT LEAST 80% ACTIVE INGREDIENT (e.g. AGRAL® 90, AG-SURF®) at the rate of 0.25% (v/v) MUST BE ADDED to the spray solution (2.5 L of surfactant per 1000 L of spray solution).

BROADLEAF WEED and GRASS CONTROL POST-EMERGENCE APPLICATIONS

DuPont (tm) Imazethapyr 240 SL, by early post-emergence application up to the 4 true leaf stage of the weeds, will control:

Chickweed	Cleavers
Green foxtail	Hemp-nettle
Redroot pigweed	Shepherd's purse
Stinkweed	Smartweed
Volunteer canola	Wild buckwheat**
Wild mustard	Wild oats*

* For control of wild oats apply DuPont (tm) Imazethapyr 240 SL between the 2-4 leaf stage of the weed.

** Suppression only

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with water. Using a calibrated measuring device, add the required amount of DuPont (tm) Imazethapyr 240 SL while agitating the spray solution. While the solution remains agitating, add the required amount of non-ionic surfactant. Fill the remainder of the tank with water.

WARNINGS

An interval of 60 days must follow the DuPont (tm) Imazethapyr 240 SL application before field peas are harvested.

Field peas treated with DuPont (tm) Imazethapyr 240 SL may be fed to livestock 30 days after application. DO NOT over-apply DuPont (tm) Imazethapyr 240 SL. Over application may result in injury particularly if the crop is under stress.

DO NOT let spray drift contaminate crops other than those being sprayed.

ONLY ONE (1) application of DuPont (tm) Imazethapyr 240 SL may be made during the season.

DO NOT tank-mix DuPont (tm) Imazethapyr 240 SL.

In cases of crop failure, only field peas may be re-planted in the year of application.

DO NOT APPLY BY AERIAL APPLICATION - GROUND APPLICATION ONLY.

Over-spray or drift to important wildlife habitats such as shelterbelts, water bodies, wetlands and woodlots, vegetated ditch banks, hedgerows & other cover on the edge of the field should be avoided. Leave a 15-meter buffer zone between the last spray swath and the edge of any of these habitats.

DIRECTIONS FOR USE ON NEWLY SEEDED PURE STAND ALFALFA FOR FORAGE OR SEED PRODUCTION

DuPont (tm) Imazethapyr 240 SL is a selective herbicide that can be applied as an early post-emergence treatment in newly seeded pure stand alfalfa in the year of establishment. DuPont (tm) Imazethapyr 240 SL may be applied when the crop has developed at least one (1) fully expanded trifoliate leaf and up to and including the 4 leaf stage of susceptible weeds. Do not spray before all weeds have emerged. When DuPont (tm) Imazethapyr 240 SL is applied early post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

CROPS

DuPont (tm) Imazethapyr 240 SL may be used on pure stand alfalfa in the year of establishment in the black, grey wooded and irrigated brown soil zones.

USE

DuPont (tm) Imazethapyr 240 SL is intended for use on pure stand alfalfa in the year of establishment, on stands that will remain for 3 or more years.

ROTATIONAL CROPS

In the event of crop failure, only field peas may be replanted during the same season as the DuPont (tm) Imazethapyr 240 SL application. Do not apply 2 applications of DuPont (tm) Imazethapyr 240 SL within the same year.

If necessary, spring wheat, lentils, field peas or alfalfa may be planted the season following a DuPont (tm) Imazethapyr 240 SL application in the black, grey wooded and irrigated brown soil zones. Barley may also be planted in the black and grey wooded soil zones the season following application.

Conduct a field bioassay (a test strip grown to maturity) BEFORE growing any other crop.

RATES

Apply 50 grams active (a.e.) in 100 to 400 litres of water per hectare.
Equivalent to 210 mL of product per hectare.

NOTE: For early post-emergence applications a NONIONIC SURFACTANT CONTAINING AT LEAST 80% ACTIVE INGREDIENT (AGRAL 90, AG-SURF) at the rate of 0.25% (v/v) MUST BE ADDED to the spray solution (e.g., 2.5 L of surfactant per 1000 L of spray solution).

BROADLEAF WEED and GRASS CONTROL

EARLY POST-EMERGENCE APPLICATIONS

DuPont (tm) Imazethapyr 240 SL, by early post-emergence application (up to and including the 4 leaf stage of susceptible weeds) will control:

Broadleaf weeds: Redroot pigweed, Wild mustard, Stinkweed, Volunteer canola and Green smartweed

In addition, DuPont (tm) Imazethapyr 240 SL provides partial control of and reduces competition from:

Common groundsel, Green foxtail, Shepherd's purse

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with water. Using a calibrated measuring device, add the required amount of DuPont (tm) Imazethapyr 240 SL while agitating the spray solution. While the solution remains agitating, add the required amount of non-ionic surfactant. Fill the remainder of the tank with water.

WARNINGS

Apply only in the year of establishment and only after the crop has one fully developed trifoliolate leaf.

DO NOT over-apply DuPont (tm) Imazethapyr 240 SL. Over application may result in injury particularly if the crop is under stress.

DO NOT let spray drift contaminate crops other than those being sprayed.

ONLY ONE (1) application of DuPont (tm) Imazethapyr 240 SL may be made during the life of the alfalfa stand.

DO NOT tank-mix DuPont (tm) Imazethapyr 240 SL.

In cases of crop failure, only field peas may be re-planted in the year of application.

DO NOT graze or harvest for forage until 14 days after treatment.

Mature seed from treated plants should not be used for human consumption.

Over-spray or drift to important wildlife habitats such as shelterbelts, water bodies, wetlands, woodlots, vegetated ditch banks, hedgerows and other cover should be avoided. Leave a 15-meter buffer zone between the last spray swath and the edge of any of these habitats.

DO NOT APPLY BY AERIAL APPLICATION - GROUND EQUIPMENT ONLY.

DIRECTIONS FOR USE ON ESTABLISHED ALFALFA FOR SEED PRODUCTION PURPOSES

CROP	WEEDS CONTROLLED	RATE	DIRECTIONS
Established alfalfa for seed production	Stinkweed, wild mustard, volunteer canola, redroot pigweed and suppression of green foxtail	50 g a.e./ha	One application per year; post-emergent; ground application; apply before alfalfa reaches 30 cm. Do not use DuPont (tm) Imazethapyr 240 SL in the last year of the alfalfa stand.

DIRECTIONS FOR USE ON DRY BEANS (PINTO, PINK, RED)

CROP	PEST	RATE	DIRECTIONS
Dry beans – pinto, pink, red	Hairy nightshade	50 g a.e./ha	One application per year; ground sprayer; apply in 100 - 400 L water/ha; post-emergence; apply up to the 2nd trifoliolate leaf stage of the dry bean; up to 6 leaf stage of hairy nightshade; black, grey wooded and irrigated brown soil zones only; PHI of 75 days.

DIRECTIONS FOR USE ON CHICKLING VETCH (FOR SEED PRODUCTION ONLY)

CROP	PEST	RATE	DIRECTIONS
Chickling vetch/ grass pea	Labelled weeds	50 g a.e./ha	One application per year; post-emergent; ground application; PHI 60 days; apply at the 5-7 leaf stage of chickling vetch/grass pea.

DIRECTIONS FOR USE ON SOYBEANS (MANITOBA ONLY)

CROP	PEST	RATE	DIRECTIONS
Soybeans	Annual grass and broadleaf weeds	50 g a.e./ha plus non-ionic surfactant at 0.25% v/v	Apply as post-emergent application once per season at 1 to 3 leaves of soybean growth; ground application only; PHI 85 days.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, DuPont (tm) Imazethapyr 240 SL is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to DuPont (tm) Imazethapyr 240 SL and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of DuPont (tm) Imazethapyr 240 SL or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local DuPont representative or the DuPont hotline at 1-800-667-3925.

PRECAUTIONS

- KEEP OUT OF REACH OF CHILDREN.
- This product may be harmful if swallowed, inhaled or absorbed through the skin.
- Do not contaminate food or feed products.
- Avoid contact with eyes, skin and clothing.
- Avoid breathing vapour or spray mist. Use with adequate ventilation.
- May cause eye damage. May cause skin irritation.

- Wear dust and/or splash-proof goggles or face shield and chemically resistant gloves when mixing, loading and during application, clean-up and repair.
- Wear long-sleeved shirt and long-legged pants when handling.
- Do not eat, drink or smoke when using.
- Wash hands and face before eating, drinking, smoking and using the toilet.
- Wash thoroughly with soap and water after handling.
- Take a shower IMMEDIATELY after work.
- Store and wash all protective clothing separately from normal laundry.
- Clean protective equipment (gloves, goggles, face shield) upon removal with soapy water.
- Clean spray equipment thoroughly after use. Wash in detergent and hot water before reuse.
- Wear freshly laundered clothes daily.
- If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.
- DO NOT APPLY BY AIR.

STORAGE

1. Store the leftover product in original tightly closed container.
2. Keep product from freezing. DO NOT store below 0°C. If the product is exposed to temperatures below 0°C during shipment or storage, make sure the product has thawed completely, and shake the container vigorously.
3. DO NOT ship or store the product near food, feed, seed and fertilizers.
4. Store the product in cool, dry, locked, well-ventilated areas without floor drain.

DISPOSAL

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.