

FURIOUS

I N S E C T I C I D E

ACTIVE INGREDIENT:

Geraniol Oil	30.00%
Peppermint oil	1.00%
Cottonseed oil	0.10%
Rosemary oil	0.01%

OTHER INGREDIENTS:* 68.89%

TOTAL: **100.00%**

*Canola Oil, Water

How can we help?

1-833-476-7761

MANUFACTURED BY:

GroPro[®]

12600 West Frontage Road,
Burnsville MN 55337

Phone: 1-833-476-7761

www.groproag.com

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

“SEE LABEL BOOKLET FOR PRECAUTIONARY
STATEMENTS”

**ENVIRONMENTALLY SAFE
When Used as Directed**

Intended for use by commercial applicators. For use in greenhouses, ornamentals, cannabis and on turf related to golf courses, sod farms, sport fields, residential, institutional, municipal, commercial, and other turf grass areas. This product has not been registered by the United States Environmental Protection Agency. GroPro represents that this product qualifies for exemption from registration under FIFRA.

APPROVED FOR USE ON LOW-THC CANNABIS OR MEDICAL MARIJUANA.

NET CONTENTS: ■ **64 fl oz (1.8925 L)**



TURF AND ORNAMENTAL

GroPro[®]

PRECAUTIONARY STATEMENTS

- Do not get in eyes, on skin or on clothing.
- Wash hands, forearms, and other exposed areas thoroughly after handling.
- Do not eat, drink or smoke when using this product.
- Do not breathe vapors, mist, or spray.

FIRST AID

IF IN EYES: • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • **Do not** induce vomiting unless told to do so by the poison control center or doctor. • **Do not** give anything to an unconscious person.

IF INHALED: • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice.

POISONING HOTLINE

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. **Do not** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity \leq 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

IN CASE OF EMERGENCY

In case of large-scale spill of this product, call:

- 911

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)

Steps to take if this material is released or spilled:

- In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

PRODUCT INFORMATION

FURIOUS® is an oil based concentrate formulation containing the various active ingredients that are classified as essential oil class of chemistry.

When used as directed, FURIOUS® provides knockdown and residual control of soft body insects on the crops listed on this label. FURIOUS® is a highly active contact insecticide on egg, nymph, and adult stage soft body insects. See Pests Controlled list for insects controlled.

FURIOUS® is compatible with other chemicals but a tank compatibility test is suggested as GROPRO has not been able to test every tank mix on the market.

FURIOUS® is not systemic or translaminar; therefore, thorough coverage of plant surfaces is important for optimum performance.

RESISTANCE MANAGEMENT

Geraniol, the main active ingredient in FURIOUS®, belongs to the group of essential oils classified under FIFRA 25B.

No known resistance is known to the essential oil class but to reduce the potential for developing resistance, rotate to a insecticides with a different mode of action. Monitor treated insect populations for resistance development. Read product label before applying any insecticide and follow label directions. Contact your local extension specialist, certified crop advisor, and/or GROPRO representative for additional resistance management or IPM recommendations.

PLANT TOLERANCE

Plant tolerance to FURIOUS® is not fully known, due to the large number of plant species and their associated varieties or cultivars, and due to variable growing conditions, it is impossible to test every plant for tolerance to this product.

Neither the manufacturer or seller has determined whether this product can be used without injury on all varieties of plants. It is the responsibility of the user to determine if this product can be used without causing plant injury prior to commercial use. In a small test area, test the specified rates on plants for phytotoxicity prior to large-scale use. The end user assumes all risks arising from application of FURIOUS® in a manner inconsistent with the label requirements.

PESTS CONTROLLED

When used according to label instructions, FURIOUS® is effective in controlling the following insects:

Common Name	
Annual bluegrass weevil	Black Margined Aphid
Asiatic Garden Beetle	Asian Citrus Physlid
Billbugs	Black Fly
Black Turfgrass atenius	Soft Scales
Cutworms	Avocado Lacebug
European Chafer	Pine Tip Moth Larvae
European Crane Fly	Royal Palm Bugs
Green June Beetle	Fungus Gnats
Japanese Beetle	Root Mealybug
Northern Masked Chafer	Root Weevil Complex
Chinch bugs	Banks Grass Mite
Mole Crickets	Brown Almond Mite
Aphids	Brown Wheat Mite
Thrips	Citrus Red Mite
Mites	European Reed Mite
Leafhoppers	Spider Mites
Glassy Winged Sharp Shooters	Texas Citrus
Sawfly	Armyworms
Mealy Bugs	Fruit Fly
Whiteflies	Seed Maggot
Leafminers	Root Maggot
Adelgids	Wire Worm
White Grub	Cucumber Beetle
San Jose Scales	
Black Vine Weevil Larvae	
Yellow Pecan Aphid	

APPLICATION INSTRUCTIONS

For maximum effectiveness, apply FURIOUS® at the first sign of mites before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing mite populations.

Apply to foliage using properly calibrated ground sprayers (refer to Crop-specific Application Instructions table). Thorough and uniform spray coverage of foliage, with direct contact to target pest, is required for effective control. Applications made at lower volumes may result in less than thorough coverage which can lead to slower activity and/or less control than those made with higher spray volumes. Consult with your local or state extension personnel for advice on insecticide use and selection.

Apply FURIOUS® in sufficient volume of water to ensure thorough coverage of foliage. For thorough coverage, use higher water volume.

Apply FURIOUS® in minimum water volume per acre as indicated in Crop-specific Application Instructions table. Because FURIOUS® is not systemic or translaminar, thorough coverage of plant surfaces is necessary for effective control. Applying FURIOUS® at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

CONTROLLING SPRAY DRIFT

Variable Winds. DO NOT apply in variable wind conditions.

Wind Direction. Wind direction (e.g. away from nontarget areas) must be based on the average direction (not instantaneous).

Wind Speed. Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy, and equipment specifications determine drift potential at any given wind speed. DO NOT apply when winds are greater than 15 mph.

Inversion Restriction. DO NOT make applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light-to-no wind.

Sensitive Areas. To ensure the protection of threatened or endangered species, it is important to maintain spray drift loadings below levels of concern for any area adjacent to the application site that is not excluded as possible habitat for these organisms. The following restrictions apply:

- Buffer distance - 15 feet
- Release height - 20 inches
- Droplet size - A combination of spray nozzles and appropriate pressure must be selected to provide ASABE standard S571.1 droplet size category of fine (DV0.5 of ≥ 180 microns) or coarser than fine.
- Maximum wind speed - 15 mph

CLEANING SPRAY EQUIPMENT

Before application, start with clean, well-maintained application equipment. Following spray application, thoroughly clean all application equipment. Drain application equipment of any excess product. Thoroughly rinse application equipment and flush hoses, boom, and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. **DO NOT** clean equipment near wells, water sources, or desirable vegetation. Dispose of waste rinsewater in accordance with local regulations. Spraying equipment must be cleaned thoroughly before and after applying this product, particularly if a product with the potential to injure crops was used before FURIOUS®.

ADDITIVES AND TANK MIXING INFORMATION

Shake container well before use. FURIOUS® can be tank mixed with most recommended fungicides, insecticides, liquid fertilizers, biological control products, adjuvants, and additives. The use of additives or adjuvants can improve the performance of FURIOUS®. However, all varieties and cultivars have not been tested with all possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which GROPRO has conducted testing. Physical incompatibility, reduced disease control, or crop injury can result from mixing FURIOUS® with other products.

Before using any tank mix (fungicides, insecticides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated (including plant cultivars) to ensure that a phytotoxic response will not occur as a result of application. Evaluate for crop response 3 to 7 days before making an application to the entire crop.

When an adjuvant is to be used with this product, GROPRO recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Consult a GROPRO representative or local agricultural authorities for more information concerning additives.

MIXING ORDER

1. Water - Begin by agitating a thoroughly clean sprayer tank $\frac{3}{4}$ full of clean water.
2. Agitation - Maintain constant agitation throughout mixing and application.
3. Inductor - If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA bags - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Essential Oil products (such as FURIOUS®).
6. Water-soluble products.
7. Water-soluble additives (such as AMS or UAN when applicable).
8. Make sure each component is thoroughly mixed and suspended before adding tank mix partners. Maintain constant agitation during application.

RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply if rain is expected before the spray is dry for 1 hour. FURIOUS® is rainfast 1 hour after drying.

APPLICATION INSTRUCTIONS

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
TURF GRASS	See Pests Controlled under Product Information section	22 to 33 fluid ounces per acre OR 0.50 - 0.75 fluid ounces per gallon for 1000 sq ft	0

Start applications when conditions are favorable for pest. Continue applications every 10-14 days depending on weather conditions and the development of the pest population.

For control of grubs, billbugs, annual bluegrass weevil, and European crane fly make application prior to egg hatch of the target pest.

For Chinch bugs, make application prior to or during the hatching of the first instar nymphs.

For control of Mole Crickets make application prior to or during peak egg hatch hatch.

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

When pest pressure is high use shorter 7-10 day intervals between sprays.

For backpack applications mix fluid ounces needed for control per 1000 square feet in 1-1.5 gallon of water.

For all applications a use of a surfactant is suggested to improve efficiency.

APPLICATION INSTRUCTIONS (continued)

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
Ornamental Trees Shrubs Evergreens Flowers Foliage Plants Groundcovers Interior plantscapes	See Pests Controlled under Product Information section	22 to 48 fluid ounces per acre OR 0.50 - 1.10 fluid ounces per gallon for 1000 sq ft	0

Start applications when conditions are favorable for pest. Continue applications every 10-14 days depending on weather conditions and the development of the pest population. GROPRO recommends establishing treatments prior to high pest populations as this will increase control.

For control of grub larvae such as Japanese beetle larvae, chafers, *Phyllophaga* spp., Asiatic garden beetle, oriental beetle. **DO NOT** apply in less than 0.75 fluid ounces in 2 gal of water per 1000 square feet. Irrigate to thoroughly incorporate into the top soil profile.

For soil drench uniformly apply a dosage of no less than 1.10 fluid ounces in 10 gal of water per 1000 square feet as a drench to be applied around the base of the tree root zone.

For flowers and groundcovers apply as a broadcast treatment and incorporate into the soil before planting or apply after petal fall or before bloom.

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

APPLICATION INSTRUCTIONS (continued)

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
Pome Fruit Apple Crabapple Loquat Mayhaw Oriental pear Pear (Oriental) Quince	See Pests Controlled under Product Information section	22 to 33 fluid ounces per acre OR 0.50 - 0.75 fluid ounces per gallon for 1000 sq ft	0

For knockdown and residual control of insects in pome fruit, apply FURIOUS® at first sign of infestation.

DO NOT make FURIOUS® applications at intervals longer than 10-14 days.

For maximum effectiveness, apply FURIOUS® at the first sign of insects, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing insect populations.

FURIOUS® can be applied via foliar and soil allowing for total insect control options.

FURIOUS® has a ZERO (0) Pre-Harvest Interval (PHI).

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

APPLICATION INSTRUCTIONS (continued)

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
Avocado (around perimeter of residential or commercial areas)	See Pests Controlled under Product Information section	22 to 33 fluid ounces per acre OR 0.50 - 0.75 fluid ounces per gallon for 1000 sq ft	0

For knockdown and residual control of insects in Avocado, apply FURIOUS® at first sign of infestation.

DO NOT make FURIOUS® applications at intervals longer than 10-14 days.

For maximum effectiveness, apply FURIOUS® at the first sign of insects, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing insect populations.

FURIOUS® can be applied via foliar and soil allowing for total insect control options.

FURIOUS® has a ZERO (0) Pre-Harvest Interval (PHI).

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

APPLICATION INSTRUCTIONS (continued)

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
Vines (around perimeter of residential or commercial areas)	See Pests Controlled under Product Information section	22 to 48 fluid ounces per acre OR 0.50 - 1.10 fluid ounces per gallon for 1000 sq ft	0

For knockdown and residual control of insects in vines, apply FURIOUS® at first sign of infestation.

DO NOT make FURIOUS® applications at intervals longer than 10-14 days.

For maximum effectiveness, apply FURIOUS® at the first sign of insects, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing insect populations.

FURIOUS® can be applied via foliar and soil allowing for total insect control options.

FURIOUS® has a ZERO (0) Pre-Harvest Interval (PHI).

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

APPLY IN MORNING OR EVENING, NEVER APPLY DURING THE HOTTEST PART OF THE DAY.

APPLICATION INSTRUCTIONS (continued)

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
Tree Nuts Group Almonds Beech nut Brazil nut Butternut Cashew Chestnut Chinquapin Filbert Hickory nut Macadamia nut Pecan Pistachio Walnuts (black and English) (around perimeter of residential or commercial areas)	See Pests Controlled under Product Information section	22 to 48 fluid ounces per acre OR 0.50 - 1.10 fluid ounces per gallon for 1000 sq ft	0

For knockdown and residual control of insects in nuts, apply FURIOUS® at first sign of infestation.

DO NOT make FURIOUS® applications at intervals longer than 10-14 days.

For maximum effectiveness, apply FURIOUS® at the first sign of insects, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing insect populations.

FURIOUS® can be applied via foliar and soil allowing for total insect control options. FURIOUS® has a ZERO (0) Pre-Harvest Interval (PHI).

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

APPLICATION INSTRUCTIONS (continued)

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
Citrus and Citrus Hybrids Orange (sweet and sour) Calamondin Grapefruit Kumquat Lemon Lime Pummelo Tangerine Tangelo (around perimeter of residential or commercial areas)	See Pests Controlled under Product Information section	52 to 48 fluid ounces per acre OR 0.60 - 1.10 fluid ounces per gallon for 1000 sq ft	0

For knockdown and residual control of insects in citrus fruit, apply FURIOUS® at first sign of infestation.

DO NOT make FURIOUS® applications at intervals longer than 10-14 days.

For maximum effectiveness, apply FURIOUS® at the first sign of insects, and before the population increases. Application should be timed to coincide with locally recommended treatment threshold levels in developing insect populations.

FURIOUS® can be applied via foliar and soil allowing for total insect control options. FURIOUS® has a ZERO (0) Pre-Harvest Interval (PHI).

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

APPLICATION INSTRUCTIONS (continued)

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
Grass areas of fields Forest Nurseries Sod	See Pests Controlled under Product Information section	25 to 48 fluid ounces per acre OR 0.60 - 1.10 fluid ounces per gallon for 1000 sq ft	0

Start applications when conditions are favorable for pest. Continue applications every 10-21 days depending on weather conditions and the development of the pest population. GROPRO recommends establishing treatments prior to high pest populations as this will increase control.

For control of Grub Larvae such as Japanese beetle larvae, chafers, beetle, Asiatic garden beetle, oriental beetle. **DO NOT** apply in less than 1.10 fluid ounces in 2 gal of water per 1000 square feet. Irrigate to thoroughly incorporate into the top soil profile.

For soil drench uniformly apply a dosage of no less than 1.10 fluid ounce in 10 gal of water per 1000 square feet as a drench to be applied around the base of the tree root zone.

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

APPLICATION INSTRUCTIONS (continued)

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
Nursery and Greenhouse Fabaceae family Brassicaceae family Liliaceae family Lamiaceae family Cucurbitaceae family Solanaceae family	See Pests Controlled under Product Information section	22 to 48 fluid ounces per acre OR 0.50 - 1.10 fluid ounces per gallon for 1000 sq ft	0

Start applications when conditions are favorable for pest. Continue applications every 10-14 days depending on weather conditions and the development of the pest population. GROPRO recommends establishing treatments prior to high pest populations as this will increase control.

For control of soil pests apply a soil drench uniformly using no less than 1.1 fluid ounces in 10 gal of water per 1000 square feet as a drench to be applied in pots or containers to saturate root zone.

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

APPLICATION INSTRUCTIONS (continued)

CROP	PEST	RATE	MINIMUM TIME FROM APPLICATION TO ENTRY
Cannabis	See Pests Controlled under Product Information section	22 to 32 fluid ounces per acre OR 0.50 - 0.75 fluid ounces per gallon for 1000 sq ft	0

Start applications when conditions are favorable for pest. Continue applications every 10-14 days depending on weather conditions and the development of the pest population. GROPRO recommends establishing treatments prior to high pest populations as this will increase control.

For control of soil pests apply a soil drench uniformly using no less than 0.75 fluid ounces in 5 gal of water per 1000 square feet as a drench to be applied in pots or containers to saturate root zone.

FOR ALL APPLICATIONS THE USE OF A SURFACTANT IS RECOMMENDED.

CONDITIONS OF SALE AND WARRANTY

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of GROPRO or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

GROPRO warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GROPRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND GROPRO EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GROPRO AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

GROPRO and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of GROPRO.


For control of insects via soil or foliar in turfgrass (including sod farms), landscape ornamentals, on ornamental and fruit and nut trees, on ornamental and vegetable plants in greenhouses, nurseries, commercial, residential, and interior plantscapes.


**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle (If you do not understand this label, find someone to explain it to you in detail)

See inside for complete First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty, and state-specific crop and/or use site restrictions.

FURIOUS® is a registered trademark of GROPRO





FURIOUS

I N S E C T I C I D E

ACTIVE INGREDIENT:

Geraniol Oil	30.00%
Peppermint oil	1.00%
Cottonseed oil	0.10%
Rosemary oil	0.01%
OTHER INGREDIENTS:*	68.89%
TOTAL:	100.00%

*Canola Oil, Water

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

"SEE LABEL BOOKLET FOR PRECAUTIONARY
STATEMENTS"

**How can we help?
1-833-476-7761**

MANUFACTURED BY:

GroPro^{US}

12600 West Frontage Road,
Burnsville MN 55337
Phone: 1-833-476-7761
www.groproag.com

ENVIRONMENTALLY SAFE
When Used as Directed

Intended for use by commercial applicators. For use in greenhouses, ornamentals, cannabis and on turf related to golf courses, sod farms, sport fields, residential, institutional, municipal, commercial, and other turf grass areas. This product has not been registered by the United States Environmental Protection Agency. GroPro represents that this product qualifies for exemption from registration under FIFRA.

APPROVED FOR USE ON LOW-THC CANNABIS OR MEDICAL MARIJUANA.

NET CONTENTS: ■ 64 fl oz (1.8925 L)



TURF AND ORNAMENTAL

GroPro^{US}