

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms. For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

Lambda Cyhalothrin

GROUP

3

INSECTICIDE

Grizzly Too CS

Insecticide

ACTIVE INGREDIENT:

Lambda-cyhalothrin: 22.88%

Other Ingredients: 77.12%

Total: 100.0%

This product contains 2.08 pounds of active ingredient per gallon and is a capsule suspension.
Contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye.
- Call a poison control center or doctor for treatment advice.

If swallowed:

- Immediately call a poison control center or doctor.
- **DO NOT** induce vomiting unless told to do so by a poison control center.
- **DO NOT** give any liquid to the person.
- **DO NOT** give anything by mouth to an unconscious person.

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For 24 hour Medical or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), call CHEMTREC toll free at 1-800-424-9300.

NOTE TO PHYSICIAN

Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

EPA Reg. No. 42750-385-1381

EPA Est. No. _____

Net Contents: _____ gallons

MANUFACTURED FOR
Winfield Solution, LLC
PO BOX 64589
St. Paul MN 55164

|

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION/PRECAUCIÓN

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber, \geq 14 mils polyvinyl chloride \geq 14 mils or viton \geq 14 mils.
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFS 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife.

For terrestrial uses: **DO NOT** apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater.

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds if bees or other pollinating insects are foraging the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT mix or allow contact with oxidizing agents or reducing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Chemical resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber, \geq 14 mils polyvinyl chloride \geq 14 mils or viton \geq 14 mils.
- Protective eyewear

DO NOT apply this product in a way that it will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult State/Tribal agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

INFORMATION ON USE OF THIS PRODUCT

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or aerial equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air or 10 gallons per acre by ground unless otherwise specified in this label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher listed application volumes and/or higher listed use rates may improve initial and residual control.

For cutworm control, this product may be applied before, during, or after planting. For soil-incorporated applications, use higher listed rates for improved control.

RESISTANCE MANAGEMENT

This product is a Group 3 Insecticide (contains the active ingredient lambda-cyhalothrin). Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control, and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

APPLICATION INSTRUCTIONS

SPRAY DRIFT PRECAUTIONS BUFFER ZONES

Vegetative Buffer Strip

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing this product onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers:

Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21 pp.

www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf

In the State of New York, a 25 ft. vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for both aerial or ground application. For aerial applications, the 25 ft. vegetated non-cropped buffer strip for runoff protection would be part of the larger 150 ft. buffer strip (or 450 ft. buffer strip for ULV application) required for spray drift.

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast)

DO NOT apply within 25 feet of aquatic habitats (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

Buffer Zone for ULV Aerial Application

DO NOT apply within 450 feet of aquatic habitats (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

Buffer Zone for Non-ULV Aerial Application

DO NOT apply within 150 feet of aquatic habitats (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

SPRAY DRIFT REQUIREMENTS

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition. **DO NOT** apply when the wind velocity exceeds 15 mph.

Temperature Inversion

DO NOT make aerial or ground applications into temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices.

The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. **DO NOT** release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downward. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

TANK MIX APPLICATION

When tank mixing with any other agricultural products, this product may be added last to avoid nozzle plugging. Fill the tank with 1/2 - 2/3 volume of the mixing diluent. All other products should be fully dispersed in the mixing diluent before adding the recommended rate of this product to the tank. Add the remainder of the mixing diluent volume. It is recommended that mixing and spray equipment have continuous agitation for best results. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

While this product has good flexibility for tank mixing with other agricultural products, a jar test for physical compatibility is recommended for untried mixtures, using proper ratios and mixing sequences of all ingredients to be included in the mixture.

This product is an aqueous based formulation. **DO NOT** use non-emulsifiable oils in combination with this product. If adjuvants are used, use only:

- Nonionic Surfactant (NIS) containing at least 75% surface agent, or
- Nonphytotoxic Crop Oil Concentrate (COC), including once-refined Vegetable Oil Concentrate (VOC), or,
- Methylated Sunflower Oils (MSO) containing a minimum of 17% emulsifier.

Adjuvants other than NIS or COC may be used providing the product meets the following criteria:

- Contains only EPA exempt ingredients.
- Is nonphytotoxic to the target crop.
- Is compatible in mixture. (May be established through a jar test.)
- Is supported locally for use with this product on the target crop through proven field trials and through university and extension recommendations.

In addition, the following may be used as diluents:

- Crop Oil Concentrate
- Methylated Sunflower Oils
- Urea-Ammonium Nitrate

DO NOT use the following in combination with this product as diluents or adjuvants:

- Non-emulsifiable oils
- Diesel Fuel
- Straight Mineral Oil

CHEMIGATION

Sprinkler Irrigation Application

Apply this product at rates and timing described elsewhere in this label. As local recommendations differ, consult your local State Extension Service or other local experts for recommendations on adjuvant or diluent types, tank mixes, and mixing instructions. These recommendations should be proven, through university and extension field trials, to be effective with this product applied by chemigation.

Check the irrigation system to insure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period.

Apply by injecting the labeled rate of this product into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1-0.2 acre-inch of water. In general, use the least amount of water required for proper distribution and coverage. It is recommended that the product be injected into the main irrigation line ahead of a right angle turn in the line to ensure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

In addition to the above instructions, if application is being made during a normal irrigation set of a stationary sprinkler, the labeled rate of this product for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

Use Precautions – Sprinkler Irrigation Applications

1. Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move. **DO NOT** apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
3. If you have any questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers, or other experts.
4. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label prescribed safety devices for public water systems are in place.
5. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
6. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and are capable of being fitted with a system interlock.
12. Any alternatives to the above required safety devices must conform to the list of EPA-approved alternative devices.

13. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment or non-uniform distribution of treated water.

CROP SPECIFIC USE INSTRUCTIONS

[Note to reviewer: The following statement may be added to Crop Specific Use Directions as per CDPR requirement]

[Not approved for use in California]

ALFALFA, ALFALFA GROWN FOR SEED

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|---|---|
| Alfalfa Caterpillar Army Cutworm Cutworm spp. Green Cloverworm Leafhopper spp. Looper spp. Three-cornered Alfalfa Hopper Velvetbean Caterpillar Webworm spp. | 0.92-1.54 fl. oz. (0.015-0.025 lb.a.i) | Use scouting to determine need for applications. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of foliage. |
| Alfalfa Seed Chalcid (Adult) Alfalfa Weevil Armyworm Bean Leaf Beetle (Adult) Blister Beetle spp. Blue Alfalfa Aphid Clover Leaf Weevil spp. Clover Root Borer (Adult) Clover Root Curculio spp. (Adult) Clover Stem Borer (Adult) Corn Earworm Cowpea Aphid Cowpea Curculio (Adult) Cowpea Weevil (Adult) Cucumber Beetle spp. (Adult) Egyptian Alfalfa Weevil Fall Armyworm ¹ Grape Colaspis (Adult) Grasshopper spp. Green June Beetle (Adult) Green Peach Aphid ³ Japanese Beetle (Adult) Meadow Spittlebug Mexican Bean Beetle Pea Aphid Pea Weevil (Adult) Plant Bug spp. Including Lygus spp. ³ Spotted Alfalfa Aphid Stink Bug spp. Thrips spp. ⁴ Sweet Clover Weevil (Adult) Western Yellowstriped Armyworm | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | Apply in at least 2 gals./A by air or 10 gals./A by ground. In situations of dense foliage and/or high pest populations, use 5-10 gals./A by air or 20 gals./A by ground and higher listed use rates. Also use higher listed rates for improved residual control. Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2-3 days following application. Avoid direct application to bee shelters. ¹ Use higher listed rates for large larvae. ² Suppression only. ³ See resistance statement under Product Information. ⁴ Does not include Western Flower Thrips. |

| | | |
|--|---------------------------------|--|
| Whitefringed Beetle spp. (Adult) Yellowstriped Armyworm | | DO NOT apply more than 0.03 lb. a.i. (0.12 pt. or 1.84 fl. oz. of product)/A per cutting. |
| Beet Armyworm ^{1,3} Blotch Leafminer ³ Spider Mites ² | 1.84 fl. oz. (0.03 lb. a.i.) | DO NOT apply more than 0.12 lb. a.i. (0.46 pt. or 7.37 fl. oz. of product)/A per season. DO NOT apply within 1 day of harvest for forage or within 7 days of harvest for hay. |

CANOLA

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|--|--|--|
| Armyworm spp. Cabbage Seedpod Weevil Cutworm spp. Diamondback Moth Flea Beetle Grasshoppers Looper spp. Lygus Bug | 0.92-1.84 fl. oz. (0.015-0.03 lb.a.i) | Use scouting to determine need for applications, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. Apply by air or ground with enough water to obtain full coverage of foliage. |
| Cabbage Aphid | 1.84 fl. oz. (0.03 lb. a.i.) | For air applications, apply a minimum of 2 gals. of water/A. DO NOT apply more than 0.09 lb. a.i. (0.35 pt. or 5.53 fl. oz. of product)/A per year. DO NOT apply within 7 days of harvest. |

CEREAL GRAINS – CORN (AT PLANT): FIELD CORN, POPCORN, SEED CORN, SWEET CORN

| Pests | | Product Rate Per Acre | | USE DIRECTIONS | | |
|---|--------|--|--------|--|--------|--------|
| Corn Rootworm Larvae (Western, Northern, Southern, Mexican) Cutworm spp. Lesser Cornstalk Borer Red Imported Fire Ant ¹ Seedcorn Beetle Seedcorn Maggot Wireworm spp. ¹ White Grub spp. | | 0.31 fl. oz. (0.005 lb.a.i) | | DO NOT harvest or graze livestock or cut treated crops for feed within 21 days of at plant application. DO NOT apply more than 0.09 lb. a.i. (0.35 pt. or 5.53 fl. oz. of product)/A per crop at plant. For field corn, popcorn, and seed corn DO NOT apply more than 0.12 lb. a.i. (0.46 pt. or 7.37 fl. oz. of product)/A per crop from at plant and foliar applications. For sweet corn DO NOT apply more than 0.48 lb. a.i. (1.84 pts. or 29.49 fl. oz. of product)/A per crop from at plant and foliar applications. For Banded Applications – Make application at planting as a 5-7 inch T-band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel. For In-Furrow Applications – Make application into the seed furrow through spray nozzles or microtubes, behind the planter furrow openers and in front of the press wheel. Apply a minimum of 3 gals. finished spray/A. ¹ Suppression only. | | |
| | | 0.31 fl. oz. (0.005 lb.a.i) / 1000 ft of row ² | | | | |
| ² lbs. a.i. and fl. oz./A of this product applied at 0.31fl. oz./1,000 ft. of row for various row spacings. | | | | | | |
| Row Spacing | 40" | 38" | 36" | 34" | 32" | 30" |
| Linear ft./A | 13,068 | 13,756 | 14,520 | 15,374 | 16,335 | 17,424 |
| lbs. a.i./A | 0.067 | 0.07 | 0.075 | 0.079 | 0.084 | 0.09 |
| fl. oz./A | 4.12 | 4.30 | 4.61 | 4.85 | 5.16 | 5.53 |

CEREAL GRAINS – CORN (FOLIAR): FIELD CORN, POPCORN, SEED CORN, SWEET CORN

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|--|---|
| Corn Earworm ¹ Cutworm spp. Green Cloverworm Meadow Spittlebug Western Bean Cutworm ¹ | 0.92-1.54 fl. oz. (0.015-0.025 lb.a.i.) | Use scouting or locally prescribed corn growth stages to determine need for application, usually at intervals of 7 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds or other locally recommended methods. |
| Armyworm ² Bean Leaf Beetle Bird Cherry-Oat Aphid ³ Cereal Leaf Beetle Corn Leaf Aphid ³ Corn Rootworm Beetle (Adult beetles including Mexican, Northern, Southern, Western) English Grain Aphid ³ European Corn Borer ¹ Fall Armyworm ² Flea Beetle spp. Grasshopper spp. Hop vine Borer ¹ Japanese Beetle (Adult) Lesser Cornstalk Borer Sap Beetle (Adult) Seedcorn Beetle Southwestern Corn Borer ¹ Stalk Borer ¹ Stink Bug spp. Tobacco Budworm ^{1,4} Webworm spp. Yellowstriped Armyworm ² | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | <p>Apply by ground or air using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in at least 2 gals. of water/A.</p> <p>For chinch bug control, begin application when bugs migrate from small grains or grass weeds to small corn and direct the spray to the base of corn plants. Make additional applications at 3-5 day intervals if needed. This product may only suppress heavy infestations and/or subsequent migrations.</p> <p>For control of adult corn rootworm beetles (<i>Diabrotica</i> spp.) as part of an aerial applied corn rootworm control program use at least 1.84 fl. oz./A (0.03 lb. a.i./A).</p> |
| Beet Armyworm ⁴ Chinch Bug Green Bug ^{3,4} Mexican Rice Borer ¹ Rice Stalk Borer ¹ Southern Corn Leaf Beetle ³ Sugarcane Borer | 1.84 fl. oz. (0.03 lb. a.i.) | <p>¹For control before the larva bores into the plant stalk or ear.</p> <p>²Use higher listed rates for large larvae.</p> <p>³Suppression only.</p> <p>⁴See resistance statement under Product Information.</p> <p>DO NOT apply within 21 days of harvest.</p> <p>DO NOT allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after treatment. DO NOT feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.</p> |

| | | |
|--|--|--|
| | | <p>DO NOT apply more than 0.12 lb. a.i. (0.46 pt. or 7.37 fl. oz. of product)/A per crop from at plant and foliar applications.</p> <p>DO NOT apply more than 0.06 lb. a.i. (0.23 pt. or 3.69 fl. oz. of product) after silk initiation.</p> <p>DO NOT apply more than 0.03 lb. a.i. (0.12 pt. or 1.84 fl. oz. of product)/A after corn has reached the milk stage (yellow kernels with milky fluid).</p> |
|--|--|--|

CEREAL GRAINS – SWEET CORN (FOLIAR)

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|--|---|---|
| Aphid spp. ^{2,3} Armyworm ¹ Aster Leafhopper Beet Armyworm ^{1,3} Cereal Leaf Beetle Chinch Bug Common Cornstalk Borer Corn Rootworm Beetle (Adult beetles including Mexican, Northern, Southern, Western) Corn Earworm Cutworm spp. European Corn Borer Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Sap Beetle (Adult) Southern Armyworm ¹ Southwestern Corn Borer Spider Mite spp. ² Stink Bug spp. Tarnished Plant Bug Yellowstriped Armyworm ¹ Western Bean Cutworm Webworm spp. | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | ¹ Use higher listed rates for large larvae. ² Suppression only. ³ See resistance statement under Product Information. DO NOT apply within 1 day of harvest. DO NOT allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. DO NOT feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment. DO NOT apply more than 0.48 lb. a.i. (1.84 pts. Or 29.49 fl. oz. of product)/A per crop from at plant and foliar applications. Use scouting or locally prescribed corn growth stages to determine need for application, usually at intervals of 4 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. For best results target control before insects enter the stalk or ear. Apply ground or air using enough water and application methods to obtain full coverage of foliage and ears (if present). When applying by air, apply in at least 2 gals. of water per acre. |
| Corn Silkfly (Adult) ⁴ Green Bug ^{2,3} | 1.84 fl. oz. (0.03 lb. a.i.) | |

| | | |
|--|--|---|
| | | For control of adult corn rootworm beetles (<i>Diabrotica</i> spp.) as part of an aerial applied corn rootworm control program use a minimum of 0.025 lb. a.i. (1.54 fl. oz.)/A. |
|--|--|---|

CEREAL GRAINS – RICE AND WILD RICE

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|--|--|
| Bird Cherry-Oat Aphid Chinch Bug Fall Armyworm Grasshopper spp. Green Bug Leafhopper spp. Rice Stink Bug Rice Water Weevil (Adult) Riceworm Sharpshooter spp. True Armyworm Yellowstriped Armyworm Yellow Sugarcane Aphid | 1.54-2.46 fl. oz. (0.025-0.04 lb. a.i.) | ¹ For control before the larvae bore into the plant stalk. DO NOT release flood water within 7 days of application. DO NOT apply more than 0.12 lb. a.i. (0.46 pt. or 7.37 fl. oz. of product)/A per season. DO NOT apply more than 0.08 lb. a.i. (0.31 pt.)/A within 28 days of harvest or more than 0.04 lb. a.i. (0.15 pt.)/A within 21 days of harvest. DO NOT apply within 21 days of harvest. |
| European Corn Borer ¹ Mexican Rice Borer ¹ Rice Seed Midge ¹ Sugarcane Borer ¹ | 1.84-2.46 fl. oz. (0.03-0.04 lb. a.i.) | DO NOT use treated rice fields for the aquaculture of edible fish and crustacea. DO NOT apply as an ultra-low volume (ULV) spray. |

CEREAL GRAINS – SORGHUM (GRAIN)

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|--|--|---|
| Cutworm spp. Sorghum Midge | 0.92-1.23 fl. oz. (0.015-0.02 lb. a.i.) | ¹ Use higher listed rates for large larvae. ² For control before the larva bores into the plant stalk. |
| Armyworm Beet Armyworm ³ Corn Earworm European Corn Borer ² Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Lesser Cornstalk Borer ² Southwestern Corn Borer ² Stink Bug spp. Webworm spp. Yellowstriped Armyworm ¹ | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | ³ See resistance statement under Product Information. DO NOT apply more than 0.08 lb. a.i. (0.31 pt. or 4.91 fl. oz. of product)/A per season. DO NOT apply more than 0.06 lb. a.i. (0.23 pt. or 3.69 fl. oz. of product)/A per season after crop emergence. DO NOT apply more than 0.02 lb. a.i. (0.08 pt. or 1.23 fl. oz. of product)/A per season once crop is in soft dough stage. |

| | | |
|---|---------------------------------|--|
| | | <p>DO NOT apply within 30 days of harvest.</p> <p>Use scouting to determine need for treatment, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water and application methods to obtain full coverage of target location. When applying by air, apply in at least 2 gals. of water/A.</p> <p>For sorghum midge control, make first application when one quarter of the sorghum heads have emerged and are in tip bloom. If needed, repeat applications at 5-day intervals.</p> |
| Chinch Bug Mexican Rice Borer ² Sugarcane Borer ² | 1.84 fl. oz. (0.03 lb. a.i.) | <p>For chinch bug control, start applications when bugs migrate from small grains or grass weeds to small sorghum, directing spray to the base of sorghum plants. If needed, repeat applications at 3-5 day intervals.</p> <p>This product may only suppress heavy infestations and/or subsequent migrations.</p> |

CEREAL GRAINS – WHEAT, WHEAT HAY, TRITICALE, BARLEY, BUCKWHEAT, OATS AND RYE

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|---|---|
| Army Cutworm Cutworm spp. | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | <p>¹Best control is obtained before insects begin to roll leaves. Once crop has started to boot, this product may provide suppression only. Higher rates and increased coverage will be necessary.</p> <p>²Suppression only.</p> <p>³See resistance statement under Product Information.</p> <p>⁴Make application when adults emerge.</p> <p>DO NOT apply within 30 days of harvest.</p> |
| Armyworm Bird Cherry-Oat Aphid ¹ Cereal Leaf Beetle English Grain Aphid ¹ Fall Armyworm Flea Beetle spp. Grasshopper spp. Hessian Fly ⁴ Orange Blossom Wheat Midge Russian Wheat Aphid ¹ Stink Bug spp. Yellowstriped Armyworm | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | |
| Grass Sawfly | 1.54-1.84 fl. oz. (0.025-0.03 lb. a.i.) | |

| | | |
|---|---------------------------------|--|
| Chinch Bug Corn Leaf Aphid ² Greenbug ^{1,3} Mite spp. ² | 1.84 fl. oz. (0.03 lb. a.i.) | <p>DO NOT apply more than 0.06 lb. a.i. (0.23 pt. or 3.69 fl. oz. of product)/A per season.</p> <p>DO NOT allow livestock to graze in treated areas or harvest treated wheat forage as feed for meat or dairy animals with- in 7 days after treatment. DO NOT feed treated straw to meat or dairy animals within 30 days after last treatment.</p> <p>Use scouting to determine need for treatment, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water and application methods to obtain full coverage of foliage.</p> <p>When applying by air, apply in at least 2 gals. of water/A.</p> <p>For chinch bug control, repeat applications at 3-5 day intervals if needed. This product may only suppress heavy infestations and/or migrations.</p> <p>Because Greenbug is known to have many biotypes, it is possible that this product may only provide suppression. If this occurs, a second application using an alternative chemistry may be needed.</p> |
|---|---------------------------------|--|

COLE CROPS – BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAVALO BROCCOLO, CAULIFLOWER, CHINESE BROCCOLI (GAI LON), CHINESE CABBAGE (NAPA), CHINES MUSTARD CABBAGE (GAI CHOY), KOHLRABI

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|---|--|
| Alfalfa Looper Cabbage Looper Cabbage Webworm Cutworm spp. Imported Cabbageworm Southern Cabbageworm | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | ¹ For control of first and second instars only. ² Suppression only. ³ See resistance statement under Product Information. |
| Aphid spp. ^{2,3} Armyworm Beet Armyworm ^{1,3} | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | |

| | | |
|---|--|---|
| Corn Earworm Diamondback Moth ³ Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Meadow Spittlebug Plant Bug spp. – including Lygus spp. ³ Spider Mite spp. ² Stink Bug spp. Thrips spp. ² Vegetable Weevil (Adult) Whitefly spp. ^{2,3} Yellowstriped Armyworm | | <p>DO NOT apply within 1 day of harvest.</p> <p>DO NOT apply more than 0.24 lb. a.i. (0.92 pts. or 14.74 fl. oz. of product.)/A per season.</p> <p>Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A.</p> |
|---|--|---|

COTTON

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|--|--|
| Cutworm spp. Soybean Thrips Tobacco Thrips | 0.92-1.23 fl. oz. (0.015-0.02 lb. a.i.) | <p>¹For control of first and second instars only.</p> <p>²Suppression only.</p> <p>³See resistance statement under Product Information.</p> |
| Cabbage Looper Cotton Fleahopper Cotton Leafperforator Cotton Leafworm Lygus Bug spp. ³ | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | <p>DO NOT apply within 21 days of harvest.</p> <p>DO NOT graze livestock in treated areas.</p> |
| Bandedwing Whitefly ^{2,3} Beet Armyworm ^{1,3} Boll Weevil Brown Stink Bug Cotton Aphid ^{2,3} Cotton Bollworm European Corn Borer Fall Armyworm Green Stink Bug Southern Green Stink Bug Sweetpotato Whitefly ^{2,3} Tobacco Budworm ³ Twospotted Spider Mite ² | 1.54-2.46 fl. oz. (0.025-0.04 lb. a.i.) | <p>DO NOT apply more than 0.2 lb. a.i. (0.77 pts. or 12.29 fl. oz. of product)/A per season.</p> <p>DO NOT make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season.</p> <p>Use scouting to determine need for application, usually at intervals of 5-7 days. Base the timing and frequency of applications on when insect populations reach local economic thresholds.</p> <p>Apply ground or air using enough water to obtain full coverage of foliage.</p> |

| | | |
|--|--|---|
| | | <p>Applications may also be made with equipment adapted and calibrated for ULV sprays. This product may be mixed with once-refined vegetable oil and applied in a minimum of at least one qt. of finished spray/A.</p> <p>When bollworm/budworm infestation levels are light, 0.02 lb. a.i. (1.23 fl. oz. of product)/A may be applied in conjunction with intense field monitoring.</p> <p>For boll weevil, spray on a 3-5 day schedule.</p> <p>When applied according to the directions above for control of cotton bollworm and tobacco budworm, this product also provides ovicidal control of unhatched <i>Heliothis</i> species eggs.</p> |
|--|--|---|

CURCUBITS – CHAYOTE (FRUIT), CHINESE WAXGOURD, CITRON MELON, CUCUMBER, GHERKIN, GOURD (edible) MOMORDICA spp., MUSKMELON, PUMPKIN, SQUASH (summer and winter), WATERMELON

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|---|--|
| Armyworm spp. ¹ Blister Beetle spp. Cabbage Looper Corn Earworm Cricket spp. Cucumber Beetle spp. (Adult) Cutworm spp. Flea Beetle spp. Grasshopper spp. June Beetle spp. Leaffooted Bug Leafhopper spp. Lygus Bug spp. ¹ Melonworm Pickworm Plant Bug spp. Rindworm species complex Saltmarsh Caterpillar Squash Beetle Squash Bug spp. Squash Vine Borer spp. Stink Bug spp. Thrips spp. ^{1,2} Tobacco Budworm ¹ Webworm spp. | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | <p>Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of solution per acre. When applying by ground, apply in a minimum of 10 gals. of solution per acre.</p> <p>Use higher application volumes and/or listed application rates when foliage is dense, larvae are large, pest populations are high, size of plants increases, or weather conditions are adverse. Use higher listed rates for longer residual. Insects that tunnel or bore into leaves, stems, vines, or fruit must be controlled before penetration. Only insects (larvae and adults) exposed to the product can be controlled with foliar applications of this product.</p> |
| Aphid spp. ¹ Leafminer spp. ^{1,3} Spider Mite spp. ³ | 1.84 fl. oz. (0.03 lb. a.i.) | |

| | | |
|------------------------------|--|--|
| Whitefly spp. ^{1,3} | | <p>¹See resistance statement under Product Information.</p> <p>²Western Flower Thrips are not included.</p> <p>³Suppression only.</p> <p>DO NOT apply within 1 day of harvest.</p> <p>DO NOT apply more than 0.18 lb. a.i. (0.69 pts. or 11.06 fl. oz.)/A per season.</p> |
|------------------------------|--|--|

FRUITING VEGETABLE – TOMATO, TOMATILLO, PEPPERS (BELL AND NON BELL), EGGPLANT, GROUND CHERRY, PEPINO

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|--|--|
| Cabbage Looper Cutworm spp. Hornworm spp. | 0.92-1.23 fl. oz. (0.015-0.02 lb. a.i.) | <p>¹For control of first and second instars only.</p> <p>²Suppression only.</p> |
| Aphid spp. ^{2,3} Beet Armyworm ^{1,3} Blister Beetle spp. Colorado Potato Beetle ³ Cucumber Beetle spp. (Adult) European Corn Borer ⁴ Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Leafminer spp. ² Meadow Spittlebug Pepper Weevil (Adult) ² Plant Bug spp. Southern Armyworm ¹ Spider Mite spp. ² Stalk Borer ⁴ Stink Bug spp. Thrips ⁵ Tobacco Budworm ³ Tomato Fruitworm Tomato Pinworm Tomato Psyllid ^{2,3} Vegetable Weevil (Adult) Whitefly spp. ^{2,3} Yellowstriped Armyworm ¹ | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | <p>³See resistance statement under Product Information.</p> <p>⁴For control before the larva bores into the plant stalk or fruit.</p> <p>⁵Does not include Western Flower thrips. DO NOT apply within 5 days of harvest.</p> <p>DO NOT apply more than 0.36 lb. a.i. (1.38 pts. or 22.12 fl. oz. of product)/A per season.</p> <p>Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A.</p> |

GRASS FOR FORAGE, FODDER AND HAY – PASTURE AND RANGELAND GRASS, GRASS GROWN FOR HAY OR SILAGE AND GRASS GROWN FOR SEED

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|---|---|
| Army Cutworm Cutworm spp. Essex Skipper Range Caterpillar Striped Grass Looper | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | See additional instructions below. ¹ Best control is obtained before insects begin to roll leaves. |
| Beet Armyworm Billbug spp. ³ Bird Cherry-Oat Aphid ¹ Black Grass Bug Black Turfgrass Beetle (Adult) Blue Stem Midge Cereal Leaf Beetle Chinch Bug Crane Fly spp. Cricket spp. English Grain Aphid ¹ Fall Armyworm Flea Beetle spp. Grass Mealybug Grass Sawfly (Adult) Grasshopper spp. Green June Beetle (Adult) Greenbug ^{1,2,4} Japanese Beetle (Adult) Katydid spp. Leafhopper spp. Mite spp. ³ Russian Wheat Aphid ¹ Southern Armyworm Spittlebug spp. Stink Bug spp. Sugarcane Aphid Thrips spp. Tick spp. True Armyworm Webworm spp. Yellowstriped Armyworm | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | ² See resistance statement under Product Information. ³ Suppression only. ⁴ Greenbug is known to have many biotypes. This product may provide suppression only. A second application using alternative chemistry may be needed. Pasture and rangeland grass may be used for grazing or cut for forage 0 days after application. DO NOT cut grass to be dried and harvested for hay until 7 days after the last application. Grass grown for seed: Straw and mature seed (seed screenings) may be used as feed 7 days after the last application. Regrowth of grass grown for seed may be used for grazing, cut for forage or cut to be dried and harvested for hay. DO NOT apply more than 0.03 lb. a.i. (0.12 pt. or 1.84 fl. oz.)/A per cutting for pastures, rangeland, and grasses grown for seed. A minimum re-treatment interval (RTI) of 30 days is required for pastures and rangeland receiving 0.03 lb. a.i./A which have not been cut between applications. DO NOT apply more than 0.09 lb. a.i. (0.35 pt. or 5.53 fl. oz. of product) per acre per season. Use scouting to determine application requirements. Base the timing and frequency of applications on when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 |

| | | |
|--|--|--|
| | | <p>gals. of water per acre. When applying by ground, apply in a minimum of 7 gals. of water per acre.</p> <p>Use higher application volumes and/or listed application rates when foliage is dense, larvae are large, pest populations are high, or weather conditions are adverse. Use higher listed rates for longer residual.</p> <p>For chinch bug control, this product may only suppress heavy infestations and/or migrations. In these situations, a second application using alternative chemistry may be needed.</p> |
|--|--|--|

LEGUME VEGETABLES (BEANS AND PEAS)

EDIBLE PODDED (ONLY): *Canavalia gladiata* – sword bean; *Canavalia ensiformis* – jackbean; *Glycine max* – soybean (immature seed).

EDIBLE PODDED, SUCCULENT SHELLED OR DRIED SHELLED: *Phaseolus* spp. – includes field, kidney, lima, navy, pinto, runner, snap, tepary, and wax beans; *Vigna* spp. – includes adzuki, asparagus, moth, mung, rice, urd and yardlong beans, black-eyed pea, catjang, Chinese longbean, cowpea, Crowder pea, and Southern pea; *Pisum* spp. – includes dwarf, edible-pod, English, field, garden, green, snow, and sugar peas; *Cajanus cajan* – Pigeon pea.

SUCCULENT SHELLED OR DRIED SHELLED: *Vicia faba* – broadbean (fava bean).

DRIED SHELLED (ONLY): *Lupinus* spp. – includes grain, sweet, white and sweet white lupines; *Cicer arietinum* – chickpea (garbanzo bean), *Cyamopsis tetraganloba* – guar, *Lablab purpureus* – lablab bean (hyacinth bean), *Lens esculata* – lentils.

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|--|---|---|
| Cutworm spp. Green Cloverworm Imported Cabbageworm Mexican Bean Beetle Saltmarsh Caterpillar Velvetleaf Caterpillar | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | For edible podded and succulent shelled legume vegetables, DO NOT apply within 7 days of harvest. For dried shelled legume vegetables, DO NOT apply within 21 days of harvest. |
| Alfalfa Caterpillar Aphid spp. ⁴ Armyworm ² Bean Leaf Beetle Bean Leafskeletonizer Blister Beetle spp. Corn Earworm Corn Rootworm Beetle spp. (Adult) Cucumber Beetle spp. (Adult) Curculio and Weevil spp. ¹ (Foliage and pod feeding adults and larvae) European Corn Borer Fall Armyworm ² | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | DO NOT apply more than 0.12 lb. a.i. (0.46 pt. or 7.37 fl. oz. of product)/A per season. For succulent and dried shelled peas and beans, DO NOT graze livestock in treated areas or harvest vines for forage or hay. |

| | | |
|--|---------------------------------|---|
| Flea Beetle spp. (Adult) Flea Hopper spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Leaf-tier spp. Looper spp. Meadow Spittlebug Painted Lady Butterfly (Larva) Plant Bug spp. including Lygus spp. ⁴ Stalk Borer ¹ Stink Bug spp. Threecornered Alfalfa Hopper Thrips spp. ^{4,5} Tobacco Budworm ⁴ Webworm spp. Western Bean Cutworm Western Yellowstriped Armyworm ² Yellowstriped Armyworm ² | | Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A. ¹ For control before the larva bores into the plant stalk or pods. ² Use higher listed rates for large larvae. ³ For suppression only. |
| Beet Armyworm ^{3,4} Leafminer spp. ^{3,4} Lesser Cornstalk Borer ³ Soybean Looper ^{3,4} Spider Mite spp. ³ Whitefly spp. ^{3,4} | 1.84 fl. oz. (0.03 lb. a.i.) | ⁴ See resistance statement under Product Information. ⁵ Does not include Western Flower Thrips. |

LEGUME VEGETABLES (SOYBEANS)

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|--|---|---|
| Bean Leaf Beetle Cabbage Looper Corn Earworm Corn Rootworm Beetle (Adult beetles including Mexican, Northern, Southern, Western) Cutworm spp. Green Cloverworm Mexican Bean Beetle Painted Lady (Thistle) Caterpillar Potato Leafhopper Saltmarsh Caterpillar Soybean Aphid ⁴ Threecornered Alfalfa Hopper Thrips spp. ⁵ Velvetbean Caterpillar | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. DO NOT graze or harvest treated soybean forage, straw or hay for livestock feed. Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A. |

| | | |
|---|--|--|
| Armyworm ¹ Blister Beetle spp. European Corn Borer Fall Armyworm ¹ Grasshopper spp. Japanese Beetle (Adult) Plant Bug spp. Silverspotted Skipper Stink Bug spp. Tobacco Budworm ³ Webworm spp. Yellowstriped Armyworm ¹ | 1.54-1.84 fl. oz. (0.025-0.03 lb. a.i.) | For control of adult corn rootworm beetles (<i>Diabrotica</i> spp.) as part of an aerial applied to corn rootworm control program use at least 1.23 fl. oz./A of product (0.02 lb. a.i./A). ¹ Use higher listed rates for large larvae. ² Suppression only. ³ See resistance statement under Product Information. |
| Beet Armyworm ^{2,3} Lesser Cornstalk Borer ² Soybean Looper ^{2,3} Spider Mite spp. ² | 1.84 fl. oz. (0.03 lb. a.i.) | ⁴ Use lower rates for early season applications and/or lighter populations. ⁵ Does not include Western Flower Thrips. DO NOT apply within 30 days of harvest. DO NOT apply more than 0.06 lb. a.i. (0.23 pt. or 3.69 fl. oz of product)/A per season. |

LETTUCE (HEAD AND LEAF)

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|--|---|--|
| Alfalfa Looper Cabbage Looper Cutworm spp. Green Cloverworm Imported Cabbageworm Saltmarsh Caterpillar | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. |
| Aphid spp. ^{2,3} Armyworm Beet Armyworm ^{1,3} Corn Earworm Diamondback Moth ³ European Corn Borer Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Meadow Spittlebug Plant Bug spp. including Lygus spp. ³ Southern Armyworm Spider Mite spp. ² Stink Bug spp. Tobacco Budworm ³ Vegetable Weevil (Adult) Whitefly spp. ^{2,3} | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A. ¹ For control of first and second instars only. ² Suppression only. ³ See resistance statement under Product Information. DO NOT apply within 1 day of harvest. |

| | | |
|--|--|--|
| | | DO NOT apply more than 0.3 lb. a.i. (1.15 pts. or 18.43 fl. oz. of product)/A per season. |
|--|--|--|

ONION (BULD) AND GARLIC

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|---|---|
| Cutworm spp. Leafminer spp. (Adult) Onion Maggot (Adult) Seedcorn Maggot (Adult) | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | See additional instructions below. Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. |
| Aphid spp. ² Armyworm spp. ¹ Flower Thrips ^{2,3} Onion Thrips ³ Plant Bug spp. Stink Bug spp. Tobacco Thrips ³ Western Flower Thrips ^{2,3} | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | <p>Use the higher listed label rates as thrips population increases and avoid rescue situations.</p> <p>Apply by ground or air using enough water and application methods to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A.</p> <p>To control thrips by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v or a silicone adjuvant (follow manufacturer's use directions) may improve the deposition of the spray and increase plant coverage.</p> <p>Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds.</p> <p>Use the higher listed label rates as thrips population increases and avoid rescue situations.</p> <p>Apply by ground or air using enough water and application methods to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A.</p> <p>To control thrips by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v or a silicone adjuvant (follow manufacturer's use directions) may</p> |

| | | |
|--|--|---|
| | | <p>improve the deposition of the spray and increase plant coverage.</p> <p>¹For control of first and second instars only.</p> <p>²Suppression only.</p> <p>³See resistance statement under Product Information.</p> <p>DO NOT apply within 14 days of harvest.</p> <p>DO NOT apply more than 0.24 lb. a.i. (0.92 pts. or 14.74 fl. oz. of product)/A per season.</p> |
|--|--|---|

PEANUTS

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|--|---|--|
| Cutworm spp. Green Cloverworm Potato Leafhopper Rednecked Peanut Worm Threecornered Alfalfa Leafhopper Velvetbean Caterpillar | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | Use scouting to determine need for application, usually at intervals of 7 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. |
| Bean Leaf Beetle Corn Earworm Fall Armyworm ¹ Grasshopper spp. Southern Corn Rootworm (Adult) Stink Bug spp. Tobacco Thrips Vegetable Weevil Whitefringed Beetle (Adult) | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | <p>Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A.</p> <p>¹Use higher listed rates for large larvae.</p> <p>²Suppression only.</p> <p>³See resistance statement under Product Information</p> |
| Aphid spp. ² Beet Armyworm ^{2,3} Lesser Cornstalk Borer ² Soybean Looper ^{2,3} Spider Mite spp. ² | 1.84 fl. oz. (0.03 lb. a.i.) | <p>DO NOT apply within 14 days of harvest.</p> <p>DO NOT apply more than 0.12 lb. a.i. (0.46 pt. or 7.37 fl. oz. of product)/A per season.</p> |

POME FRUITS – APPLE, CRABAPPLE, LOQUAT, MAYHAW, ORIENTAL PEAR, PEAR, QUINCE

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|---|---|
| Apple Aphid Apple Maggot (Adult) Cherry Fruit Fly spp. (Adult) Codling Moth Green Fruitworm Japanese Beetle Leafhopper spp. Leafroller spp. Lesser Appleworm Omnivorous Leafroller Orange Tortrix Oriental Fruit Moth Pear Psylla ¹ Pear Sawfly Periodical Cicada Plant Bug spp. Plum Curculio Rosy Apple Aphid San Jose Scale (Fruit infestations only) Spirea Aphid ¹ Stink Bug spp. Tent Caterpillar spp. Tentiform Leaf Miner spp. Tree Borer spp. Tufted Apple Budworm Webworm spp. | 1.23-2.46 fl. oz. (0.02-0.04 lb. a.i.) | <p>See additional instructions below. Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds and IPM recommendations.</p> <p>Apply by ground or air using enough water to obtain full coverage of the foliage or target area. When applying by air, apply in at least 5 gals. of water per acre, but use higher volumes as appropriate for thorough coverage.</p> <p>¹Suppression only.</p> <p>DO NOT apply within 21 days of harvest.</p> <p>DO NOT apply more than 0.2 lb. a.i. (0.77 pts. or 12.29 fl. oz. of product)/A per year.</p> <p>DO NOT apply more than 0.16 lb. a.i. (0.61 pts. or 9.83 fl. oz. of product)/A per year post bloom.</p> |

STONE FRUITS – APRICOT, SWEET CHERRY, TART CHERRY, NECTARINE, PEACH, PLUM, CHICASAW PLUM, DAMSON PLUM, JAPANESE PLUM, PLUMCOT, PRUNE

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|---|---|
| American Plum Borer Apple Maggot (Adult) Black Cherry Aphid Cherry Fruit Fly spp. (Adult) Codling Moth Green Fruitworm Japanese Beetle June Beetle Leafhopper spp. Leafroller spp. Oriental Fruit Moth Peachtree Borer spp. Peach Twig Borer Pear Sawfly Periodical Cicada Plant Bug spp. Plum Curculio | 1.23-2.46 fl. oz. (0.02-0.04 lb. a.i.) | <p>Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds and IPM recommendations.</p> <p>Apply by ground or air using enough water to obtain full coverage of the foliage or target area. When applying by air, apply at least 5 gals. of water per acre but use higher volumes as appropriate for thorough coverage. DO NOT apply within 14 days of harvest.</p> <p>DO NOT apply more than 0.2 lb. a.i. (0.77 pts. or 12.29 fl. oz. of product)/A</p> |

| | | |
|---|--|---|
| Rose Chafer Stink Bug spp. Tent Caterpillar spp. Thrips spp. | | per year. DO NOT apply more than 0.16 lb. a.i. (0.61 pts. or 9.83 fl. oz. of product)/A per year post bloom. |
|---|--|---|

SUGARCANE

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|--|---|
| Mexican Rice Borer ¹ Pygmy Mole Cricket Rice Stalk Borer ¹ Sugarcane Aphid ³ Sugarcane Beetle (Adult) ² Sugarcane Borer ¹ West Indian Cranefly Yellow Sugarcane Aphid ³ | 1.54-2.46 fl. oz. (0.025-0.04 lb. a.i.) | <p>Use scouting to determine need for application, usually at intervals of 7 or more days. Base the timing and frequency of applications on when insect populations reach local economic threshold.</p> <p>Apply by ground or air using enough water to obtain full coverage of the foliage or target area. When applying by air, apply at least 2 gals. of water/A.</p> <p>¹For control before the larva bores into the plant stalk.</p> <p>²Suppression only of beetles active above ground.</p> <p>³See resistance statement under Product Information.</p> <p>DO NOT apply within 21 days of harvest.</p> <p>DO NOT apply more than 0.16 lb. a.i. (0.61 pts. or 9.83 fl. oz. of product)/A per season.</p> |

SUNFLOWER

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|-------------------------------------|---|--|
| Cutworm spp. Sunflower Beetle | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | Use scouting to determine need for application, usually at intervals of 5 or |

| | | |
|--|---|--|
| Banded Sunflower Moth Fall Armyworm ¹ Grasshopper spp. Head-Clipper Weevil (Adult) Japanese Beetle (Adult) Leafhopper spp. Meadow Spittlebug Painted Lady (Thistle) Caterpillar Seed Weevil (Adult) Spotted Cabbage Looper Stem Weevil (Adult) Stink Bug spp. Sunflower Maggot (Adult) Sunflower Moth Woolybear Caterpillar | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of sunflower heads and/or foliage. When applying by air, apply in at least 2 gals. of water/A. ¹ Use higher listed rates for large larvae. ² Suppression only. |
| Beet Armyworm ^{2,3} Spider Mite spp. ² | 1.84 fl. oz. (0.03 lb. a.i.) | ³ See resistance statement under Product Information. DO NOT apply within 45 days of harvest. DO NOT apply more than 0.12 lb. a.i. (0.46 pt. or 7.37 fl. oz. of product)/A per season. DO NOT apply more than 0.09 lb. a.i. (0.35 pt. or 5.53 fl. oz. of product)/A per season after bloom initiation. DO NOT apply as an Ultra-Low Volume (ULV) spray. |

TOBACCO

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|--|--|
| Armyworm spp. ¹ Blister Beetle spp. Cabbage Looper Corn Earworm Cucumber Beetle spp. (Adult) Cutworm spp. Grasshopper spp. Japanese Beetle (Adult) Katydid spp. Plant Bug spp. ³ Potato Tuberworm Saltmarsh Caterpillar Stinkbug spp. Tobacco Aphid spp. ^{2,3} Tobacco Budworm ³ Tobacco Flea Beetle (Adult) Tobacco Hornworm Tobacco Thrips spp. ² | 0.92-1.84 fl. oz. (0.015-0.03 lb. a.i.) | Use scouting to determine need for application, usually at intervals of 7 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of the foliage. When applying by air, apply in at least 2 gals. of water/A. ¹ For control of first and second instars only. ² Suppression only. |

| | | |
|--|--|---|
| Tomato Hornworm Tree Cricket spp. Vegetable Weevil (Adult) Webworm spp. | | ³ See resistance statement under Product Information. DO NOT apply within 40 days of harvest. DO NOT apply more than 0.09 lb. a.i. (0.35 pt. or 5.53 fl. oz. of product)/A per year. |
|--|--|---|

TREE NUTS – ALMOND, BEECH NUT, BRAZIL NUT, BUTTERNUT, CASHEW, CHESTNUT, CHINQUAPIN, FILBER (HAZLENUT), HICKORY NUT, MACADAMIA NUT (BUSH NUT), PISTACHIO, WALNUT-BLACK, WALNUT-ENGLISH (PERSIAN), PECAN

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|--|---|--|
| Ants Chinch Bug Codling Moth Filbertworm Hickory Shuckworm Leaffooted Bug Leafroller spp. Navel Orangeworm Peach Twig Borer Pecan Aphid spp. Pecan Casebearer spp. Pecan Phylloxera spp. Pecan Spittlebug Pecan Weevil Plant Bug spp. Stink Bug spp. Walnut Aphid Walnut Husk Fly spp. (Adult) | 1.23-2.46 fl. oz. (0.02-0.04 lb. a.i.) | Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of the foliage or target area. When applying by air, apply in at least 5 gals. of water per acre, but use higher listed rates as appropriate for thorough coverage. DO NOT apply within 14 days of harvest. DO NOT apply more than 0.16 lb. a.i. (0.61 pts. or 9.83 fl. oz. of product)/A per year. DO NOT apply more than 0.12 lb. a.i. (0.48 pt. or 7.37 fl. oz. of product)/A per year post bloom. |

TUBEROUS AND CORM VEGETABLES – ARRACACHA, ARROWROOT, ARTICHOKE (Chinese and Jerusalem only), CANNA (edible), CASSAVA (bitter and sweet), CHAYOUT (root), CHUFA, DASHEEN, GINGER, LEREN, POTATO, SWEET POTATO, TANIER, TURMERIC, YAM (bean and true)

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|---|---|--|
| Cutworm spp. Leafhopper spp. Saltmarsh Caterpillar Sweet Potato Hornworm Woollybear Caterpillarspp. | 0.92-1.54 fl. oz. (0.015-0.025 lb. a.i.) | Use scouting to determine need for application, usually at intervals of 7 or more days. Base the timing and frequency of applications on when insect |

| | | |
|---|---|---|
| Aphid spp. ¹ Armyworm spp. ¹ Blister Beetle spp. Colorado Potato Beetle ¹ Corn Earworm Cricket spp. Cucumber Beetle spp. (Adult) European Corn Borer Flea Beetle spp. (Adult) Grasshopper spp. Looper spp. ¹ Lygus Bug spp. ¹ Plant Bug spp. Potato Psyllid Potato Tuberworm Stink Bug spp. Sweet Potato Leaf Beetle (Adult) Sweet Potato Vine Borer Thrips spp. ^{1,2} Tortoise Beetle spp. Webworm spp. Weevil spp. (Adult) | 1.23-1.84 fl. oz. (0.02-0.03 lb. a.i.) | populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water per acre. When applying by ground, apply in a minimum of 10 gals. of water per acre. Use higher application volumes and/or application rates when foliage is dense, larvae are large, pest populations are high, plant size increases, or weather conditions are adverse. Use higher listed rates for longer residual. Insects that tunnel or bore into leaves, vines, stems, tubers, or corms must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of this product. |
| Leafminer spp. ^{1,3} Whitefly spp. ^{1,3} Spider Mite spp. ³ | 1.84 fl. oz. (0.03 lb. a.i.) | ¹ See resistance statement under Product Information. ² Does not include Western Flower Thrips. ³ Suppression only. DO NOT apply within 7 days of harvest. DO NOT apply more than 0.12 lb. a.i. (0.46 pt. or 7.37 fl. oz. of product) per acre per season. |

CONIFER AND DECIDUOUS TREES – PLANTATIONS AND NURSERIES

| Pests | Product Rate Per Acre | USE DIRECTIONS |
|--|---|--|
| Bagworm Balsam Twig Aphid Balsam Wooly Aphid Birch Leafminer Black Pine Weevil Elm Leaf Beetle European Elm Leaf Beetle Gypsy Moth Japanese Beetle June Beetle spp. Leaf Beetle spp. Leafroller spp. May Beetle spp. Mealybug spp. ¹ Pales Weevil Pine Chafer | 1.23-2.46 fl. oz. (0.02-0.04 lb. a.i.) | Use scouting to determine timing for control of exposed foliage, flower, cone, seed and bark feeding insects. Base the timing and frequency of applications on when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of target site. When applying by air, apply in at least 2 gals. of water/A. ¹ Suppression only. |

| | | |
|--|--|---|
| Pine Colaspis Beetle Pine Conelet Bug Pine Leaf Chermid Pine Needle Scale Pine Sawfly spp. Pine Tip Moth spp. Pine Tortoise Scale Pine Weevil spp. Poplar Aphid spp. Sawfly spp. Spittlebug spp. Spruce Budworm Tent Caterpillar spp. Tussock Moth spp. Webworm spp. | | DO NOT apply more than 0.24 lb. a.i. (0.92 pts. or 14.74 fl. oz. of product)/A per year. |
|--|--|---|

CONIFER AND DECIDUOUS TREES – SEED ORCHARDS

| Pests | USE DIRECTIONS |
|---|---|
| Coneworm spp. Seed Bug spp. Thrips spp. | <p>For high volume sprayers, dilute 2.46 fl. oz. of product per 100 gals. of water and apply 5-10 gals. of finished spray per tree.</p> <p>For low volume sprayers, dilute 10 fl. oz. of product per 100 gals. of water and apply 100 gals. of finished spray per acre.</p> <p>For aerial applications, apply 7.5 fl. oz. of product per acre in a minimum of 10 gals. of finished spray per acre.</p> <p>DO NOT apply more than 0.5 lb. a.i. (2 pts. or 30.72 fl. oz. of product)/A per year.</p> |

NON-AGRICULTURAL USES

NON-CROPLAND ADJACENT TO AGRICULTURAL AREAS (EXCLUDING PUBLIC LAND)

| Pests | USE DIRECTIONS |
|---|--|
| See crop instructions in sections above for specific pest and rate information. | <p>Spray non-cropland adjacent to agricultural areas to control insects which may migrate to and threaten crops. Follow the Directions for Use instructions, application rates, and spray recommendations found elsewhere on this label for the adjacent crop outlet and target pests.</p> <p>When foliage is dense/large, insect populations are high or larval stages are large, use the highest labeled rate for that crop-pest combination.</p> <p>Repeat as necessary to maintain control.</p> <p>DO NOT apply more than 0.2 lb. a.i. (0.77 pts. or 12.29 fl. oz. of product)/A per year.</p> <p>DO NOT graze livestock in treated areas.</p> |

Rate Conversion Chart

| Treated | 60 | 50 | 40 | 33 | 25 |
|-------------|------|------|-------|------|------|
| pts./A | 0.04 | 0.08 | 0.10 | 0.12 | 0.15 |
| fl. oz./A | 0.61 | 1.23 | 1.54 | 1.84 | 2.46 |
| lbs. a.i./A | 0.01 | 0.02 | 0.025 | 0.03 | 0.04 |

FOR USE ON PLANTS INTENDED FOR AESTHETIC PURPOSES OR CLIMATIC MODIFICATION AND BEING GROWN IN INTERIOR PLANSAPES, ORNAMENTAL GARDENS OR PARKS, OR ON LAWNS OR GROUNDS

- **DO NOT** apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. During application, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 1. Treatment to soil or vegetation around structures;
 2. Applications to lawns, turf, and other vegetation;
 3. Applications to building foundations, up to a maximum height of 3 feet above grade;
 4. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
 5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;

6. Applications made through the use of a coarse, low-pressure spray to only those portions of surfaces that are directly above bare soil., lawn, turf, mulch or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters, in order to control occasional invaders or aggregating pests.

- **DO NOT** water the treated area to the point of run-off.
- **DO NOT** make applications during rain.
- **DO NOT** apply directly to sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, except as directed by this label.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store in a cool, dry place. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

CONTAINER HANDLING:

Containers < 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Minibulk Containers: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follow: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Bulk Containers: Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes.

Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When the container is empty, replace the cap and seal all openings that have been opened during use; and return to the point of purchase, or to a designated location named at the time of purchase of this product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. **DO NOT** transport if this container is damaged or leaking. If the container is damaged or leaking, call CHEMTREC. If the container is damaged and leaking or material has been spilled, follow these procedures:

- Cover spill with absorbent material.
- Sweep into disposal container.

- Wash area with detergent and water and follow with clean water rinse.
- **DO NOT** allow to contaminate water supplies.
- Dispose of according to instructions.

If not returned to the point of purchase or to a designated location, clean empty container as instructed above and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations on liability, DO NOT use the product, and return it unopened to the Seller, and the purchase price will be refunded.

All trademarks are the property of their respective owners.

