



GUNSLINGER® AMP

PASTURE HERBICIDE

Specimen Label

For control of broadleaf weeds, including invasive and noxious weeds, and certain woody plants on rangeland, permanent grass pastures (including grasses grown for hay*), and Conservation Reserve Program (CRP) acres, non-cropland areas such as rights-of-way, roadsides, non-irrigation ditch banks, and natural areas such as wildlife management areas including seasonally dry flood plains, deltas, marshes, prairie potholes, or vernal pools, natural recreation areas, campgrounds, trailheads and trails, and grazed areas in and around these noncrop sites.

*Hay from grass treated with GUNSLINGER AMP PASTURE HERBICIDE within the preceding 18-months can only be used on the farm or ranch where the product is applied unless allowed under specific use directions for certain states on this label.

Not for Sale, Distribution, or Use in New York State. Not For Sale, Distribution, or Use in the San Luis Valley of Colorado.

ACTIVE INGREDIENTS:

Trisopropylammonium salt of 2-pyridine carboxylic acid, 4-amino-3,6-dichloro- 8.24%
 Dimethyl amine salt of (2,4-dichlorophenoxy) acetic acid 41.26%
OTHER INGREDIENTS: 50.50%
TOTAL: 100.00%
 Acid Equivalents:
 aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) – 4.28% - 0.41 lb/gal (50 g/L)
 2,4-D [(2,4-dichlorophenoxy) acetic acid] – 34.25% - 3.33 lb/gal (400 g/L)

EPA Reg. No. 81927-81

KEEP OUT OF REACH OF CHILDREN DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

Manufactured for:
 Alligare, LLC
 1565 5th Avenue
 Opelika, AL 36801

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils, when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.
- Chemical resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate

See engineering controls for additional requirements

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protections Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

AMINOPYRALID	GROUP 4	HERBICIDE
2,4-D	GROUP 4	HERBICIDE

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protections Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)].

User Safety Recommendations

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates and may be toxic to fish. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Take care to minimize the incidental overspray along the shoreline when applying to terrestrial plants at the water's edge or to water in areas where surface water is present. Do not apply directly to intertidal areas below the mean high water mark, except as permitted on this label. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not For Sale, Distribution, or Use in New York State.

Not For Sale, Distribution, or Use in the San Luis Valley of Colorado.

Not for use on pastures in Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, and Vermont. All other labeled uses are permitted in these states including grazed areas in and around these sites.



Light gray = states where use in pastures is not permitted
 Dark gray = NY where the product is not registered

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Protective eyewear
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS does not pertain to nonagricultural use on sites, such as, rangeland, permanent grass pastures, or non-cropland. See the Agricultural Use Requirements section below for information where the WPS applies.

Entry Restrictions for Non-WPS Uses: For applications on rangeland and permanent grass pastures, and non-cropland areas, do not enter or allow worker entry into treated areas until sprays have dried.

Product Information

GUNSLINGER® AMP PASTURE HERBICIDE controls broadleaf weeds, including invasive and noxious weeds, and certain woody plants on rangeland, permanent grass pastures (including grasses grown for hay*), and Conservation Reserve Program (CRP) acres, non-cropland areas such as rights-of-way, roadsides, non-irrigation ditch banks, and natural areas such as wildlife management areas including seasonally dry flood plains, deltas, marshes, prairie potholes, or vernal pools, natural recreation areas, campgrounds, trail-heads and trails, and grazed areas in and around these non-crop sites.

***Hay from grass treated with GUNSLINGER AMP PASTURE HERBICIDE within the preceding 18-months can only be used on the farm or ranch where the product is applied unless allowed under specific use directions for certain states on this label.**

Weed Resistance Management

For resistance management, GUNSLINGER AMP PASTURE HERBICIDE is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Avoid the consecutive use of this product or other target site of action Group 4 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.
- Scout fields after application to verify that the treatment was effective and to monitor weed populations for early signs of resistance development.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Report any incidence of non-performance of this product against a particular weed species to your Alligare LLC retailer, representative or call 888-255-4427. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemicals means to remove escapes, as practical, with the goal of preventing further seed production.

Development of plant populations resistant to this herbicide mode of action is usually not a problem on rangeland, permanent grass pastures, Conservation Reserve Program (CRP), or non-cropland sites since these sites receive infrequent pesticide applications.

Use Precautions

- Applications made during periods of intense rainfall, to soils saturated with water, surfaces paved with materials such as asphalt or concrete, or soils through which rainfall will not readily penetrate may result in runoff and movement of GUNSLINGER AMP PASTURE HERBICIDE. Injury to crops may result if treated soil and/or runoff water containing GUNSLINGER AMP PASTURE HERBICIDE is washed, or moved onto land used to produce crops. Exposure to GUNSLINGER AMP PASTURE HERBICIDE may injure or kill susceptible crops and other plants, such as grapes, soybeans, tobacco, sensitive ornamentals.
- **Seeding Grasses:**
 - **Preemergence:** Grasses may be reseeded in the fall following an application of GUNSLINGER AMP PASTURE HERBICIDE applied in the spring or early summer.
 - **Postemergence:** During the season of establishment, GUNSLINGER AMP PASTURE HERBICIDE should be applied only after perennial grasses are well established (have developed a good secondary root system and show good vigor). Most perennial grasses are tolerant to GUNSLINGER AMP PASTURE HERBICIDE at this stage of development. GUNSLINGER AMP PASTURE HERBICIDE may suppress certain established grasses, such as smooth bromegrass (*Bromus inermis*), especially when plants are stressed by adverse environmental conditions. Plants should recover from this transient suppression with the onset of environmental conditions favorable to grass growth and upon release from weed competition.
- **Field Bioassay Instructions:** In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, rainfall pattern or drainage. The field bioassay can be


initiated one year after the last application of aminopyralid in that field. Observe the test crop for symptoms of herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), epinasty, and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the intended rotational crop; plant only to wheat, corn, forage grasses, native grasses or grasses grown for hay.

Consult with an Alligare, LLC representative if you do not understand the "Use Precautions and Restrictions." Call (888-255-4427) for more information.

IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS

- Carefully read the section "**Restrictions in Hay or Manure Use.**"
- It is mandatory to follow the "**Use Precautions and Restrictions**" section of this label.
- Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- Hay can only be used on the farm or ranch where product is applied unless allowed under specific use directions for certain states on this label.
- Consult with an Alligare, LLC representative if you do not understand the "Use Precautions and Restrictions". Call (888) 255-4427

Forage and Manure Management



Pasture and Rangeland Restrictions

- Do not use grasses treated with GUNSLINGER AMP PASTURE HERBICIDE in the preceding 18-months for hay intended for export outside the United States.
- Hay from areas treated with GUNSLINGER AMP PASTURE HERBICIDE in the preceding 18-months CANNOT be distributed or made available for sale off the farm or ranch where harvested unless allowed under specific use directions for certain states on this label.
- Hay from areas treated with GUNSLINGER AMP PASTURE HERBICIDE in the preceding 18-months CANNOT be used for silage, haylage, baylage and green chop unless allowed under specific use directions for certain states on this label.
- Do not move hay made from grass treated with GUNSLINGER AMP PASTURE HERBICIDE within the preceding 18-months off farm unless allowed under specific use directions for certain states on this label.
- Do not use hay or straw from areas treated with GUNSLINGER AMP PASTURE HERBICIDE within the preceding 18-months or manure from animals feeding on hay treated with GUNSLINGER AMP PASTURE HERBICIDE in compost.
- Do not use grasses treated with GUNSLINGER AMP PASTURE HERBICIDE in the preceding 18-months for seed production.

Restrictions for All Uses

- This product is not intended for reformulation or repackaging into other end-use products.
- Do not treat frozen soil where runoff could damage sensitive plants.
- **Maximum seasonal rate:** Apply no more than 2.1 pints (34 fl oz) (0.87 lb acid equivalent 2,4-D) per acre per use season.
- Use 2 or more gallons of spray solution per acre
- Do not make more than two applications per year
- Do not apply within 30 days of previous application
- If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable
- **Maximum Application Rate:** Do not broadcast apply more than 2.1 pints (34 fl oz) per acre of GUNSLINGER AMP PASTURE HERBICIDE per year. The total amount of GUNSLINGER AMP PASTURE HERBICIDE applied broadcast, as a re-treatment, and/or spot treatment per year must not exceed 2.1 pints (34 fl oz) per acre. Spot treatments may be applied at an equivalent broadcast rate of up to 4.2 pints (68 fl oz) per acre of GUNSLINGER AMP PASTURE HERBICIDE per annual growing season; however, not more than 50% of an acre may be treated at that rate.
- **Grazing and Haying Restrictions:** Do not harvest forage for hay within 7 days of GUNSLINGER AMP PASTURE HERBICIDE application. Cutting hay too soon after spraying weeds can compromise the weed control. Wait 14 days prior to cutting grass hay to allow for maximum herbicide activity.
- **Do not use this product for impregnation on dry fertilizer, unless specified in an Alligare, LLC state specific product bulletin.**
- Do not apply this product on lawns, turf, ornamental plantings, urban walkways, driveways, tennis courts, golf courses, athletic fields, commercial sod operations, or other high-maintenance, fine turfgrass areas, or similar areas.
- **Transfer of Animals Feeding on GUNSLINGER AMP PASTURE HERBICIDE Treated Forage:** Do not transfer animals grazing or feeding on hay to areas where sensitive broadleaf crop occurs without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- **Restrictions in Hay or Manure Use:**
 - Do not use aminopyralid-treated plant residues, including hay or straw from areas treated within the preceding 18-months, in compost, mulch or mushroom spawn.
 - Do not use manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days, in compost, mulch or mushroom spawn.
 - Do not spread manure from animals that have grazed or consumed forage or hay from treated areas within the previous 3 days on land used for growing broadleaf crops.
 - Manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days may only be used on pasture grasses, grass grown for seed, wheat and corn.
 - Do not plant a broadleaf crop (including soybeans, sunflower, tobacco, vegetables, field beans, peanuts, and potatoes) in fields treated in the previous year with manure from animals that have grazed forage or eaten hay harvested from aminopyralid-treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.
 - To promote herbicide decomposition, plant residues should be evenly incorporated in the surface soil or burned. Breakdown of aminopyralid in plant residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.
- **Grazing Poisonous Plants:** Herbicide application may increase palatability of certain poisonous plants. Do not graze treated areas until poisonous plants are dry and no longer palatable to livestock.
- **Seeding Legumes:** Do not plant forage legumes until a soil bioassay has been conducted to determine if aminopyralid residues remaining in the soil will adversely affect the legume establishment.
- **Crop Rotation:** Do not rotate non-cropland to cropland for one year following an application of GUNSLINGER AMP PASTURE HERBICIDE. Cereals and corn can be planted one year after treatment. Most broadleaf crops are more sensitive and can require at least 2 years depending on the crop and environmental conditions. Do not plant a broadleaf crop until an adequately sensitive field bioassay shows that the level of aminopyralid present in the soil will not adversely affect that broadleaf crop.
- **GUNSLINGER AMP PASTURE HERBICIDE is highly active against many broadleaf plant species.** Do not use this product on areas where loss of desirable broadleaf forage plants, including legumes, cannot be tolerated.
- Trees adjacent to or in a treated area can occasionally be affected by root uptake of GUNSLINGER AMP PASTURE HERBICIDE through movement into the soil. Do not apply GUNSLINGER AMP PASTURE HERBICIDE within the root zone of desirable trees unless such injury can be tolerated. Use special caution near roses, and leguminous trees such as locusts, redbud, mimosa, and caragana.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Do not contaminate water intended for irrigation or domestic purposes.** Do not treat inside banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes.

Grass Harvested for Hay Intended for Distribution or Sale off the Farm or Ranch and Grass Harvested for Silage, Haylage, Baylage, or Green Chop Intended for Use On the Farm or Ranch

(For use only in the states of AL, AR, AZ, CO, FL, GA, ID, KS, KY, LA, MO, MS, MT, ND, NE, NV, NM, OK, SD, TN, TX, UT, WY)

Restrictions

- **The Applicator must provide the land manager with a copy of the Alligare, LLC Stewardship instructions regarding uses of forage from areas treated with aminopyralid.**
- **Do not use grasses treated with GUNSLINGER AMP PASTURE HERBICIDE in the preceding 18-months for hay intended for export outside the United States.**
- **Do not use hay or straw from areas treated with GUNSLINGER AMP PASTURE HERBICIDE within the preceding 18-months, or manure from animals feeding on hay treated with GUNSLINGER AMP PASTURE HERBICIDE, in compost.**
- **Grazing and Haying Restrictions:** Do not harvest forage for hay within 7 days of GUNSLINGER AMP PASTURE HERBICIDE application. Cutting hay too soon after spraying weeds can compromise weed control. Wait 14 days prior to cutting grass hay to allow for maximum herbicide activity.
- **Transfer of Animals Feeding on GUNSLINGER AMP PASTURE HERBICIDE-Treated Forage:** Do not transfer animals grazing or feeding on hay from areas treated with GUNSLINGER AMP PASTURE HERBICIDE to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- **Grazing Poisonous Plants:** Herbicide application may increase palatability of certain poisonous plants. Do not graze treated areas until poisonous plants are dry and no longer palatable to livestock.
- **Restrictions in Hay or Manure Use:**
 - Do not use treated plant residues, including hay or straw from areas treated within the preceding 18-months in compost, mulch or mushroom spawn.
 - Do not use manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days, in compost, mulch or mushroom spawn.
 - Do not spread manure from animals that have grazed or consumed forage or hay from treated areas within the previous 3 days on land used for growing broadleaf crops.
 - Manure from animals that have grazed forage or eaten hay harvested from aminopyralid-treated areas within the previous 3 days may only be used on pasture grasses, grass grown for seed, wheat and corn.
 - Do not plant a broadleaf crop in fields treated in the previous year with manure from animals that have grazed forage or eaten hay harvested from aminopyralid-treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid residues in the soil is at level that is not injurious to the crop to be planted.
 - To promote herbicide decomposition, plant residues should be evenly incorporated in the surface soil or burned. Breakdown of aminopyralid in plant residues or manure is more rapid under warm, moist soil conditions and may be accelerated by supplemental irrigation.
- **Preharvest Interval:** Do not cut forage for hay within 7 days of application. For program lands, such as CRP, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Crop Rotation:** Do not rotate non-cropland to cropland for one year following an application of GUNSLINGER AMP PASTURE HERBICIDE. Do not plant a broadleaf crop until an adequately sensitive field bioassay shows that the level of aminopyralid present in the soil will not adversely affect that broadleaf crop.
- **GUNSLINGER AMP PASTURE HERBICIDE is highly active against many broadleaf plant species.** Do not use this product on areas where loss of desirable broadleaf forage plants, including legumes, cannot be tolerated.
- Trees adjacent to or in a treated area can occasionally be affected by root uptake of GUNSLINGER AMP PASTURE HERBICIDE through movement into the soil. Do not apply GUNSLINGER AMP PASTURE HERBICIDE within the root zone of desirable trees unless such injury can be tolerated. Use special caution near roses, and leguminous trees such as locusts, redbud, mimosa, and caragana.

Field Bioassay Instructions: In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, rainfall pattern or drainage. The field bioassay can be initiated one year after the last application of aminopyralid in that field. Observe the test crop for symptoms of herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the intended rotational crop; plant only to wheat, forage grasses, native grasses or grasses grown for hay.

Spray Drift Management

Avoid application under conditions that may allow spray drift because very small quantities of spray, which may not be visible, may injure susceptible crops. This product should be applied only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, non-target crops and other plants) is minimal (e.g., when wind is blowing away from the sensitive areas). A drift control aid may be added to the spray solution to further reduce the potential for drift. If a drift control aid is used, follow the use directions and precautions on the manufacturer's label. Do not use a thickening agent with Microfoil, Thru-Valve booms, or other spray delivery systems that cannot accommodate thickened spray solutions.

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, fruit trees, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that may not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Aerial Application

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Ground Boom Application

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Non-Cropland Areas

GUNSLINGER AMP PASTURE HERBICIDE may be applied alone or in tank mix combination to non-cropland areas, such as non-irrigation ditch banks, industrial and storage areas, airports, roadsides, railroad and utility rights-of-way, including grazed areas on these sites as an aerial or ground broadcast treatment, as a spot application, or as a high volume foliar application (see Application Methods section). Refer to the Broadleaf Weeds Controlled section for application rates specified for specific broadleaf weeds.

Restrictions:

Postemergence (annual and perennial weeds):

- Limited to 2 applications per year
- Maximum of 2.1 pints (34 fl oz) (0.87 lb ae/acre 2,4-D)/acre per application per year
- Minimum of 30 days between applications

Postemergence (woody plants)

- Limited to 1 application per year
- Maximum of 2.1 pints (34 fl oz) (0.87 lb ae/acre 2,4-D)/acre per year

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Restrictions for Non-Irrigation Canal Ditchbank Application and Terrestrial Weeds near and up to the Water's Edge

Use Rate Restrictions:

Limited to 2 applications per year
 Minimum of 30 days between applications
 Maximum of 2.1 pints (34 fl oz) (0.87 lb ae)/acre per broadcast application.
 Do not apply more than 2.1 pints (34 fl oz) per acre per year.

Spot treatments may be applied at an equivalent broadcast rate of up to 4.2 pints (68 fl oz) of GUNSLINGER AMP PASTURE HERBICIDE (1.75 lbs acid equivalent) per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate.

Do not use on small canals with a flow rate less than 10 cubic feet per second (CFS) where water will be used for drinking purposes. CFS may be estimated by using the formula

below. The approximate velocity needed for the calculation can be determined by observing the length of time that it takes a floating object to travel a defined distance. Divide the distance (ft.) by the time (sec.) to estimate velocity (ft. per sec.). Repeat 3 times and use the average to calculate CFS.

$$\text{Average Width (ft.)} \times \text{Average Depth (ft.)} \times \text{Average Velocity (ft. per sec.)} = \text{CFS}$$

For ditchbank weeds:

Do not allow boom spray to be directed onto water surface.
 Do not spray across stream to opposite bank.

For shoreline weeds:

Allow no more than 2 foot overspray onto water.

Sprayer Clean-Out Instructions

It is recommended to use separate spray equipment on highly sensitive crops such as tobacco, soybeans, potatoes, peanuts and tomatoes.

Do not use spray equipment used to apply GUNSLINGER AMP PASTURE HERBICIDE for other applications to land planted to, or to be planted to, crops or desirable sensitive plants, unless it has been determined that all residues of this herbicide have been removed by thorough cleaning of equipment.

Equipment used to apply GUNSLINGER AMP PASTURE HERBICIDE should be thoroughly cleaned before reusing to apply any other chemicals as follows:

- Rinse and flush application equipment thoroughly after use. Dispose of rinse water away from water supplies.
- Rinse a second time, adding 1 quart of household ammonia or tank cleaning agent for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 minutes). Let the solution stand for several hours, preferably overnight.
- Flush the solution out of the spray tank through the boom.
- Rinse the system twice with clean water, recirculating and draining each time.
- Spray nozzles and screens should be removed and cleaned separately.

Application Methods

Apply the specified rate of GUNSLINGER AMP PASTURE HERBICIDE as a coarse low-pressure spray. **Do not apply this product with mist blower systems that deliver very fine spray droplets. Use of mist blower equipment can reduce weed control and increase spray drift potential.**

Spray volume should be sufficient to uniformly cover foliage. Increase spray volume to ensure thorough and uniform coverage when target vegetation is tall and/or dense. To enhance foliage wetting and coverage, an approved non-ionic agricultural surfactant may be added to the spray mixture as specified by the surfactant label.

Ground Broadcast Application: Higher spray volumes (greater than 10 gallons per acre) generally provides better coverage and better control, particularly in dense and/or tall foliage.

Aerial Broadcast Application: Do not apply less than 2 gallons per acre total spray volume. Five gallons per acre or greater will generally provide better coverage and better control, particularly in dense and/or tall foliage.

High-Volume Foliar Application: High volume foliar treatments may be applied at rates equivalent to a maximum of 2.1 pints (34 fl oz) per acre per annual growing season. Use sufficient spray volume to thoroughly and uniformly wet foliage and stems.

Spot Application: Spot treatments may be applied at rates equivalent to broadcast-applied rate of up to a maximum of 4.2 pints (68 fl oz) on 50% of the treated field. Spray volume should be sufficient to thoroughly and uniformly wet weed foliage. Repeat treatments may be made, but the total amount of GUNSLINGER AMP PASTURE HERBICIDE applied must not exceed 2.1 pints (34 fl oz) per acre per year (see comments in the Use Precautions and Restrictions section above on Maximum Application Rate).

Table 1: Amount of GUNSLINGER AMP PASTURE HERBICIDE (in fl oz) to mix in 3 gallons of water

GPA	GUNSLINGER AMP PASTURE HERBICIDE Amount (in fluid oz) To Mix In 3 Gal of Water With Various Application Rates		
	19 fl oz/A	24 fl oz/A	34 fl oz/A
20	2.9	3.6	5.1
30	1.9	2.4	3.4
40	1.4	1.8	2.6
50	1.1	1.4	2.0
60	1.0	1.2	1.7
70	0.8	1.0	1.4
80	0.7	0.9	1.3
90	0.6	0.8	1.1
100	0.6	0.7	1.0

Table 2: Application rates in the table below are based on treating an area of 1000 sq ft. An area of 1000 sq ft is about 10.5 by 10.5 yards in size. Mix the amount of GUNSLINGER AMP PASTURE HERBICIDE (fl oz or milliliters) corresponding to the desired broadcast rate in 0.5 to 2.5 gallons of water, depending upon the spray volume required to treat 1000 sq ft. A delivery volume of 0.5 gallons per 1000 sq ft is equivalent to 22 gallons per acre and 2.5 gallons per 1000 sq ft is equivalent to 109 gallons per acre.

Amount of GUNSLINGER AMP PASTURE HERBICIDE per 1000 sq ft to Equal Broadcast Rate			
Broadcast Rate		Amount of GUNSLINGER AMP PASTURE HERBICIDE per 1000 sq ft	
(fl oz/acre)	(pt/acre)	(fl oz)	(mL)
19	1.2	0.44	13
24	1.5	0.55	16
34	2.1	0.78	23

Note: 1 mL = 1 cc and 1 fluid ounce (fl oz) = 29.6 milliliters (mL) = 2 tablespoons = 6 teaspoons

To calculate the amount of GUNSLINGER AMP PASTURE HERBICIDE for areas larger than 1000 sq ft: Multiply the table value (fl oz or milliliters) by the area to be treated in "thousands" of square feet. For example, if the area to be treated is 3500 sq ft, multiply the table value by 3.5 (3500 sq ft divided by 1000 sq ft = 3.5).

Mixing Instructions

Mixing with Water

To prepare the spray, add about half the required amount of water in the spray tank. Then, with agitation, add the specified amount of GUNSLINGER AMP PASTURE HERBICIDE and other registered tank mix herbicides. Finally, with continued agitation, add the rest of the water and additives such as surfactants or drift control and deposition aids.

Addition of Surfactants or Adjuvants on All Labeled Use Sites: The addition of a high quality non-ionic surfactant (of at least 80% active ingredient) at 0.25 to 0.5 % volume per volume (1 to 2 quarts per 100 gallons of spray) is recommended to enhance herbicide activity under adverse environmental conditions (such as, high temperature, low relative humidity, drought conditions, dusty plant surfaces) or when weeds are heavily pubescent or more mature.

Tank Mixing with Other Herbicides

GUNSLINGER AMP PASTURE HERBICIDE at rates of up to 2.1 pints (34 fl oz) per acre may be mixed with labeled rates of other labeled herbicides including Alligare Triclopyr 4 (EPA Reg. No. 81927-11), Alligare Cleargraze Pasture Herbicide (EPA Reg. No. 81927-65), Boulder 6.3 (EPA Reg. No. 91927-54), Alligare Triclopyr 3 (EPA Reg. No. 81927-13), Alligare Triumph 22K Herbicide (EPA Reg. No. 81927-18) to broaden the spectrum of weeds and brush controlled or to improve control of certain weeds. GUNSLINGER AMP PASTURE HERBICIDE may be applied in tank mix combination with labeled rates of other herbicides provided: (1) the tank mix product is labeled for the timing and method of application for the use site to be treated and (2) mixing is not prohibited by the label of the registered tank mixed products, and (3) that the tank mix combination is physically compatible (see tank mix compatibility testing below). When tank mixing, use only in accordance with the restrictions, precautions and limitations on the respective product labels.

- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Do not exceed specified application rates. If products containing the same active ingredient are mixed, do not exceed the maximum allowable active ingredient use rates.
- For direct injection or other spray equipment where the product formulations will be mixed in undiluted form, special care should be taken to ensure tank mix compatibility.
- Always perform a jar test to ensure the compatibility of products to be used in tank mixture.

Tank Mixing Precautions:

- For products packaged in water soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment has been adequately cleaned. (See Sprayer Clean-Out instructions.)
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

Tank Mix Compatibility Testing: Perform a jar test prior to mixing in a spray tank to ensure compatibility of GUNSLINGER AMP PASTURE HERBICIDE and other pesticides or carriers. Use a clear glass jar with lid and mix ingredients in the same order and proportions as will be used in the spray tank. The mixture is compatible if the materials mix readily when the jar is inverted several times. The mixture should remain stable after standing for 1/2 hour or, if separation occurs, should readily remix if agitated. An incompatible mixture is indicated by separation into distinct layers that do not readily remix when agitated and/or the presence of flakes, precipitates, gels, or heavy oily film in the jar. Use of an appropriate compatibility aid may resolve mix incompatibility.

Mixing with Sprayable Liquid Fertilizer Solutions

GUNSLINGER AMP PASTURE HERBICIDE is usually compatible with liquid fertilizer solutions. It is anticipated that GUNSLINGER AMP PASTURE HERBICIDE will not require a compatibility agent for mixing with fertilizers; however, a compatibility test (jar test) should be made prior to large scale batch mixing. Jar tests are particularly important when a new batch of fertilizer or pesticide is used, when water sources change, or when tank mixture ingredients or concentrations are changed. Compatibility may be determined by mixing the spray components in the desired order and proportions in a clear glass jar before large scale mixing of spray components in the spray tank. Use of a compatibility agent could be used to help obtain and maintain a uniform spray solution during mixing and application. When

mixing GUNSLINGER AMP PASTURE HERBICIDE in liquid fertilizer, mix GUNSLINGER AMP PASTURE HERBICIDE in water (in a 1:1 ratio at the minimum) and add to the spray tank first, then add the liquid fertilizer to the spray tank. **Note:** The lower the temperature of the liquid fertilizer, the greater the likelihood of mixing problems. **Mixing GUNSLINGER AMP PASTURE HERBICIDE in N-P or N-P-K liquid fertilizer solutions is more difficult than mixing with straight nitrogen fertilizer and should not be attempted without first conducting a successful compatibility jar test.** Agitation in the spray tank must be vigorous to be comparable with jar test agitation. Apply the spray mixture the same day it is prepared while maintaining continuous agitation. Rinse the spray tank thoroughly after use.

Suggested Mixing Procedure:

- With continuous vigorous agitation dilute herbicide with water (1 part herbicide to 2 parts water) before adding to liquid nitrogen fertilizer solution.
- Apply as soon as mixing is complete, maintaining continuous, vigorous agitation throughout mixing and application without interruption.
- Application during very cold (near freezing) weather is not advisable. The likelihood of mixing or compatibility problems with liquid fertilizer increases under cold conditions.
- Do not store the spray mixture.

Note: Foliar-applied liquid fertilizers themselves can cause yellowing of the foliage of forage grasses and other vegetation.

Use Rates and Timing

Do not use GUNSLINGER AMP PASTURE HERBICIDE if loss of legumes species or other broadleaf species cannot be tolerated.

GUNSLINGER AMP PASTURE HERBICIDE may be applied postemergence as a broadcast spray or as a spot application to control weeds listed on this label; weeds other than those listed may also be controlled by this herbicide. When a rate range is given, use a higher rate in the range to control weeds at advanced growth stages or under less-than-favorable growing conditions (e.g., drought stress) or for longer residual control. Best weed control results are obtained when spray volume is sufficient to provide uniform coverage of treated plants. For optimum uptake and translocation of the herbicide, avoid mowing, haying, shredding, burning or soil disturbance in treated areas for at least 7 days following application.

For most species, 2 hours between application and rainfall provides a sufficient amount of time to avoid loss in weed control due to herbicide wash-off of the treated foliage.

GUNSLINGER AMP PASTURE HERBICIDE also provides preemergence control of germinating seeds or emerging seedlings of susceptible weeds and re-growth of certain perennial weeds following application. Weed establishment following GUNSLINGER AMP PASTURE HERBICIDE application will depend upon application rate, season of application, and growing condition.

GUNSLINGER AMP PASTURE HERBICIDE can provide long-term control of weeds. The length of control is dependent upon the application rate, condition and growth stage of target weeds, environmental conditions at and following application, and the density and vigor of competing desirable vegetation. Long-term broadleaf weed control is most effective where forage grasses are allowed to recover from overgrazing, drought, etc., and compete with weeds.

GUNSLINGER AMP PASTURE HERBICIDE can be an important component of integrated vegetation management programs designed to renovate or restore desired non-cropland plant communities. To maximize and extend the benefits of weed control provided by GUNSLINGER AMP PASTURE HERBICIDE, it is important that vegetation management practices, including grazing management, biological control agents, replanting, fertilization, prescribed fire, reseeding with desirable plants, etc., be used to increase the competitiveness of desired forages. Used as part of an integrated management program, GUNSLINGER AMP PASTURE HERBICIDE can serve as a catalyst for rapid improvement of rangeland, permanent grass pasture, and CRP, and non-cropland sites by alleviating the adverse competitive effect of weeds on the yield and quality of forages and other desirable plant species. Agricultural and natural resources specialists with federal and state government agencies can provide guidance on best management practices and development of integrated vegetation management systems.

Broadleaf Weeds Controlled

Early to mid-spring applications. GUNSLINGER AMP PASTURE HERBICIDE can be applied at rates between 0.8 and 1.2 pints (13 to 19 fl oz) product per acre in early to mid-spring when weeds are less than 2 inches tall; applications in this rate range are most effective when conditions are favorable for plant growth.

The following weeds will be controlled at 1.2 to 2.1 pints (19 to 34 fl oz) product per acre. For best results, apply when weeds are actively growing and conditions are favorable for plant growth. Use a higher rate in the rate range when growing conditions are less than favorable, when weeds are mature, when weed foliage is tall and dense or when residual control is important. GUNSLINGER AMP PASTURE HERBICIDE also provides preemergence control of germinating seeds or seedlings of susceptible weeds that emerge following application. Increasing application rate to the high end of the rate range specified will extend period of residual control.

Table 3: Broadleaf Weeds Controlled

Weed Species			
Common Name	Scientific Name	Life Cycle ^c	Plant Family
Rate Range: 1.2 to 1.5 pints (19 to 24 fl oz) product per acre			
Bedstraw	<i>Galium</i> spp.	perennial	Rubiaceae
bedstraw, smooth	<i>Galium mollugo</i>	perennial	Rubiaceae
broomweed, annual ^a	<i>Amphichyris dracunculoides</i>	annual	Asteraceae
carrot, wild ^a	<i>Daucus carota</i>	biennial	Apiaceae
Cinquefoil, hoary	<i>Potentilla argentea</i>	perennial	Rosaceae
cinquefoil, sulfur ^{a,b}	<i>Potentilla recta</i>	perennial	Rosaceae
clover, sweet	<i>Melilotus officinalis</i>	biennial	Fabaceae

Weed Species			
Common Name	Scientific Name	Life Cycle ^c	Plant Family
clover, white	<i>Trifolium repens</i>	perennial	Fabaceae
cocklebur ^a	<i>Xanthium strumarium</i>	annual	Asteraceae
croton, woolly ^{a,b}	<i>Croton capitatus</i>	annual	Euphorbiaceae
crownvetch ^a	<i>Securigera varia</i>	perennial	Fabaceae
daisy, oxeye ^{a,b}	<i>Leucanthemum vulgare</i>	perennial	Asteraceae
falsedandelion, Carolina ^a	<i>Pyrrhopappus carolinianus</i>	annual/ biennial	Asteraceae
fleabane, annual ^a	<i>Erigeron annuus</i>	annual	Asteraceae
horsenettle, Carolina ^{a,b}	<i>Solanum carolinense</i>	perennial	Solanaceae
lettuce, prickly ^a	<i>Lactuca serriola</i>	annual	Asteraceae
pokeweed, common	<i>Phytolacca americana</i>	perennial	Phytolaccaceae
ragweed, common ^{a,b}	<i>Ambrosia artemisiifolia</i>	annual	Asteraceae
ragweed, western	<i>Ambrosia psilostachya</i>	perennial	Asteraceae
ragwort, tansy ^{a,c}	<i>Senecio jacobaea</i>	perennial	Asteraceae
sneezeweed, bitter ^a	<i>Helenium amarum</i>	annual	Asteraceae
starthistle, yellow ^{a,c}	<i>Centaurea solstitialis</i>	annual	Asteraceae
thistle, bull ^{a,b}	<i>Cirsium vulgare</i>	biennial	Asteraceae
thistle, musk ^{a,b}	<i>Carduus nutans</i>	biennial	Asteraceae
thistle, plumeless ^{a,b}	<i>Carduus acanthoides</i>	biennial	Asteraceae
vervain, blue ^a	<i>Verbena hastata</i>	perennial	Asteraceae
vervain, hoary ^a	<i>Verbena stricta</i>	perennial	Asteraceae
vetch, common ^a	<i>Vicia sativa</i>	annual	Fabaceae
woodsorrel, yellow ^a	<i>Oxalis stricta</i>	perennial	Oxalidaceae
wormwood, absinth ^{a,b}	<i>Artemisia absinthium</i>	perennial	Asteraceae
Rate Range: 1.5 to 2.1 pints (24 to 34 fl oz) product per acre			
acacia spp. ^{a,†}	<i>Acacia</i> sp.	perennial	Fabaceae
actinomeris, wingstem	<i>Verbesina alternifolia</i>	perennial	Asteraceae
beebalm, pony ^{a,b} (horse mint)	<i>Monarda pectinata</i>	annual	Lamiaceae
blackberry spp. ^{a,†}	<i>Rubus</i> sp.	perennial	Rosaceae
Blackbrush ^{a,†}	<i>Acacia rigidula</i>	perennial	Fabaceae
bullnettle, Texas ^f	<i>Cnidocolus texanus</i>	perennial	Euphorbiaceae
amaranth, spiny ^a	<i>Amaranthus spinosus</i>	annual	Amaranthaceae
burdock, common ^{a,b}	<i>Arctium minus</i>	biennial	Asteraceae
buttercup, hairy ^a	<i>Ranunculus sardous</i>	perennial	Ranunculaceae
buttercup, tall ^{a,b}	<i>Ranunculus acris</i>	perennial	Ranunculaceae
camphorweed ^a	<i>Heterotheca subaxillaris</i>	annual	Asteraceae
chickweed, common ^a	<i>Stellaria media</i>	annual	Caryophyllaceae
chicory ^a	<i>Cichorium intybus</i>	perennial	Asteraceae
Coneflower, upright prairie ^g	<i>Ratibida columnifera</i>	perennial	Asteraceae
cudweed, purple	<i>Gnaphalium purpureum</i>	annual	Asteraceae
dandelion, common ^a	<i>Taraxacum officinale</i>	perennial	Asteraceae
dock, broadleaf ^a	<i>Rumex obtusifolius</i>	perennial	Polygonaceae
dock, curly ^a	<i>Rumex crispus</i>	perennial	Polygonaceae
dogfennel ^c	<i>Eupatorium capillifolium</i>	perennial	Asteraceae
evening primrose, cutleaf ^d	<i>Oenothera lacinata</i>	annual	Asteraceae
false dandelion, Carolina ^a	<i>Tragopogon dubius</i>	biennial	Asteraceae
fiddleneck, common	<i>Amsinckia intermedia</i>	annual	Boraginaceae
fireweed	<i>Epilobium angustifolium</i>	perennial	Onagraceae
fleabane, annual ^a	<i>Erigeron annuus</i>	annual	Asteraceae
goldenrod, Canada ^a	<i>Solidago canadensis</i>	perennial	Asteraceae
goldenrod, Missouri ^a	<i>Solidago missouriensis</i>	perennial	Asteraceae
goldenrod, rigid	<i>Solidago rigida</i>	perennial	Asteraceae
gumweed, curlycup	<i>Grindelia squarrosa</i>	biennial	Asteraceae
hawkweed, orange ^{a,b}	<i>Hieracium aurantiacum</i>	perennial	Asteraceae
hawkweed, yellow ^{a,b}	<i>Hieracium pratense</i>	perennial	Asteraceae
henbit ^a	<i>Lamium amplexicaule</i>	annual/ biennial	Lamiaceae
honeylocust	<i>Gleditsia triacanthos</i>	perennial	Fabaceae
horehound [†]	<i>Marrubium vulgare</i> L.	perennial	Lamiaceae
horseweed ^a	<i>Conyza canadensis</i>	annual	Asteraceae

Weed Species			
Common Name	Scientific Name	Life Cycle ^c	Plant Family
Huisache ^{a,†}	<i>Acacia farnesiana</i>	perennial	Fabaceae
indigo, blue	<i>Baptisia australis</i>	perennial	Fabaceae
ironweed, tall	<i>Vernonia gigantea</i>	perennial	Asteraceae
ironweed, western	<i>Vernonia baldwinii</i>	perennial	Asteraceae
knapweed	<i>Centaurea</i> spp.	biennial	Asteraceae
knapweed, brown	<i>Centaurea jacea</i>	perennial	Asteraceae
knapweed, diffuse ^{a,b}	<i>Centaurea diffusa</i>	biennial/ perennial	Asteraceae
knapweed, Russian ^{a,b}	<i>Acroptilon repens</i>	perennial	Asteraceae
knapweed, spotted ^{a,b}	<i>Centaurea stoebe</i>	biennial/ perennial	Asteraceae
knapweed, squarrose	<i>Centaurea virgata</i>	biennial/ perennial	Asteraceae
kudzu ^{a,b}	<i>Pueraria montana</i>	perennial	Fabaceae
lambsquarters, common ^a	<i>Chenopodium album</i>	annual	Chenopodiaceae
lespedeza, annual	<i>Lespedeza striata</i>	annual	Fabaceae
locust, black	<i>Robinia pseudoacacia</i>	woody perennial	Fabaceae
marshelder, annual ^a	<i>Iva annua</i>	annual	Asteraceae
mayweed, scentless ^a	<i>Tripleurospermum perforate</i>	annual	Asteraceae
mayweed, stinking ^{a,b}	<i>Anthemis cotula</i>	annual	Asteraceae
medic, black ^a	<i>Medicago lupulina</i>	perennial	Fabaceae
mesquite, honey ^{a,†}	<i>Prosopis glandulosa</i>	perennial	Fabaceae
Mexican-tea	<i>Dysphania ambrosioides</i>	annual/ perennial	Chenopodiaceae
mullein ^e	<i>Verbascum</i> spp.	biennial	Scrophulariaceae
nightshade, silverleaf [†]	<i>Solanum elaeagnifolium</i>	perennial	Solanaceae
partridgepea ^a	<i>Chamaecrista fasciculata</i>	annual	Fabaceae
plantain, broadleaf ^a	<i>Plantago major</i>	perennial	Plantaginaceae
plantain, buckhorn ^a	<i>Plantago lanceolata</i>	perennial	Plantaginaceae
pricklyash, lime [†]	<i>Zanthoxylum fagara</i>	perennial	Fabaceae
ragweed, false	<i>Parthenium hysterophorus</i>	annual	Asteraceae
Rose, Macartney [†]	<i>Rosa bracteata</i>	perennial	Fabaceae
rose, multiflora ^e	<i>Rosa multiflora</i>	perennial	Rosaceae
sicklepod ^a	<i>Cassia obtusifolia</i>	perennial	Fabaceae
sida, prickly [†]	<i>Sida spinosa</i>	annual	Malvaceae
smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	annual	Polygonaceae
snow-on-the-mountain [†]	<i>Euphorbia marginata Pursh</i>	annual	Euphorbiaceae
soda apple, tropical ^{a,b}	<i>Solanum viarum</i>	perennial	Solanaceae
sowthistle, perennial ^{a,b}	<i>Sonchus arvensis</i>	perennial	Asteraceae
sowthistle, prickly ^a	<i>Sonchus asper</i>	annual	Asteraceae
Spanish needles	<i>Bidens bipinnata</i>	annual	Asteraceae
starthistle, yellow ^{a,b}	<i>Centaurea solstitialis</i>	annual	Asteraceae
sumac, smooth	<i>Rosa glabra</i>	perennial	Anacardiaceae
sunflower, common ^a	<i>Helianthus annua</i>	annual	Asteraceae
tallow tree, Chinese ^{a,†}	<i>Triadica loureiroi</i>	perennial	Euphorbiaceae
Teasel ^a	<i>Dipsacus</i> spp.	biennial	Dipsacaceae
thistle, Canada ^{a,b}	<i>Cirsium arvense</i>	perennial	Asteraceae
thistle, scotch	<i>Onopordum acanthium</i>	biennial	Asteraceae
yarrow, common ^a	<i>Achillea millefolium</i>	perennial	Asteraceae

^aThese plants are indicated to be invasive in the USDA-NRCS, PLANTS Database (<http://plants.usda.gov/index.html>).

^bPlants designated as noxious weeds in at least one state (PLANTS Database, USDA-NRCS, <http://plants.usda.gov/index.html>).

^cSpot treatment at rates up to 4.2 pints (68 fl oz) per acre of GUNSLINGER AMP PASTURE HERBICIDE may be particularly effective against dense patches of perennial broadleaf plants.

^dapply during rosette stage

^esee specific use directions below for multiflora rose.

^fapply at flowering stage

^gapply when actively growing before flowering

[†]Suppression only

Specific Use Directions

Multiflora rose, individual plant treatment - Use 2.1 pints (34 fl oz) of GUNSLINGER AMP PASTURE HERBICIDE in 100 gal of water with 2 pints (32 fl oz) or 0.25% v/v of a non-ionic surfactant. Or, 1.5 pints (24 fl oz) of GUNSLINGER AMP PASTURE HERBICIDE can be tank mixed with the appropriately labeled amount of Alligare Triclopyr 4 (EPA Reg. No. 81927-11). Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing. Spot treatments may be applied at an equivalent broadcast rate of up to 4.2 pints (68 fl oz) of GUNSLINGER AMP PASTURE HERBICIDE per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate.

Multiflora rose, broadcast treatment: 1.5 to 2.1 pints (24 to 34 fl oz) per acre of GUNSLINGER AMP PASTURE HERBICIDE can be tank mixed with the appropriately labeled amount of Alligare Triclopyr 4 (EPA Reg. No. 81927-11) per acre. Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing.

Control of Terrestrial Weeds near and up to the Water's Edge

GUNSLINGER AMP PASTURE HERBICIDE can be used to treat terrestrial weeds that extend up to the water's edge. **Do not apply directly to water.** This product must not be used to treat vegetation standing in the water. When controlling terrestrial weed species near and up to the water's edge, take precautions to minimize incidental overspray to the adjacent water. Consult local public water control authorities before applying this product near public waters. Permits may be required to treat such areas. Apply the specified rate, listed in Table 2, of GUNSLINGER AMP PASTURE HERBICIDE as a coarse low-pressure spray as ground broadcast or spot applications. Do not apply aerially for control of weeds growing at or near the water's edge. Spray volume should be sufficient to uniformly cover foliage. Increase the spray volume to ensure thorough and uniform coverage when target vegetation is tall and/or dense. It is also permissible to treat target weeds within dry non-irrigation ditches and seasonally dry transitional areas between upland and lowland sites (such as flood plains, deltas, marshes, prairie potholes or vernal pools), but only at times when those sites are dry and are forecasted or managed by water control systems to remain dry for at least 2 weeks following application.

STORAGE AND DISPOSAL

Do not contaminate water, food, feed or fertilizer by storage or disposal.

Pesticide Storage: If this product is exposed to subfreezing temperatures, the active ingredient may crystallize and settle out of solution. Under these conditions the product should be warmed to at least 40°F and agitated well to dissolve any crystallized active ingredient prior to use.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Container Handling:

[Nonrefillable Containers 5 gallons or less:] [Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Nonrefillable containers larger than 5 gallons:] [Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Refillable containers larger than 5 gallons:] [Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.]

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

Warranty: Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. To the extent consistent with applicable law, no such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

Limitation of Liability: To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

GUNSLINGER® is a registered trademark of Alligare, LLC

EPA 20201002