

Harvista™ 1.3 SC

Protects fruit and vegetable crops from the various effects of ethylene.

ACTIVE INGREDIENT:	% By WT.
1-Methylcyclopropene	1.3%
OTHER INGREDIENTS:	98.7%
TOTAL:	100.0%

Produced for: AgroFresh Inc.
510-530 Walnut Street, Suite 1350
Philadelphia PA 19106, USA
Tel: 1-866-206-1001

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unused within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

[See [front][back][side][inside] panel for First Aid Instructions and [Leaflet][Booklet] for complete Precautionary Statements and Directions for Use.

FIRST AID	
If in eyes	<ul style="list-style-type: none">▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.▪ Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">▪ Take off contaminated clothing.▪ Rinse skin immediately with plenty of water for 15-20 minutes.▪ Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">▪ Move person to fresh air.▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.▪ Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none">▪ Call a poison control center or doctor immediately for treatment advice.▪ Have person sip a glass of water for 15-20 minutes.▪ Call a poison control center or doctor for treatment advice.
POISON CONTROL CENTER 1-800-222-1222	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378. For emergencies, call the poison control center 1-800-222-1222.	

Lot No. on container

Net Contents (on container):

EPA Reg. No. 71297-17

Net content on package

EPA Est. No. 71297-WA-1

110197_20220009US

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to Supplemental Labeling entitled AGRICULTURAL USE REQUIREMENTS in the DIRECTIONS FOR USE section of the labeling for information about this standard.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking chewing gum using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Avoid breathing vapor.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (made of any waterproof material)
- Protective eyewear (such as goggles, safety glasses, or face shield)

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been fogged or heavily contaminated with the concentrated product. Do not reuse them.

USER SAFETY RECOMMENDATIONS: Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing / PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial use: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. This pesticide is moderately toxic to fish and aquatic invertebrates.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 4 hours** unless wearing the appropriate PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Long -sleeved shirt and long pants
- Chemical-resistant gloves (made of any waterproof material)
- Shoes plus socks
- Protective eyewear (such as goggles, safety glasses, or face shield)

Read the entire label before using this product.

APPLICATION INSTRUCTIONS

HARVISTA 1.3 SC is packaged in plastic containers, such as pails and totes. While the package is opened, stand upwind from the container. **DO NOT ADD ANY WATER TO THE CONTAINER.**

Rate, time and mode of application: Refer to instructions in crop tables.

Plant Safety: This product has been tested on numerous plant varieties with no phytotoxic effects. However, since this product has not been tested on all plant varieties or in combination with all available adjuvants or tank mixes, the manufacturer recommends testing on a small number of plants to check for adverse plant effects before applying on a larger number of plants.

Incompatibility: Do not use with products containing copper.

For treatment of FRUITS AND VEGETABLES.

- This product may be used with a nonionic surfactant or a wetting agent which is added to the water.
- Do NOT allow product to contact any COPPER

CROP	Application Timing	RATE fl oz/acre	Restrictions
Apple Pear	3-21 Days Before Anticipated Harvest (Fruit Ripening)	48 to 242 fl oz (60-300 g ai/ha)	Do not apply more than 242 fl oz /acre per crop. Allow a minimum of 3 days between treatment and harvest

(continued)

CROP	Application Timing	RATE fl oz/acre	Restrictions
Blueberry	3-21 Days Before Anticipated Harvest (Fruit Ripening)	48 to 242 fl oz (60-300 g ai/ha)	Do not apply more than 242 fl oz /acre per crop. Allow a minimum of 3 days between treatment and harvest
Kiwifruit	3-21 Days Before Anticipated Harvest (Fruit Ripening)	12 to 48 fl oz (15-60 g ai/ha)	Do not apply more than 48 fl oz /acre per crop. Allow a minimum of 3 days between treatment and harvest
BRASSICA LEAFY VEGETABLES (Crop Group 5):[*] Broccoli[*]; Broccoli, Chinese[*]; Broccoli raab[*]; Brussels sprouts[*]; Cabbage[*]; Cabbage, Chinese (bok choy) [*]; Cabbage, Chinese (napa) [*]; Cabbage, Chinese mustard[*]; Cauliflower[*]; Cavalo broccoli[*]; Collards[*]; Kale[*]; Kohlrabi[*]; Mustard Greens[*]; Mustard Spinach[*]; Rape Greens[*].	3-14 days before anticipated harvest	12 to 242 fl oz (15-300 g ai/ha)	Do not apply more than 242 fl oz /acre per crop. Allow a minimum of 0 days between treatment and harvest.
CUCURBIT VEGETABLES (Crop Group 9) [*]: Melon[*]: Cantaloupe[*], Citron melon[*], Honeydew[*], Muskmelon[*] and Watermelon[*]; Cucumber[*]; Gherkin[*]; Summer Squash[*]; cultivars, varieties, and/or hybrids of these[*].	3-21 days before anticipated harvest (Fruit Ripening)	121 to 242 fl oz (150-300 gai/ha)	Do not apply more than 242 fl oz /acre per crop. Allow a minimum of 3 days between treatment and harvest

(continued)

CROP	Application Timing	RATE fl oz/acre	Restrictions
Persimmon[*].	3-21 days before anticipated harvest (Fruit Ripening)	121 to 242 fl oz (150-300 gai/ha)	Do not apply more than 242 fl oz / acre per crop. Allow a minimum of 3 days between treatment and harvest.
STONE FRUIT (Crop Group 12-12A) [*]: Capulin[*]; Cherry, black[*]; Cherry, Nanking[*]; Cherry, sweet[*]; Cherry, tart[*]; cultivars, varieties, and/or hybrids of these.[*]	Single application - apply from 10% full bloom to 50% full bloom Two applications - apply first application at about 10% full bloom; apply second application at about 50% full bloom. For applying the second application using ground equipment, dilute nurse tank with Magnesium sulfate is recommended before applying AF-701	121 to 242 fl oz (150-300 gai/ha)	Do not apply more than 242 fl oz / acre per crop. Allow a minimum of 3 days between treatment and harvest
STONE FRUIT (Crop Group 12-12B) [*]: Peach[*]; Nectarine[*] STONE FRUIT (Crop Group 12-12C) [*]: Apricot[*]; Apricot, Japanese[*]; Jujube, Chinese[*]; Plum[*]; Plum, American[*]; Plum, beach[*]; Plum, Canada[*]; Plum, cherry[*]; Plum, Chickasaw[*]; Plum, Damson[*]; Plum, Japanese[*]; Plum, Klamath[*]; Plumcot[*];	3-21 days before anticipated harvest (Fruit Ripening)	121 to 242 fl oz (150-300 gai/ha)	Do not apply more than 242 fl oz / acre per crop. Allow a minimum of 3 days between treatment and harvest

(continued)

CROP	Application Timing	RATE fl oz/acre	Restrictions
Plum, prune[*]; Sloe[*]; cultivars, varieties, and/or hybrids of these.*]			
Table grapes [*]	Apply before grape veraison 3-21 days before anticipated harvest (Fruit Ripening)	12 to 48 fl oz (15-60 gai/ha)	Do not apply more than 48 fl oz / acre per crop. Allow a minimum of 3 days between treatment and harvest

[Not for use in California]

APPLES: HARVISTA 1.3 SC decreases the rate of fruit maturation, which allows for an extended optimal quality harvest period of 7 to 14 days. Depending on the cultivar, orchard conditions, and grower objectives, one or more of the following benefits may be provided by this delay: improved harvest management, reduction of preharvest fruit drop, reduced fruit ethylene production, additional time for color and size development, maintenance of fruit firmness and acid levels, delayed starch hydrolysis, delayed onset and incidence of watercore, and enhanced storage potential.

PEARS: HARVISTA 1.3 SC provides one or more of the following benefits depending on the cultivar, orchard conditions, and grower objectives: improved harvest management, reduction of preharvest fruit drop, reduced ethylene production, additional time for increased fruit size, maintenance of fruit firmness and enhanced storage firmness.

CHERRIES: HARVISTA 1.3 SC enhances fruit set.

BLUEBERRY: HARVISTA 1.3 SC provides one or more of the following benefits depending on the cultivar, orchard conditions, and grower objectives: improved harvest management, reduction of preharvest fruit drop, reduced ethylene production, additional time for increased fruit size, maintenance of fruit firmness and enhanced storage firmness.

KIWIFRUIT: HARVISTA 1.3 SC provides one or more of the following benefits depending on the cultivar, orchard conditions and grower objective: improved harvest management, increased fruit size maintenance of fruit firmness before harvest, reduction of preharvest fruit drop, maintenance of fruit quality, and enhanced storage potential.

BRASSICA LEAFY VEGETABLES (Crop Group 5) [*]: AF-701 provides one or more of the following benefits depending on the cultivar, field conditions and grower objectives: improved harvest management, maintenance of floret color before harvest, maintenance of floret quality and enhanced storage potential.

CUCURBIT VEGETABLES (Crop Group 9) [*]: AF-701 provides one or more of the following benefits depending on the cultivar, orchard conditions and grower objectives: Improved harvest management, increased fruit size, maintenance of fruit firmness before harvest, maintenance of fruit quality and enhanced storage potential.

PERSIMMON[*]: AF-701 provides one or more of the following benefits depending on the cultivar, orchard conditions and grower objectives: Improved harvest management, increased fruit size, maintenance of fruit firmness before harvest, maintenance of fruit quality and enhanced storage potential.

STONE FRUIT (Crop Group 12-12B)*]: AF-701 provides one or more of the following benefits depending on the cultivar, orchard conditions and grower objectives : Improved harvest management, increased fruit size, maintenance of fruit firmness before harvest, maintenance of fruit quality and enhanced storage potential. CHERRIES*]: AF-701 enhances fruit set.

TABLE GRAPES*]: AF-701 provides one or more of the following benefits depending on the cultivar, orchard conditions and grower objectives: improve harvest management, increased soluble solids, maintenance of fruit firmness before harvest, maintenance of fruit quality and enhanced storage potential.

GROUND APPLICATION

Apply the correct amount of HARVISTA 1.3 SC from the injector nurse tank. For best results, add an organosilicone surfactant as a wetting agent to the water in the main sprayer tank and use a minimum of 2 gallons of water per acre using an in-line injector system. See application timing by crop in the rate table. Follow manufacturer's recommendation for the proper concentration of wetting agent to be used. DO NOT apply when conditions favor drift from the target area. Flat-fan nozzles are preferred. Select nozzles and pressure combinations which deliver moderate (50-300 um range) spray droplets.

AERIAL APPLICATION

Apply the correct amount of HARVISTA 1.3 SC from the injector nurse tank. For best results, add an organosilicone surfactant as a wetting agent to the water in the main sprayer tank and use a minimum of 5 gallons of water per acre via helicopter or airplane equipped with an in-line injector system. See application timing by crop in the rate table. Follow manufacturer's recommendation for the proper concentration of wetting agent to be used. DO NOT apply when conditions favor drift from the target area. Flat-fan nozzles are preferred. Select nozzles and pressure combinations which deliver coarse (250-375 um range) spray droplets. Select nozzles and pressure combinations that maximize canopy penetration.

Restrictions/Precautions:

- Spray at a height of fifteen (15) feet or less above the tree canopy unless a greater height is required for safety.
- Nozzles must always point down, perpendicular to the ground.

Chemical Injector Systems

In-line injector systems are specifically designed for use with liquid formulations. This equipment can improve spray efficiency, reduce chemical waste, and facilitate easier clean out. Be sure injector nurse tank is drained and free of all water. Set injector to deliver the recommended amount of HARVISTA 1.3 SC per acre. Follow the manufacturer's recommendation for proper setting and calibration of the injector.

IMPORTANT CONSIDERATIONS FOR USE OF HARVISTA 1.3 SC

- Do not apply when temperatures exceed 95 degrees.
- Do not use overhead mist cooling or irrigation equipment for at least 1 hour following application.
- Do not apply if rain is expected within 1 hour of application.

- Combination effects and potential interactions between HARVISTA 1.3 SC and products containing naphthalene acetic acid (NAA) or ethephon have not been thoroughly evaluated. Use of these products on crops previously treated with HARVISTA 1.3 SC may reduce the benefits provided by HARVISTA 1.3 SC at harvest or after storage.
- Do not apply in winds exceeding 10mph
- Do not allow product to contact any copper or mix with products containing copper.
- Test this product in order to determine its suitability prior to use.
- HARVISTA 1.3 SC has a 3 day preharvest interval (PHI) for apple, pear, blueberry, kiwifruit, curcubit vegetable, persimmon, stone fruit, and table grape. • AF-7-01 has a 0 day preharvest interval (PHI) for brassica leaf vegetables.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original packaging in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill container smaller than 2 (two) gallons. Offer for recycling, if available.

For non refillable container, triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with concentrated salt water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure a second time with salt water and a third time with water.

For containers greater than 50 lbs: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with concentrated salt water. Replace and tighten closures. Tip container on its side and roll it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure a second time with salt water and a third time with water.

Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by the State and local ordinances.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of AGROFRESH INC. or the seller is authorized to vary in any way. Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of AGROFRESH INC. and the seller. The buyer or user of this product assumes all such inherent risks. Subject to the foregoing inherent risks, AGROFRESH INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD "AS IS" TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. AGROFRESH INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE. IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT AGROFRESH INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS, TO THE EXTENT REQUIRED BY APPLICABLE LAW:

AGROFRESH INC.

ATTENTION: LAW DEPARTMENT
510-530 WALNUT STREET – SUITE 1350
PHILADELPHIA, PA 19106

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, OR STRICT LIABILITY.

Harvista is a registered trademark of AgroFresh Inc.

NOTES

NOTES

NOTES

Harvista™ 1.3 SC

Protects fruit and vegetable crops from the various effects of ethylene.

ACTIVE INGREDIENT:	% By WT.
1-Methylcyclopropene	1.3%
OTHER INGREDIENTS:	98.7%
TOTAL:	100.0%

Produced for: AgroFresh Inc.
510-530 Walnut Street, Suite 1350
Philadelphia PA 19106, USA
Tel: 1-866-206-1001

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unused within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original packaging in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill container smaller than 2 (two) gallons. Offer for recycling, if available. For non refillable container, triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with concentrated salt water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure a second time with salt water and a third time with water.

Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by the State and local ordinances.

See [front][back][side][inside] panel for First Aid Instructions and [Leaflet][Booklet] for complete Precautionary Statements and Directions for Use.

FIRST AID	
If in eyes	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> ▪ Move person to fresh air. ▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. ▪ Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advice.
POISON CONTROL CENTER 1-800-222-1222	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378. For emergencies, call the poison control center 1-800-222-1222.	

Lot No. on container
EPA Reg. No. 71297-17
EPA Est. No. 71297-WA-1

Net Contents (on container):
Net content on package

110197_20220009US