

- For initial treatments, apply daily for 2 or 3 days. As infestations decline, repeat the application at 14 to 21 day intervals if needed, if infestation persists, or if prescribed by veterinarian. The duration of activity may vary because of environmental conditions (e.g. rainfall) and management factors (husbandry, bathing or shampooing).
- Do not wet the animal's skin.
- Do not apply to areas where animal has no hair. Avoid direct contact with the animal's skin.
- Securely cover feedstuffs, troughs, bins, or other food and water receptacles while spraying.

Table 1. Application Instructions of Horse and Pony Spray for direct applications as mist or wipe-On application to horses and ponies.

| Treatment Method | Pest Controlled | Application rate, fl.oz. of product per adult animal | Retreatment interval, days | Instructions |
|--|---------------------------|--|----------------------------|---|
| Ready-To-Use Use undiluted as a localized or spot treatment. | Horse flies or deer flies | 3.25 to 6.5 fl.oz. per application | 14 to 21 days | Apply to head, neck, shoulders, flanks, legs, belly and croup (rump). |
| | Stable fly | 3.25 to 6.5 fl.oz. per application | 14 to 21 days | Stable flies bite chiefly about the legs. Apply to legs, ankles and belly. |
| | Horn fly | 3.25 to 6.5 fl.oz. per application | 14 to 21 days | Horn flies hover over the backs, withers, and belly. Apply to withers, flanks and belly. |
| | Gnats | 3.25 to 6.5 fl.oz. per application | 14 to 21 days | Gnats hover about the eyes, ears, and nostrils. Treat face, ears, poll, jaw, and neck. Avoid contact with eyes and muzzle (mouth and nostrils). |
| | Face fly | 3.25 to 6.5 fl.oz. per application | 14 to 21 days | Face flies cluster on the head. Treat face, ears, poll, jaws and neck. Avoid contact with eyes and muzzle. |

Whole Body Spray:

- Dilute concentrate with water for use as a whole body treatment for mosquitoes, ticks, and horse sucking louse (lice).
- Measure the proper amount of product, pour the product into the sprayer, and add the appropriate amount of water.
- Use a 1:4 ratio with water by mixing 1 gallon of product with 4 gallons of water (one gallon of product provides five gallons of spray solution). Apply 1 to 2 pints of spray emulsion per animal.
- Treat animal thoroughly or cover the entire surface area of the animal (See Table 2). Avoid contact with eyes and muzzle (mouth and nostrils).
- Repeat the application at 14 to 21 day intervals if needed, if infestation persists, or if prescribed by veterinarian.
- Securely cover feedstuffs, troughs, bins, or other food and water receptacles while spraying.

Table 2. Application Instructions of Horse and Pony Spray for direct applications as whole body sprays to horses and ponies.

| Treatment Method | Pest Controlled | Application rate, fl.oz. of product per adult animal | Retreatment interval, days | Instructions |
|---|----------------------------|--|----------------------------|---|
| Concentrate Diluted with water as whole body spray. | Mosquitoes | Use 1:4 dilution | 14 to 21 days | Treat entire animal or apply for complete coverage. |
| | Horse sucking louse (lice) | Use 1:4 dilution | 14 to 21 days | Lice are most common about the head, neck, and base of the tail. Even if few lice are found, treat entire body. |
| | Ticks | Use 1:4 dilution | 14 to 21 days | Treat the animal thoroughly with 1 to 2 pints of spray applied per animal. |

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. To prevent cross-contamination during storage, keep separate from other pesticides, fertilizers, seeds, food, and feed.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS OF A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Gordon's® logo is a registered trademark of PBI-Gordon Corporation.

967/11-2018 AP080417
EPA REG. NO. 2217-479



MANUFACTURED BY
PBI/GORDON CORPORATION
P.O. BOX 860350
SHAWNEE, KANSAS 66286
www.GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.