# **Z-MAX**°

## **GUARANTEED ANALYSIS**

Sulfur (S)	5.00%
5.00% Combined Sulfur (S)	
Copper (Cu)	0.50%
Manganese (Mn)	2.00%
Zinc (Zn)	8.00%

Derived from: Copper Sulfate, Manganese Sulfate, Zinc Sulfate.

#### **DIRECTIONS FOR USE**

• Shake well before use. • Designed to be applied by both soil and/or foliar application. • Z-MAX® should be applied when tissue and soil analysis identifies the need and proper rate of application for this product. • Best results will be obtained when application is concentrated in the active root zone or on the leaf surface. • Can be applied in combination with compatible plant growth regulators, pesticides, or other liquid fertilizers. • If compatibility is in question, jar test a small quantity. • Apply foliar sprays with sufficient water to ensure uniform coverage without running off leaf surfaces. • Do not apply this product foliarly in concentrations greater than 5%. • For maximum results, apply in the early morning or in the evening when air temperatures are optimum for foliar application. • Do not apply HUMA GRO® nutrients during the hottest time of the day or to plants under drought or flood stress. • Application timing, intervals and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop. • Applications can be made as often as every 7 to 10 days as needed. • Consult your local HUMA GRO® Representative or other agricultural specialist for crop-specific recommendations. • Do not store this product below 50°F (10°C) or above 90°F (30°C).

This product is intended as a supplement to a complete fertilization program and will not, by itself, provide all the nutrients normally required by agricultural crops.

#### CONDITIONS OF SALE:

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions and other factors are beyond the control of the seller.

CAUTION: KEEP OUT OF REACH OF CHILDREN

### METHOD OF APPLICATION

SUGGESTED RATE PER ACRE

Foliar band application at 50% coverage

up to 1 quart / - -

Foliar broadcast or sprinklers: solid, set, pivot, linear (100% speed)

up to 2 quarts / up to 1 gallon

Soil banded or injected through drip tape or micro sprinklers

up to 2 quarts / up to 1 gallon

Soil broadcast spray incorporated, flood or furrow irrigated

up to 1 gallon / up to 2 gallons

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

A clear to slightly hazy, greenish blue, moderately acidic liquid having a slight characteristic odor. The liquid and mists may cause irritation to the eyes, skin and respiratory tract. This product may be toxic by ingestion or inhalation of high mist concentrations.



CLASSIFICATION: HAZARD CATEGORY 4

SIGNAL WORD: WARNING
HAZARD STATEMENT: H302 Harmful if swallowed.
PRECAUTIONARY STATEMENT: P301+P312 IF SWALLOWED, call a POISON
CENTER or doctor/physician if you feel unwell: P330 Rinse mouth;
P264 Wash hands thoroughly after handling: P270 Do not eat, drink or smoke when using this product.

CLASSIFICATION: MILD SKIN IRRITATION – HAZARD CATEGORY 3 SIGNAL WORD: WARNING

HAZARD STATEMENT: H316 - WARNING - Causes mild skin irritation PRECAUTIONARY STATEMENT: P332+P313 If skin irritation occurs, get medical attention/advice.

**11.70 lb per gallon @ 68° F.** 1.40 kg per liter @ 20° C. pH 2.5 - 3.5 1 liter • Net Wt. 3.09 lb 1 liter • Net Wt. 1.40 kg LOT# 12345678



Produced by

BIO HUMA NETICS, INC. © 2005-2016

1331 W. Houston Ave., Gilbert, AZ 85233 • 480.961.122

HG-280-151208