

Imazethapyr 2 SL

GROUP 2 HERBICIDE

Herbicide For use on Alfalfa, Clover, Peas, Beans,
Peanuts and Soybeans

ACTIVE INGREDIENT: % BY WT.

Ammonium salt of imazethapyr (\pm)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid*22.87%

OTHER INGREDIENTS: 77.13%

TOTAL: 100.0%

*Equivalent to 21.6% or 2 pounds per U.S. gallon or 240 grams per liter of imazethapyr acid.

EPA Reg. No. 66222-248

**EPA Est. No. 37429-GA-001^{BT}; 37429-GA-002^{BO};
37429-GA-003^{BV}**

Letter(s) in lot number correspond(s) to superscript
in EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For First Aid, Precautionary Statements, and Directions for Use,
see inside of this booklet.

How can we help? 1-866-406-6262

Net Contents
1 gallon



HERBICIDE

ADAMA
ESSENTIALS

FIRST AID

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.

In case of spills, fire, leaks or accidents call 1-800-535-5053.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemically resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves such as butyl rubber ≥ 14 mils, or natural rubber ≥ 14 mils, or neoprene rubber ≥ 14 mils, or nitrile rubber ≥ 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, and chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Groundwater Advisory and Proper Handling Instructions

This chemical has properties and characteristics associated with chemicals detected in groundwater. This use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes or reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum, 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Do not apply this product through any type of irrigation system.

This product must be used in a manner that will prevent back siphoning in wells, spills or improper disposal of excess pesticide spray mixture.

PHYSICAL AND CHEMICAL HAZARDS

Do not use with or store near oxidizing agents.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves such as butyl rubber ≥ 14 mils, or natural rubber ≥ 14 mils, or neoprene rubber ≥ 14 mils, or nitrile rubber ≥ 14 mils
- Shoes plus socks

PRODUCT INFORMATION

Do not use or sale this product in Long Island, New York State.

Label directions must be with the applicator when treatment takes place, and must be read and followed in full. Application of Imazethapyr 2 SL in any way that is not in accordance with these directions may cause crop injury. Any such injury is the responsibility of the applicator.

Treatment with Imazethapyr 2 SL will provide residual control of listed germinating target species when there is sufficient moisture.

Crop Growth Following Treatment

Normal growth of rotational crops should take place following applications of Imazethapyr 2 SL. However, it is impossible to anticipate and eliminate all risk factors brought about by varying environmental and agronomic conditions. Rotational crop injury therefore may result from treatment with Imazethapyr 2 SL.

A combination of treatment with this product and certain conditions, such as high organic matter in the soil, low soil pH, heavy soil texture or low rainfall, may cause damage to crops that are subsequently planted.

Mode of Action

Treatment with Imazethapyr 2 SL provides control through the AHAS/ALS enzyme inhibiting mode of action. Application of Imazethapyr 2 SL works by uptake of the treatment by target species through foliage and/or roots and then translocates quickly to the growing points. For optimal mode of action, soil must be moist prior to application.

Applications of Imazethapyr 2 SL may cause internode shortening and/or yellowing of desirable vegetation. These effects, when they occur, are temporary and normal growth should resume 1-2 weeks following treatment. Sugar beets and other vegetable crops are susceptible to residues of Imazethapyr 2 SL in the soil.

RESISTANCE MANAGEMENT

Imazethapyr 240SL is a Group 2 Herbicide (contains the active ingredient Imazethapyr). Following many years of continuous use of this product and chemically related products biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected and weeds controlled by this product are expected to be present along with resistant biotypes we recommend the use of this product in combinations or in sequence with other registered herbicides which are not solely a Group 2 Herbicide. If only resistant biotypes are expected to be present use a registered herbicide which is not solely a Group 2 Herbicide Consult with your state Agricultural Extension Service for specific recommendations.

Weed Resistance

Some listed weeds have developed naturally occurring biotypes which will not be controlled by applications of Imazethapyr 2 SL or other products that have a similar mode of action, such as sulfonyleureas, sulfonamides and pyrimidyl benzoates.

Where naturally resistant biotypes occur, control can be achieved by sequentially applying or tank mixing this product with a registered product with a different mode of action.

Replanting

When replanting an area previously treated with Imazethapyr 2 SL, the following crops may be planted:

	lima beans	peanuts
Southern peas	soybeans	

Restrictions

- Do not rework the soil any deeper than the treated zone.
- Do not apply this product a second time.

Refer to individual crop sections and the Rotational Crop section for minimum replanting intervals following treatment.

USE DIRECTIONS

Containers containing Imazethapyr 2 SL must be closed securely in order to prevent contamination and spills. Application equipment must be drained and cleaned thoroughly prior to mixing the application solution and treatment. Application equipment must also be drained and thoroughly cleaned following treatment to avoid contamination and future crop injury.

Adjuvants

When applying Imazethapyr 2 SL as a postemergence treatment, it must be combined with a fertilizer solution (see Fertilizer Solution table below) and **one** of the following adjuvants:

Adjuvant	Directions
Crop Oil Concentrate ¹ (vegetable or petroleum based)	When target species are under stress from temperature or moisture, use methylated seed oil at the rate of 1.0% v/v (i.e. 1 gallon of methylated seed oil per 100 gallons of spray application solution)
	Use crop oil concentrate at a rate of 1.25% v/v (i.e. 1.25 gallons of crop oil concentrate per 100 gallons of spray application solution)

Adjuvant	Directions
Surfactant	Combine Imazethapyr 2 SL with a non-ionic surfactant with a minimum of 80% active ingredient, at the rate of 0.25% v/v (i.e. 0.25 gallons (1 quart) of surfactant per 100 gallons of spray application solution). A dry surfactant or organo-silicone surfactant may be combined with Imazethapyr 2 SL instead of a non-ionic surfactant.

¹Do not apply Imazethapyr 2 SL with a crop oil concentrate when treating edible legume vegetables.

Fertilizer Solution^{1,2}

Combine Imazethapyr 2 SL with a nitrogen based liquid fertilizer (for example: 28%N, 32%N, 10-34-0). Add 1.25-2.5 gallons of fertilizer solution per 100 gallons of application spray solution. When target species are under stress from temperature and/or moisture, use the higher rate of fertilizer in the specified rate range.

As an alternative to a fertilizer solution, a spray grade ammonium sulfate may be combined with Imazethapyr 2 SL at a rate of 12-15 pounds per 100 gallons of spray application solution

¹Do not use fertilizer solution in California

²Fertilizer solution is not required for applications south of Interstate Highway 40, except in New Mexico, Oklahoma and Texas.

Restriction

- Do not use fertilizer solution in California.

Mixing Procedure

1. Fill mix tank half full with clean water.
2. Add the specified amount of Imazethapyr 2 SL while agitating the solution.
3. Add specified adjuvants while continuing agitation.
4. Fill the remaining volume with clean water.

Tank Mixtures

Imazethapyr 2 SL may be tank mixed with registered organo-phosphate or carbamate insecticide products. When applied in crops, temporary crop damage may result. Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Tank Mix Procedure

1. Fill tank with clean water.
 2. Combine soluble packet products whilst agitating and thoroughly mix.
 3. Add ingredients not in soluble packets:
 - Dispersible granules (DG)
 - Dry flowables (DF)
 - Wettable powders (WP)
 - Liquid flowables
- Continue agitation and mix thoroughly.

4. Add Imazethapyr 2 SL and other aqueous solution product(s). Continue agitation and mix thoroughly.
5. Add emulsifiable concentrate (EC) products. Continue agitation and mix thoroughly.
6. Add crop oil or surfactant as appropriate (see Adjuvants section above). Continue agitation and mix thoroughly.
7. Add fertilizer solution as appropriate (see Adjuvants section above). Continue agitation and mix thoroughly.
8. Fill the remaining tank volume with clean water. Continue agitation and mix thoroughly.

SPRAY APPLICATIONS

Restriction

- Do not apply Imazethapyr 2 SL when wind speed exceeds 10 mph or when spray may drift to sensitive crops (e.g. sugar beets and leafy vegetables).

Ground Applications

Apply Imazethapyr 2 SL in a minimum of 10 gallons of water per acre. Apply solution at 20-40 psi, at a sufficient boom height to ensure uniform coverage of target species foliage.

When applying Imazethapyr 2 SL to no-till crop areas, use a minimum of 20 gallons of water per acre for sufficient coverage of target species. Apply Imazethapyr 2 SL in higher volume where there is dense crop residue and/or target species foliage.

Restrictions

- For postemergence applications, only use flat fan nozzles.
- Do not overlap spray applications.

Low Volume Spray Application

Use low volume spray application equipment to apply Imazethapyr 2 SL to soybeans. Treat target species before they reached the specified height. For an effective treatment, calibrate application equipment to ensure spray coverage is sufficient and uniform.

Apply Imazethapyr 2 SL in a minimum of 10 gallons of water per acre at a pressure of 40-60 psi.

Aerial Application

Unless otherwise directed, Imazethapyr 2 SL may be applied by air in a minimum of 5 gallons of water per acre.

For optimal effectiveness when applying Imazethapyr 2 SL as a postemergence treatment, add a fertilizer solution and a non-ionic surfactant or crop oil concentrate to the application solution.

Non Ionic Surfactant	Combine 1 quart per 100 gallons of application solution.
Crop Oil Concentrate	Combine 1.25 gallons per 100 gallons of application solution.
Liquid Fertilizer¹	Combine 1.25 gallons per 100 gallons of application solution.
¹ Do not use fertilizer solution in California	

Follow drift management directions in order to avoid contact with and damage to crops.

Note: drift management directions do not apply to dry formulation applications, public health uses or forestry treatments.

Restrictions

- Avoid contact with non-target species through drift or otherwise. Applicators are responsible for assessing application conditions and equipment in order to avoid drift.
- On the boom, the distance of the outer most nozzles must not exceed 75% the length of the wingspan or rotor.
- Spray nozzles must always be parallel with the air stream and must point backwards.
- Spray nozzles must not be pointed downwards more than 45 degrees.

More restrictive directions imposed by states must be followed.

Aerial Drift Reduction Advisory Information:

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets. Increase water volume to at least 10 gallons of water per acre if grass foliage or crop canopy is dense.
- Pressure - Do not exceed the nozzle manufacturer's specific pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the specified practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than 75% of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

The applicator is responsible for any loss or damage that results from spraying this product in a manner other than instructed in this label. In addition, the applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

APPLICATION INSTRUCTIONS

POSTEMERGENCE APPLICATIONS

When applied as a postemergence treatment, Imazethapyr 2 SL will provide control of existing grasses and broadleaf weeds in conventional production systems as well as conservation tillage areas. Target species absorb the treatment through foliage and roots and will either die or stop growing. Imazethapyr 2 SL will also provide control of target species that emerge following application.

Temperatures under 50°F will reduce the efficacy of an application of Imazethapyr 2 SL. For optimal effectiveness and to minimize the risk of crop damage, delay application for 48 hours after the temperature has reached 50°F for a minimum of 10 hours.

Apply this product during early postemergence when target species are less than 3 inches in height and actively growing, and a minimum of one hour prior to irrigation/rainfall. Treat target species when most of them have reached the specified growth stage.

For optimal effectiveness, combine Imazethapyr 2 SL with a crop oil concentrate or a surfactant and a nitrogen-based fertilizer. Refer to the Adjuvants section above.

For optimal control of target species, carry out tillage 7-10 after application. Tillage will improve residual control of target species, in particular when conditions are dry.

DOUBLE CROP SOYBEANS NO TILLAGE/MINIMUM TILLAGE SYSTEMS

Applied during early postemergence, treatment with Imazethapyr 2 SL will control existing weeds and provide residual control of most weed species in the following crops:

- Soybeans in no tillage/minimum tillage systems
- Soybeans in double crop production systems

Apply Imazethapyr 2 SL before or after crop emergence in accordance with the specified weed height at the time of application.

Where weeds are greater than the specified size and Imazethapyr 2 SL is applied before crop emergence, combine the application solution with a contact herbicide in order to improve control (refer to the Preemergence Applications section).

SOIL APPLICATIONS

Target species absorb the treatment through foliage and roots and will either die or stop growing. Imazethapyr 2 SL will also provide control of target species that emerge following application.

Apply Imazethapyr 2 SL in conservation tillage areas designed to meet conservation compliance requirements in order to achieve control of listed species.

Imazethapyr 2 SL can be applied in soybeans as a preplant incorporated, early preplant, or preemergence treatment. Imazethapyr 2 SL may also be used to treat no-tillage, minimum tillage and conventional systems.

For optimal effectiveness, rainfall/irrigation is required to ensure that there is soil moisture (to a depth of 2 inches) so that treatment with Imazethapyr 2 SL can move into the weed germination zone. The required level of rainfall irrigation will depend on existing soil moisture, organic content and soil texture.

If the moisture level remains insufficient 7 days after treatment, carry out cultivation in order to control emerging weeds. As the moisture content returns to adequate levels, Imazethapyr 2 SL will provide residual control of listed germinating species. The location of the root system in the soil and species type will determine the effectiveness of application on established weeds.

Soil Applications With Liquid Fertilizers

Imazethapyr 2 SL can be applied in liquid fertilizers to soybeans on its own or in combination with Outlook® or Prowl® 3.3 EC. Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Carry out a jar mixture compatibility test on the proposed mixture/combination before mixing on a larger scale prior to application.

Apply treatment solution with ground equipment in a minimum of 20 gallons of liquid fertilizer per acre.

Restriction

- Any tank mixture containing trifluralin must only be applied to soybeans.

PREEMERGENCE (SURFACE APPLICATIONS)

Apply Imazethapyr 2 SL in production tillage systems up to 45 days before planting and at planting in no tillage, reduced tillage or in conventional systems. Imazethapyr 2 SL may also be applied as an after planting and preemergence treatment.

Reduced or No-Till Tillage Systems

Treat with Imazethapyr 2 SL prior to planting, during planting or after planting in a minimum of 20 gallons of water per acre. For optimal effectiveness, ensure coverage is thorough and uniform. Use a higher volume solution where weeds are dense or where there is dense crop residue.

For enhanced control of grasses, tank mix Imazethapyr 2 SL Prowl® 3.3 EC or Outlook®.

For optimal treatment of existing target species, tank mix Imazethapyr 2 SL with Gramoxone® Extra, Roundup Ultra® or a registered product containing 2,4-D (for application as an early pre-planting treatment - refer to 2,4-D label for restrictions). Product(s) can be tank mixed with Imazethapyr 2 SL alone or in combination with Prowl® 3.3 EC or Outlook®.

Restriction

- Do not tank mix Imazethapyr 2 SL with Gramoxone® Extra, Roundup Ultra® or a registered product containing 2,4-D if there are no existing target species at the time of treatment.

INCORPORATED APPLICATIONS – PREPLANT TREATMENT

Apply Imazethapyr 2 SL incorporated to a depth of 1-2 inches after land preparation and bed formation with a rolling cultivator (if crops planted on beds) or PTO-driven equipment or a rolling cultivator.

Keep Imazethapyr 2 SL at a depth of 1 to 2 inches in the finished beds.

When applying Imazethapyr 2 SL for control of nutsedge in peanuts, incorporate the product into the soil with two passes of the incorporation equipment. In order to minimize the risk of streaking, make the second pass at an offset angle to the first pass.

Restriction

- When treating soybeans, treatment must be applied a minimum of 45 days before planting.

FEDERAL CONSERVATION RESERVE PROGRAM AND AGRICULTURAL RESERVE PROGRAM LAND SEEDED TO FORAGE LEGUME SPECIES

Apply Imazethapyr 2 SL to control many annual grass and broadleaf weeds in Agricultural Reserve Programs (Set-Aside) and Conservation Reserve Programs, as well as areas seeded to forage legumes.

Treatment with Imazethapyr 2 SL may cause a reduction in growth of legumes, however these effects are temporary, and desirable crops will benefit from reduced competition from weed species.

Restrictions

- Do not allow grazing/feeding of treated legumes.
- Treated legumes must not be cut for use as forage/hay.
- Treated legumes must not be harvested for use as feed.
- Seed from treated legumes must not be used for sprouting.
- Do not exceed a maximum of 1 application of Imazethapyr 2 SL per year.

Cover Crops

Apply Imazethapyr 2 SL to cover crops, including:

Legumes	
Alfalfa	Lupin
Birdsfoot trefoil	Milk vetch
Clover	Sainfoin
Crown vetch	Trefoil
Kudzu	Velvet bean
Lespedeza	Vetch

Restriction

- Where cover crops have been planted into areas previously treated with Imazethapyr 2 SL for control of target species in soybeans, do not treat the cover crop until the following spring.

Conservation Reserve Program: Postemergence Applications

Apply Imazethapyr 2 SL to cover crops in Conservation Reserve Program areas at a rate of 4 fluid ounces per acre as a postemergence treatment to established legumes. Apply as a fall treatment or in the spring before they reach the specified height for control.

Legume seedlings that have 3 fully expanded trifoliate leaves (minimum) may also be treated with Imazethapyr 2 SL.

See the Weeds Controlled table in the Soybean section for information.

EDIBLE LEGUME VEGETABLES

Because crop maturity may be delayed by application of Imazethapyr 2 SL, crop harvest will also be delayed. Growth, crop quality and yield may also be affected.

In order to minimize risk of crop damage, lima beans, peas and lentils must be planted at 0.5 inches depth minimum.

Restrictions

- Do not apply Imazethapyr 2 SL as a postemergence treatment after flowering occurs in order to avoid crop damage.
- If planting is delayed, do not apply Imazethapyr 2 SL if frost is possible prior to crop maturity.
- Do not apply Imazethapyr 2 SL if there are cold/wet conditions prevailing or if they are forecast within 7 days of treatment.
- Imazethapyr 2 SL must only be applied in areas where good agronomic methods have been carried out, such as:
 - Crop rotation
 - Good soil fertility
 - Insect management
 - Disease management
 - Tillage to prevent hardpans and compaction

CROP USE DIRECTIONS

ALFALFA AND CLOVER

Application of Imazethapyr 2 SL will control a broad range of grass and broadleaf weeds in alfalfa and clover. Clover and alfalfa develop tolerance to applications of this product once it is at/beyond the 3 trifoliate growth stage. Yellowing or minor height reduction may occur following treatment, particularly where application is made at temperatures of 40°F or less.

Apply Imazethapyr 2 SL as a broadcast postemergence application to actively growing weeds in alfalfa and clover at the rate of 3-6 fluid ounces per acre. Apply before target species exceed 3 inches in height.

Target species suffering from stress (e.g., due to drought or extremes of temperature) are less susceptible to treatment.

Restrictions

- For application in alfalfa and clover as a postemergence application only.
- Do not exceed a maximum of 6 fluid ounces of Imazethapyr 2 SL per acre per year (0.094 lbs. of imazethapyr acid equivalent per acre per year).
- Do not exceed a maximum of 4 fluid ounces of Imazethapyr 2 SL per acre in North Dakota or Minnesota north of Highway 210.
- Do not exceed a maximum of 4 fluid ounces of Imazethapyr 2 SL per acre in the last year of the stand.
- Treated alfalfa or clover must not be grazed, fed or harvested for a minimum of 30 days following application.

Seedlings

Treatment with Imazethapyr 2 SL may cause a reduction in growth. This effect is temporary.

Apply Imazethapyr 2 SL to seedling alfalfa and clover as a post emergence application when most weeds are 1-3 inches in height and seedlings are at or beyond the second trifoliate stage

When treating mustards or other low growing target species, treat with Imazethapyr 2 SL before the rosette is greater than 3 inches.

Restriction

- For application in alfalfa and clover as a postemergence application only.
- Apply Imazethapyr 2 SL to alfalfa or clover grown for seed before bud formation.

Established Alfalfa/Clover

Apply Imazethapyr 2 SL in established dormant or semi-dormant clover or alfalfa before any significant growth/regrowth (i.e. less than 3 inches regrowth). Applications delayed until after significant regrowth has taken place may not allow treatment to reach target species. Make applications in the spring or in the fall, or treat between cuttings.

Replanting Restriction

- Do not plant clover or alfalfa for a minimum of 4 months following treatment with Imazethapyr 2 SL.

See the Rotational Crop section for more information.

Weeds Controlled (Postemergence Application)

For optimal suppression/control, apply Imazethapyr 2 SL before target species reach the specified height below.

Weeds Controlled: Broadleaf Weeds

Weed	App. Rate (fl. oz./acre)	Max. Size (inches)	Weed	App. Rate (fl. oz./acre)	Max. Size (inches)	Weed	App. Rate (fl. oz./acre)	Max. Size (inches)	Weed	App. Rate (fl. oz./acre)	Max. Size (inches)
Artichoke, Jerusalem	3	*	Beets, Wild	3	4 in.	Bedstraw, Catchweed	3		Buckwheat, Wild	3	
	4	6 in.		4	5 in.		4	3 in.		4	3 in.
	6	8 in.		6	6 in.		6	4 in.		6	4 in.
Chickweed, Common	3	*	Chickweed, mouseear	3	*	Cocklebur, common	3	*	Cress, Hoary	3	
	4	3 in.		4	3 in.		4	8 in.		4	*
	6	4 in.		6	3 in.		6	8 in.		6	*
Dandelion	3		Dock, Broadleaf (seedling)	3		Dock, Curly (seedling)	3		Dodder†	3	
	4	*		4			4			4	
	6	5*		6	6*		6	6*		6	*
Fiddleneck	3		Filaree, Redstem	3		Filaree, Whitestem	3		Fleabane, Rough	3	
	4			4	*		4	*		4	3 in.
	6	4*		6	3 in.		6	3 in.		6	3 in.
Fixweed	3	*	Goosefoot, Nettleleaf	3	*	Groundsel, Common	3		Henbit	3	
	4	3 in.		4	3 in.		4			4	*
	6	4 in.		6	4 in.		6	3*		6	3 in.
Jimsonweed	3		Knotweed, Prostrate	3		Kochia, (non-ALS resistant)	3	*	Lambs quarter, (common) †	3	
	4	3 in.		4	*		4	3 in.		4	*
	6	4 in.		6	4 in.		6	3 in.		6	2*
Lettuce, Minors	3		Mallow, Common	3		Marshelder	3		Morning-glory, Entireleaf	3	
	4	3 in.		4	3 in.		4	4 in.		4	*
	6	4 in.		6	3 in.		6	6 in.		6	3 in.
Morning-glory, Ivyleaf	3		Morning-glory, Pitted	3		Morning-glory, Small-flower	3	*	Morning-glory, Tall	3	
	4	*		4	*		4	3 in.		4	*
	6	3 in.		6	3 in.		6	4 in.		6	3 in.
Mustard, Tumble	3	3 in.	Mustard, Wild	3	3 in.	Mustard, Black	3	3 in.	Nettle, Burning	3	
	4	3 in.		4	3 in.		4	3 in.		4	3 in.
	6	4 in.		6	4 in.		6	4 in.		6	4 in.
Nightshade, Black	3	3 in.	Nightshade, Eastern Black	3	3 in.	Nightshade, Hairy	3	3 in.	Oxtongue, Bristly	3	
	4	3 in.		4	3 in.		4	3 in.		4	
	6	4 in.		6	4 in.		6	4 in.		6	3*

Weed	App. Rate (fl. oz./acre)	Max. Size (inches)	Weed	App. Rate (fl. oz./acre)	Max. Size (inches)	Weed	App. Rate (fl. oz./acre)	Max. Size (inches)	Weed	App. Rate (fl. oz./acre)	Max. Size (inches)
Pennycress, Filled	3	3 in.	Pep-perweed, Field	3	3 in.	Pep-perweed, Virginia	3	*	Pigweed, redroot	3	4 in.
	4	3 in.		4	3 in.		4	3 in.		4	6 in.
	6	4 in.		6	4 in.		6	3 in.		6	8 in.
Pigweed, Smooth	3	4 in.	Pigweed, Spiny	3		Radish, Wild	3		Ragweed, Common	3	
	4	6 in.		4	6 in.		4	*		4	2 in.
	6	8 in.		6	8 in.		6	4 in.		6	3 in.
Ragweed, Giant	3		Redmaids	3		Rocket, London	3	3 in.	Rocket, Yellow	3	*
	4	3 in.		4	3 in.		4	4 in.		4	3 in.
	6	3 in.		6	4 in.		6	6 in.		6	4 in.
Rock-purslane, Desert	3		Shepherds-purse	3	3 in.	Smart-weed, Ladys-thumb	3	*	Smart-weed, Pennsylvania	3	*
	4			4	3 in.		4	3 in.		4	3 in.
	6	3 in.		6	4 in.		6	4 in.		6	4 in.
Smart-weed, Swamp (seedling)	3		Sprurge, Prostrate	3		Sprurge, Spotted	3		Sprurge, Petty	3	
	4	3 in.		4	*		4	*		4	3 in.
	6	4 in.		6	3 in.		6	3 in.		6	4 in.
Spurry, Corn	3		Sunflower, Common	3	*	Swinecress	3		Tansy-mustard, Green	3	3 in.
	4	3 in.		4	4 in.		4	3 in.		4	3 in.
	6	3 in.		6	6 in.		6	3 in.		6	4 in.
Tansy-mustard, Pinnate	3	3 in.	Thistle, Russian	3	*	Velvetleaf	3	*	Watercress, Creeping	3	
	4	3 in.		4	3 in.		4	3 in.		4	2 in.
	6	4 in.		6	3 in.		6	4 in.		6	3 in.
Watercress	3		Willow-weed, Panicle	3							
	4	3 in.		4	3 in.						
	6	3 in.		6	3 in.						

‡ For optimal effectiveness, combine Imazethapyr with a methylated seed oil or crop oil concentrate following dodder emergence but before/immediately after attachment. Imazethapyr 2 SL will provide control of many grass weed species. For optimal effectiveness under heavy pressure from grass weeds, apply Imazethapyr 2 SL sequentially with a grass herbicide (e.g. Poast Plus®)

† Apply at the 1-2 leaf stage of growth.

∞ Imazethapyr 2 SL will provide control of emerged woolly cupgrass only.

Ω Apply Imazethapyr 2 SL to quackgrass that is growing actively and less than 7 inches in height for suppression only.

Weeds Controlled: Grass Weeds and Sedges

Weed**	App. Rate (fl. oz./acre)	Max. Size (inches)	Weed**	App. Rate (fl. oz./acre)	Max. Size (inches)	Weed**	App. Rate (fl. oz./acre)	Max. Size (inches)	Weed**	App. Rate (fl. oz./acre)	Max. Size (inches)
Barnyard-grass	4	*	Bluegrass, Annual	4		Canary-grass, Littleseed	4	*	Cereal, volunteer barley	4	*
	6	3 in.		6	3*		6	3*		6	4*
Cereal, volunteer oats	4	*	Cereal, volunteer wheat	4	*	Crabgrass, Large	4	*	Crabgrass, Smooth	4	*
	6	4*		6	4*		6	3 in.		6	3 in.
Cupgrass, Woolly [∞]	4	3 in.	Foxtail, Giant	4	6 in.	Foxtail, Green	4	3 in.	Foxtail, Yellow	4	3 in.
	6	3 in.		6	6 in.		6	4 in.		6	3 in.
Johnson-grass, Seeding	4	8 in.	Johnson-grass, Rhizome	4	*	Millet, Wild Proso	4	*	Nutsedge, Yellow	4	*
	6	8 in.		6	6-12*		6	3 in.		6	6*
Nutsedge, Purple	4	*	Oats, Wild	4	*	Rice, Red	4	3 in.	Shatter-cane	4	8 in.
	6	6*		6	4*		6	4 in.		6	10 in.
Signal-grass, broadleaf	4	*	Quack-grass ^Ω	4							
	6	8 in.		6	7*						

* Suppression only

**Imazethapyr 2 SL will provide control of many grass weed species. For optimal effectiveness under heavy pressure from grass weeds, apply Imazethapyr 2 SL sequentially with a grass herbicide (e.g. Poast Plus[®])

Tank Mixtures

Imazethapyr 2 SL may be tank mixed with other registered products where control is required of target species not listed in this table. Imazethapyr 2 SL may be tank mixed with the following products:

Buctril [®]	2,4-DB	Poast [®]
Poast Plus [®]	Prism [®]	Select [®]

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Establishing Stands

Apply Imazethapyr 2 SL before target species reach the specified height in the Weeds Controlled table and after clover or alfalfa has 2 fully expanded trifoliate leaves. Treat spring, summer or fall seeded clover or alfalfa.

Clover or alfalfa may be inter-seeded with oats. This method will reduce soil erosion and assist in stand establishment. Treating clover or alfalfa inter-seeded with oats will greatly reduce the growth of oats, or will kill the oats, allowing clover/alfalfa to establish with reduced soil erosion. Treat oat inter-seeded clover/alfalfa with Imazethapyr 2 SL when the oats are at the 3-4 leaf stage of growth.

Established Alfalfa/Clover (Growing)

Apply Imazethapyr 2 SL to clover/alfalfa after cutting for control of listed weeds during the season. Following cutting, remove hay from the treatment area and treat excessive clover/alfalfa regrowth with this product.

Treatment with Imazethapyr 2 SL will also reduce competition from present perennial grasses such as bromes, fescues, orchardgrass, and timothy grass.

Established Alfalfa/Clover (Dormant)

Apply this product to dormant alfalfa or clover in the fall following the last cutting. Also apply this product in the spring to dormant alfalfa or clover, or as alfalfa or clover breaks dormancy. Apply spring treatments prior to excessive alfalfa or clover growth (less than 3 inches), to reduce spray interference.


Restriction

- Treated clover or alfalfa must not be grazed, fed or harvested for a minimum of 30 days following treatment with Imazethapyr 2 SL.

NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO, LIMA, AND SMALL WHITE TYPE DRY BEANS, LENTILS, WHITE LUPINS, CHICKPEAS (GARBANZO BEANS), DRY EDIBLE PEAS, ENGLISH AND SOUTHERN PEAS

Directions for use in the above crops in states east of and including: North Dakota, South Dakota, Wyoming, Colorado, and New Mexico (except the states east of and including: Vermont, Massachusetts, and Connecticut).

Refer to the map below for geographical use area.



Crops must be a minimum of 3 inches in height or have a minimum of one trifoliate leaf prior to a postemergence application of Imazethapyr 2 SL. If crops are treated prior to this stage, delayed maturity and/or reduced crop growth will result.

The following species are more sensitive to treatment with Imazethapyr 2 SL:

- Olathe
- Pinto variety UI-111

Restrictions

- Imazethapyr 2 SL must not be applied through an irrigation system of any type.
- Do not apply Imazethapyr 2 SL to Domino variety black turtle beans.
- Imazethapyr 2 SL must not be applied as a postemergence treatment to:
 - Chickpeas
 - Lentils
 - Lima beans
 - White lupins
- Do not exceed a maximum of one application of Imazethapyr 2 SL per year.
- Do not exceed a maximum application rate to peas and beans (except Southern peas) in this area of 3 fluid ounces per acre of Imazethapyr 2 SL per year (0.047 pound of imazethapyr acid equivalent per acre per year).
- Do not exceed a maximum application rate to Southern peas of 4 fluid ounces per acre of Imazethapyr 2 SL per year (0.063 pound of imazethapyr acid equivalent per acre per year).
- For postemergence application of Imazethapyr 2 SL, do not combine this product with petroleum oils, crop oils or methylated seed oils.
- Only nonionic surfactant can be used with Imazethapyr 2 SL as a spray additive.
- There must be a minimum of 30 days between treatment with Imazethapyr 2 SL and harvest of:
 - English peas
 - Southern peas
 - Succulent lima beans
- There must be a minimum of 60 days between treatment with Imazethapyr 2 SL and harvest of:
 - Chick peas
 - Dry edible peas
 - Lentils
 - Any other listed bean or pea types

NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO AND SMALL WHITE DRY BEANS, DRY EDIBLE PEAS, ENGLISH AND SOUTHERN PEAS

Preemergence

Apply Imazethapyr 2 SL as a preemergence treatment at a rate up to 3 fluid ounces per acre to the following:

- Dry beans
- Dry edible peas
- English peas

Apply Imazethapyr 2 SL as a preemergence treatment in the period from immediately after planting to 3 days after planting, at a rate up to 4 fluid ounces per acre to southern peas.

Imazethapyr 2 SL may be applied in a tank mixture with a registered grass herbicide. Imazethapyr 2 SL can also be applied as a preemergence application after a registered grass herbicide has been applied as a preplant incorporated treatment.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Preplant Incorporated

Apply Imazethapyr 2 SL as a preplant incorporated treatment at a rate up to 3 fluid ounces per acre to the following:

- Dry beans: navy, great northern, red kidney, black turtle, cranberry, pinto and small white type dry beans
- Dry edible peas
- English peas

Apply Imazethapyr 2 SL as a preplant incorporated treatment within a week of planting, at a rate up to 4 fluid ounces per acre to southern peas.

Imazethapyr 2 SL may be tank mixed with a registered grass herbicide when used as a preplant incorporated treatment.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Early Postemergence Applications

The Imazethapyr 2 SL application solution must be combined with a nonionic surfactant that contains a minimum of 80% active ingredient at the rate of 2 pints per 100 gallons of spray solution.

Apply Imazethapyr 2 SL as an early postemergence treatment at a rate - up to 3 fluid ounces per acre to the following:

- Dry beans that are at the one fully expanded trifoliolate leaf growth stage or beyond.
- Dry edible peas that are a minimum of 3 inches in height, but prior to flowering and prior to 5 nodes.
- English peas that are a minimum of 3 inches in height, but prior to flowering and prior to 5 nodes.

Apply Imazethapyr 2 SL as an early postemergence treatment at a rate up to 4 fluid ounces per acre to southern peas that are a minimum of 3 inches in height, but prior to flowering and prior to 5 nodes.

For control of weed species not listed in these directions, Imazethapyr 2 SL may be tank mixed with Basagran®. When Imazethapyr 2 SL is mixed with Basagran®, antagonism causes a reduction in the control of grasses. A nitrogen-based fertilizer may be added to the application solution, but only in tank mixtures containing Basagran®.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Restrictions

- Application of trifluralin before treatment with Imazethapyr 2 SL will make crop injury likely more severe.
- In Michigan and the DelMarVa peninsula (Delaware, Maryland, and Virginia), do not exceed a maximum of 2 fluid ounces of Imazethapyr 2 SL per acre when applying to sands or loamy sand soils.
- North of Highway 210 in Minnesota or in North Dakota: do not exceed a maximum of 2 fluid ounces per acre of Imazethapyr 2 SL.

LIMA BEANS, CHICKPEAS (GARBANZOS), LENTILS AND WHITE LUPINS

Restrictions

- Do not apply Imazethapyr 2 SL to white lupins that are grown in sand or loamy sand soils.
- In Michigan and the DelMarVa peninsula (Delaware, Maryland, and Virginia), do not exceed a maximum of 2 fluid ounces of Imazethapyr 2 SL per acre when applying to sands or loamy sand soils.
- North of Highway 210 in Minnesota or in North Dakota: do not exceed a maximum of 2 fluid ounces per acre of Imazethapyr 2 SL.

Preemergence Application

Apply Imazethapyr 2 SL as a preemergence application in the period between: immediately following planting to 3 days following planting, at a broadcast rate up to 3 fluid ounces per acre.

Imazethapyr 2 SL may be tank mixed with a registered grass herbicide or apply Imazethapyr 2 SL after a preplant incorporated treatment with a registered grass herbicide.

Preplant Incorporated Application

Apply Imazethapyr 2 SL within a week before planting as a preplant incorporated treatment at a broadcast rate of up to 3 fluid ounces per acre.

Imazethapyr 2 SL may be tank mixed with a registered grass herbicide when used as a preplant incorporated treatment.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Weeds Controlled

Weeds	Application Rate*
Mustard, wild	Broadcast rate of 2 fluid ounces per acre
Nightshade, black (suppression only)	
Nightshade, Eastern black (suppression only)	
Mustard, wild	Broadcast rate of 3 fluid ounces per acre**
Nightshade, black	
Nightshade, Eastern black	
Nightshade, hairy	
Pigweed, redroot	
* When applied as an early postemergence, incorporated preplant or preemergence treatment.	
**Apply postemergence treatments when weeds are less than 2 inches in height.	

The following tables list weeds controlled or suppressed when Imazethapyr 2 SL is applied at a broadcast rate of 4 fluid ounces per acre in southern peas only as a postemergence application. In some cases, Imazethapyr 2 SL may be applied as a soil application (see table footnotes). In assessing the growth stage of weed species (leaf stage), do not count cotyledon leaves.

Broadleaf Weeds Controlled

Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)
Anoda, Spurred†	4	1 to 3	Artichoke, Jerusalem	8	6 to 10	Buffalo-bur†Ω			Carpet-weed†		
Cocklebur, Common†Ω	8	1 to 8	Galinsoga†			Jimson-weed†‡	4	1 to 3	Kochia (non-ALS resistant)†	4	1 to 3
Lambs quarters†‡	*	1 to 2	Mallow, Veince∞			Marshelder†	4	1 to 3	Morning-glory, entireleaf∞	2	1 to 2
Morning-glory, ivyleaf∞	2	1 to 2	Morning-glory, pitted∞	2	1 to 2	Morning-glory, small-flower†	4	1 to 3	Morning-glory, tall∞	2	1 to 2
Mustard, Species†	4	1 to 3	Nightshade, Black†	4	1 to 3	Nightshade, Eastern Black†	4	1 to 3	Nightshade, Hairy†	4	1 to 3
Pigweed, Redroot†	4	1 to 4	Pigweed, Smooth†	4	1 to 4	Pigweed, Spiny†	4	1 to 4	Poinsettia, Wild†		
Puncturevine†			Purslane, Common†			Pusley, Florida†			Sida, Prickly†‡		
Ragweed, Common∞	4	1 to 3	Ragweed, Giant∞	4	1 to 3	Sage, Barnyard	*	1 to 3	Smartweed, Ladys-thumb†	4	1 to 3
Smartweed, Pennsylvania†	4	1 to 3	Spurge, Prostrate†	4	1 to 3	Spurge, Spotted†	4	1 to 3	Starbur, Bristly	2	1 to 2
Sunflower, Common†‡	4	1 to 3	Thisite, Canada	*	1 to 3	Velvetleaf†‡	4	1 to 3			

* Suppression only.

† Imazethapyr 2 SL will provide control when soil applied.

∞ Imazethapyr 2 SL will provide suppression when soil applied.

‡ For optimal and consistent control, apply Imazethapyr 2 SL as a preplant incorporated treatment.

Ω Only treat moderate infestation with soil application. For optimal effectiveness, application must be preplant incorporated.

Grasses and Sedges Controlled

More consistent weed species control can be achieved when Imazethapyr 2 SL is soil applied to grasses through preplant incorporated application.

Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)
Barnyard-grass [∞]	3	1 to 3	Crabgrass, Large [∞]	3	1 to 3	Crabgrass, Smooth [∞]	3	1 to 3	Cupgrass, Woolly (emerged)	3	1 to 3
Foxtail, Giant†	6	1 to 6	Foxtail, Green†	3	1 to 3	Foxtail, Robust Purple†	3	1 to 3	Foxtail, Robust White†	3	1 to 3
Foxtail, Yellow†	3	1 to 3	Goose-grass [∞]			Johnson-grass, Seedling†	6	1 to 8	Johnson-grass, Rhizome	*	1 to 8
Panicum, Fall [∞]			Panicum, Texas [∞]			Red Rice	3	1 to 3	Shatter-cane [∞]	6	1 to 8
Signalgrass, Broadleaf [∞]	4	1 to 8	Nutsedge, Purple [∞]	*	1 to 3	Nutsedge, Yellow [∞]	*				

* Suppression only.

† Imazethapyr 2 SL will provide control when soil applied.

∞ Imazethapyr 2 SL will provide suppression when soil applied.

‡ For optimal and consistent control, apply Imazethapyr 2 SL as a preplant incorporated treatment.

Ω Only treat moderate infestation with soil application. For optimal effectiveness, application must be preplant incorporated.

RED KIDNEY BEANS (For use in California only)

The application solution of Imazethapyr 2 SL must be combined with a non-ionic surfactant with at least 80% active ingredient at the rate of 2 pints of surfactant per 100 gallons of application solution.

Apply Imazethapyr 2 SL as a postemergence application at a rate of 3 fluid ounces per acre when red kidney beans are at or beyond the 1 fully expanded trifoliate leaf growth stage and when weeds are growing actively. If Imazethapyr is applied prior to the one true leaf stage of growth, delayed maturity and/or reduced crop growth will result.

Cultivate treated areas 7 to 10 days following treatment with Imazethapyr 2 SL. Cultivation will increase residual weed control, especially in dry conditions.

Restrictions

- Do not apply Imazethapyr to weeds and crops that are suffering stress from temperature, moisture, draught, etc.
- Do not apply Imazethapyr 2 SL to Kidney Beans via aerial application.
- Do not exceed a maximum of 3 fluid ounces per acre of Imazethapyr 2 SL per year (0.047 pounds of imazethapyr acid equivalent per year).
- Do not exceed a maximum of one application per year with Imazethapyr 2 SL.
- There must be a minimum of 60 days between treatment with Imazethapyr 2 SL and harvest of kidney beans.

Weeds Controlled

Treatment with Imazethapyr 2 SL as a postemergence application will control/suppress the following weeds. Make applications at or prior to the listed maximum leaf stage.

Weeds	Max. Leaf Growth Stage	Height
Kochia (non-ALS resistant)	4	1-3 inches
Mustard, Wild		
Nightshade, Black		
Nightshade, Eastern Black		
Pigweed, Redroot		
Nightshade, Hairy	4	1-2 inches

SNAP BEANS

(For use in the states of Alabama, Florida, Georgia, Illinois, Iowa, Indiana, Minnesota, Michigan, New Jersey, North Carolina and Wisconsin.)

Imazethapyr 2 SL, when applied as detailed below, will provide suppression of/will reduce competition from the following weed species:

Mustard, Wild
Nightshade, Eastern Black
Pigweed, Redroot
Purslane, Common

Preplant Incorporated Treatment

Apply Imazethapyr 2 SL within one week of planting at a rate of 1.5 fluid ounces per acre.

When applied as a preplant incorporated treatment, Imazethapyr 2 SL can be tank mixed with a registered grass herbicide. Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Preemergence

Apply Imazethapyr 204 SL either immediately following planting or up to a maximum of one day following planting at a rate of 1.5 fluid ounces per acre.

Preemergence treatment may be made following a preplant incorporated treatment with a registered grass herbicide. Alternatively, Imazethapyr 2 SL can be tank mixed with a registered grass herbicide and applied as a preemergence treatment. Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Restrictions

- Do not exceed a maximum of 1.5 fluid ounces per acre of Imazethapyr 2 SL per year (0.023 pounds of imazethapyr acid equivalent per year).
- Do not exceed a maximum of one application of Imazethapyr 2 SL per year.
- Do not apply Imazethapyr to snap beans via aerial application.
- Imazethapyr 2 SL must not be applied after July 31st in all states listed except New Jersey.
- Imazethapyr 2 SL must not be applied in New Jersey after June 20th.
- There must be a minimum of 30 days between treatment with Imazethapyr 2 SL and harvest of snap beans.

SNAP BEANS

(Directions for use in the states of Arkansas, Missouri, Oklahoma, Texas (counties of Bailey, Castro, Lamb and Parmer only) and New Mexico (counties of Curry and Roosevelt only).)

Imazethapyr 2 SL, when applied as detailed below, will provide suppression of/will reduce competition from the following weed species:

Nightshade, Eastern Black

Pigweed, Redroot

Restrictions

- Do not exceed a maximum of 1.5 fluid ounces per acre of Imazethapyr 2 SL per year (0.023 pounds of imazethapyr acid equivalent per year).
- Do not exceed a maximum of one application of Imazethapyr 2 SL per year.
- Do not apply Imazethapyr to snap beans via aerial application.
- Imazethapyr 2 SL must not be applied after July 31st.
- There must be a minimum of 30 days between treatment with Imazethapyr 2 SL and harvest of snap beans.

Postemergence

Apply Imazethapyr as a postemergence treatment at the rate of 1.5 fluid ounces per acre in a tank mixture with Basagran®. The application solution must be a combined with a nonionic surfactant that contains a minimum of 80% active ingredient at the rate of 2 pints per 100 gallons of spray solution.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Postemergence

Make postemergence applications when snap beans have a minimum of one true leaf. Treatment prior to this growth stage will result in delayed maturity and/or reduced crop growth.

**CHICKPEAS, DRY EDIBLE PEAS, LENTILS, LIMA BEANS AND SUCCULENT PEAS,
(For use in the states of Idaho, Montana, Nevada, Oregon, Utah, and Washington.)**

Restrictions

- Do not exceed a maximum of 3 fluid ounces per acre of Imazethapyr 2 SL per year (0.047 pounds of imazethapyr acid equivalent per year) to beans and peas in the region specified.
- Do not exceed a maximum of one application of Imazethapyr 2 SL per year.
- There must be a minimum of 60 days between treatment with Imazethapyr 2 SL and harvest of chickpeas, dry edible peas, dry lima beans and lentils.
- There must be a minimum of 30 days between treatment with Imazethapyr 2 SL and harvest of succulent lima beans and succulent peas.

Preplant Treatment (minimum tillage and no-till and systems only)

Apply Imazethapyr 2 SL in the fall before planting in the spring. Moisture is necessary for the treatment to be incorporated and activated. Apply this product as a preplant treatment within 30 days prior to planting in minimum till or no-till systems at a broadcast rate of 3 fluid ounces per acre. When Imazethapyr 2 SL is applied in the fall, it must be applied prior to the ground freezing in the target area and when soil temperature at a depth of 4 inches is less than 55°F.

Weed control may be unpredictable as weather factors will influence product longevity and activity.

Restrictions

- When applying Imazethapyr 2 SL as a preplant incorporated treatment, do not incorporate to a depth greater than 3 inches.

Preplant Incorporated Treatment

Apply this product as a preplant treatment within 7 days prior to planting at a broadcast rate of 3 fluid ounces per acre.

Restrictions

- When applying Imazethapyr 2 SL as a preplant incorporated treatment, do not incorporate to a depth greater than 3 inches.

Preemergence Treatment

Apply Imazethapyr 2 SL as a preemergence treatment following planting but before crop emergence at the broadcast rate of 3 fluid ounces per acre.

In order to control dogfennel (mayweed-chamomile) and lambsquarters, Imazethapyr 2 SL may be tank mixed with Sencor® DF. Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Postemergence Treatment (Dry Edible Peas)

Apply Imazethapyr 2 SL to dry edible peas at a rate of 2 fluid ounces per acre when peas are a minimum of 3 inches in height or have a minimum of one trifoliolate leaf. Application prior to this growth stage will result in delayed maturity and/or reduced crop growth. The application solution must be a combined with a nonionic surfactant that contains a minimum of 80% active ingredient at the rate of 2 pints per 100 gallons of spray solution.

In order to control weed species not specified in this label, Imazethapyr 2 SL may be tank mixed with Basagran®. When Imazethapyr 2 SL is mixed with Basagran®, antagonism causes a reduction in the control of grasses. Add a nitrogen-based liquid fertilizer to the application solution, but only in tank mixtures containing Basagran® at a rate of 1.25 to 2.5 gallons per 100 gallons of application solution. Alternatively, combine ammonium sulfate with the application solution at the rate of 12-15 pounds per 100 gallons of spray solution.

Restriction

- Do not make postemergence applications with Imazethapyr 2 SL in chickpeas, lentils or lima beans.

Weeds Controlled

At the application rate of 3 fluid ounces per acre, Imazethapyr 2 SL will provide control of the following weed species:

Preplant Incorporated Treatment	Preemergence Treatment
Buckwheat, wild	Buckwheat, wild
Kochia (non-ALS resistant)	Kochia (non-ALS resistant)
Lambsquarters, common	Mustard, wild
Mustard, wild	Nightshade, Black
Nightshade, Black	Nightshade, Eastern black
Nightshade, Eastern black	Nightshade, Hairy
Nightshade, Hairy	Pigweed, redroot
Pigweed, redroot	Shepherdspurse

Preplant Incorporated Treatment	Preemergence Treatment
Shepherdspurse	Thistle, Russian
Thistle, Russian	

When applied as a postemergence treatment at the broadcast application rate of 2 fluid ounces per acre, Imazethapyr 2 SL will provide control/suppression of the following weed species:

- Nightshade, Black (suppression only)
- Nightshade, Eastern Black (suppression only)
- Nightshade, Hairy (suppression only)
- Mustard, Wild (control)

CHICKPEAS (For use in the states of Arizona and California.)

Restrictions

- Do not exceed a maximum of 3 fluid ounces per acre of Imazethapyr 2 SL per year (0.047 pounds of imazethapyr acid equivalent per year) to chickpeas in the region specified.
- Do not exceed a maximum of one application of Imazethapyr 2 SL per year.
- There must be a minimum of 60 days between treatment with Imazethapyr 2 SL and harvest of dry chickpeas.
- There must be a minimum of 30 days between treatment with Imazethapyr 2 SL and harvest of succulent chickpeas.

Preplant Incorporated Treatment

Apply Imazethapyr 2 SL as a preplant incorporated treatment within a week before planting at a broadcast rate of 0-3 fluid ounces per acre. When applied as a preplant incorporated treatment, Imazethapyr 2 SL may be tank mixed with a registered grass herbicide.

Preemergence Treatment

Apply Imazethapyr as a preemergence treatment during the period from immediately following planting to a maximum of 3 days following planting at a broadcast rate of up to 3 fluid ounces per acre.

Imazethapyr may be tank mixed with a registered grass herbicide. Alternatively, Imazethapyr 2 SL may be applied as a preemergence treatment sequentially after an application with a registered grass herbicide as a preplant incorporated treatment.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Weeds Controlled

Imazethapyr 2 SL will provide control of the following weed species:

Preplant Incorporated Treatment	Preemergence Treatment
Buckwheat, wild	Buckwheat, wild
Kochia (non-ALS resistant)	Kochia (non-ALS resistant)
Lambsquarters, common	Mustard, wild

Preplant Incorporated Treatment	Preemergence Treatment
Mustard, wild	Nightshade, Black
Nightshade, Black	Nightshade, Eastern black
Nightshade, Eastern black	Nightshade, Hairy
Nightshade, Hairy	Pigweed, redroot
Pigweed, redroot	Shepherdspurse
Shepherdspurse	Thistle, Russian
Thistle, Russian	

EDIBLE VEGETABLE LEGUMES: RESTRICTIONS

- Snap beans, lima beans, chickpeas (in Arizona and California), Southern peas and English peas must not be harvested for a minimum of 30 days following treatment with Imazethapyr 2 SL.
- Dry edible peas, lentils, chickpeas, red kidney beans, and all other dry bean or pea types specified in these directions dry edible peas, lentils, chickpeas, red kidney beans, and other dry bean or pea types listed on this label listed on this label must not be harvested for a minimum of 60 days following treatment with Imazethapyr 2 SL.

PEANUTS

(Not for use in California. In Arizona, use only in La Paz and Yuma counties.)

Imazethapyr 2 SL will provide control of many grasses, sedges and broadleaf weed species. When heavy pressure from grass or common lambsquarters is likely, tank mix Imazethapyr 2 SL with a registered soil-applied grass herbicide (see below).

Application Method	Imazethapyr 2SL Use Rate/Directions
Preplant Incorporated	4 fluid ounces per acre (broadcast rate)
Preemergence	
Ground Cracking	
Postemergence	
Sequential Applications: - Preplant Incorporated or Preemergence - Ground Cracking or Postemergence	2 fluid ounces per acre in a soil treatment (preplant incorporated or preemergence), then 2 fluid ounces per acre applied postemergence or at ground cracking.

Restrictions

- Do not exceed a maximum of 4 fluid ounces per acre of Imazethapyr 2 SL per year (0.063 pounds of imazethapyr acid equivalent per year) to peanuts.
- Peanuts must not be harvested for a minimum of 85 days following treatment with Imazethapyr 2 SL.
- Treated peanut hay, straw or forage must not be used for grazing or feed.
- An application of Pursuit® Plus EC must not be made in the same year as an application of Imazethapyr 2 SL. An application of Classic® may be made as a postemergence treatment after treatment with Imazethapyr 2 SL. See the Classic® label directions prior to use.

Weeds Controlled: Soil Applied/At-Crack

Apply Imazethapyr 2 SL as an at-crack treatment when the emerging peanut seedling cracks the soil, usually 10 to 14 days after planting. At the soil cracking stage, weeds are usually seedlings or have not germinated. If weed species have >2 true leaves, see the Weeds Controlled: Postemergence Treatment section below.

Imazethapyr 2 SL will provide control/suppression of the following weed species when soil applied or as a treatment applied at ground cracking:

Broadleaf Weeds	
Alligator weed*	Pigweed, Redroot
Anoda, Spurred**	Pigweed, Smooth
Buffalobur	Pigweed, Spiny
Carpetweed	Poinsettia, wild
Cocklebur, Common***	Puncturevine
Devilsclaw	Purslane, common
Galinsoga	Pusley, Florida
Jimsonweed**	Ragweed, Common (suppression only)
Lambsquarters, Common**	Ragweed, Giant (suppression only)
Morningglory, Entireleaf***	Sida, prickly (Teaweed)**
Morningglory, Ivyleaf***	Smartweed, ladystumb
Morningglory, Pitted***	Smartweed, Pennsylvania
Morningglory, smallflower	Spurge, Prostrate
Morningglory, tall***	Spurge, Spotted
Mustard Species	Spurge, Toothed
Nightshade, Black	Sunflower**
Nightshade, Eastern Black	Velvetleaf**
Nightshade, Hairy	

*Imazethapyr provides control as an at-crack treatment only.

**For optimal and consistent control, apply Imazethapyr 2 SL as a preplant incorporated treatment.

***Suppression only when soil applied.

For optimal and consistent control of grasses and sedges, apply Imazethapyr 2 SL as a preplant incorporated treatment.

Grasses and Sedges	
Alligator weed*	Pigweed, Redroot
Anoda, Spurred**	Pigweed, Smooth
Buffalobur	Pigweed, Spiny
Carpetweed	Poinsettia, wild

Grasses and Sedges	
Cocklebur, Common***	Puncturevine
Devilsclaw	Purslane, common
Galinsoga	Pusley, Florida
Jimsonweed**	Ragweed, Common (suppression only)
Lambsquarters, Common**	Ragweed, Giant (suppression only)
Morningglory, Entireleaf***	Sida, prickly (Teaweed)**
Morningglory, Ivyleaf***	Smartweed, ladysthumb
Morningglory, Pitted***	Smartweed, Pennsylvania
Morningglory, smallflower	Spurge, Prostrate
Morningglory, tall***	Spurge, Spotted
Mustard Species	Spurge, Toothed
Nightshade, Black	Sunflower**
Nightshade, Eastern Black	Velvetleaf**
Nightshade, Hairy	

*Imazethapyr provides control as an at-crack treatment only.

**For optimal and consistent control, apply Imazethapyr 2 SL as a preplant incorporated treatment.

***Suppression only when soil applied.

Weeds Controlled: Postemergence Treatment

Broadleaf weeds controlled when Imazethapyr 2 SL is applied as a postemergence treatment:

Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)
Alligator Weed	4	1 to 3	Anoda, Spurred	2	1 to 2	Buffalobur	*	1 to 3	Cocklebur, Common	8	1 to 8
Jimsonweed	4	1 to 3	Lambs-quarters, Common	*	1 to 2	Morning-glory, Entireleaf	2	1 to 2	Morning-glory, Ivyleaf	2	1 to 2
Morning-glory, Pitted	2	1 to 2	Morning-glory, Small-flower	4	1 to 3	Morning-glory, Tall	2	1 to 2	Mustard, Species	4	1 to 3
Nightshade, Black	4	1 to 3	Nightshade, Eastern black	4	1 to 3	Nightshade, Hairy	4	1 to 3	Pigweed, Redroot	8	1 to 8
Pigweed, Smooth	8	1 to 8	Pigweed, Spiny	8	1 to 8	Ragweed, Common	4	1 to 3	Ragweed, Giant	4	1 to 3
Smartweed, Ladysthumb	4	1 to 3	Smartweed, Pennsylvania	4	1 to 3	Spurge, Prostrate	4	1 to 3	Spurge, Spotted	4	1 to 3

Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)
Starbur, Bristly	2	1 to 2	Sunflower	4	1 to 3	Velvetleaf	4	1 to 3			

*Suppression only.

Grass and sedges controlled when Imazethapyr 2 SL is applied as a postemergence treatment:

Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)
Barnyard-grass	3	1 to 3	Crabgrass, Large	3	1 to 3	Crabgrass, Smooth	3	1 to 3	Cupgrass, Woolly	3	1 to 3
Foxtail, Giant	6	1 to 6	Foxtail, Green	3	1 to 3	Foxtail, Yellow	3	1 to 3	Johnson-grass, Seedling	6	1 to 8
Johnson-grass, Rhizome	*	6 to 12	Red Rice	3	1 to 3	Shattercane	6	1 to 8	Signalgrass, Broadleaf	4	1 to 6
Purple Nutsedge	3	1 to 3	Yellow Nutsedge	3	1 to 3						

*Suppression only.

In West Texas and New Mexico, wait until late cracking (most of the peanuts have emerged) before applying this product.

Weeds Controlled: Sequential Applications

Apply Imazethapyr 2 SL as a sequential treatment as follows:

1. Soil apply 2 fluid ounces of Imazethapyr 2 SL per acre (preemergence or preplant incorporated)
2. Apply 2 fluid ounces of Imazethapyr 2 SL as a postemergence or ground-crack treatment.

For weeds controlled by sequential applications, refer to the Weeds Controlled: Soil Applied/At-Crack sections and tables above in the PEANUT section.

Yellow and Purple Nutsedge: Sequential Application

Control of purple nutsedge and yellow nutsedge is increased by sequential application of Imazethapyr 2 SL. Make the second treatment prior to nutsedge exceeding the 3 leaf growth stage.

Tank Mix Combinations for Grass Species Control

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Imazethapyr 2 SL may be tank mixed with specified registered products/registered product containing specified active ingredients in order to control weeds listed below that are not specified for control by applications of Imazethapyr 2 SL alone:

Restriction

- Tank mixtures containing registered products with trifluralin as an active ingredient must be applied as a preplant incorporated treatment only.

Barnyardgrass: Tank mix Imazethapyr with Prowl® 3.3 EC, Trifluralin, Lasso®, Metolachlor, Balan™² or Sonalan™²
Crabgrass, Smooth: Tank mix Imazethapyr with Prowl® 3.3 EC, Trifluralin, Lasso®, Metolachlor, Balan™² or Sonalan™²
Crabgrass, Large: Tank mix Imazethapyr with Prowl® 3.3 EC, Trifluralin, Lasso®, Metolachlor, Balan™² or Sonalan™²
Crowfootgrass: Tank mix Imazethapyr with Prowl® 3.3 EC, Trifluralin, Balan™² or Sonalan™²

Goosegrass: Tank mix Imazethapyr with Prowl® 3.3 EC, Trifluralin, Lasso®, Metolachlor, Balan™² or Sonalan™²
Panicum, Fall: Tank mix Imazethapyr with Prowl® 3.3 EC, Trifluralin, Lasso®, Metolachlor, Balan™² or Sonalan™²
Panicum, Texas: Tank mix Imazethapyr with Prowl® 3.3 EC, Trifluralin, Balan™² or Sonalan™²

Sandbur, Field: Tank mix Imazethapyr with Prowl® 3.3 EC, Trifluralin, Balan™² or Sonalan™²

Signalgrass, broadleaf: Tank mix Imazethapyr with Prowl® 3.3 EC (preplant incorporated treatment only), Trifluralin, Lasso®, Metolachlor, Balan™² or Sonalan™²

Witchgrass: Tank mix Imazethapyr with Prowl® 3.3 EC, Trifluralin, Lasso®, Metolachlor or Sonalan™²

Tank mix Imazethapyr 2 SL with Prowl® and apply as a preplant incorporated treatment in order to suppress rhizome Johnsongrass and itchgrass.

Tank mix Imazethapyr 2 SL with a grass herbicide (e.g. Whip® or Poast Plus®) to provide control of grasses not specified in this label. However, the grass control effectiveness of a tank mix partner may be reduced by mixing with Imazethapyr 2 SL. This can be avoided by making a sequential application as follows:

1. Treat with Imazethapyr 2 SL
2. Treat with a registered postemergence grass herbicide 7 days after the Imazethapyr 2 SL treatment.

Or:

1. Treat with a registered postemergence grass herbicide.
2. Treat with Imazethapyr 2 SL 3 days after the registered postemergence grass herbicide treatment.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Tank Mix Combinations for Broadleaf Species Control

For control of broadleaf weeds not specified in this label, Imazethapyr 2 SL may be tank mixed with a registered product, for example: Basagran®, Ultra Blazer®, and products containing 2,4-DB. Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Tank mix Imazethapyr 2 SL with a registered product containing 2,4-DB in order to control the following species:

- Common ragweed
- Morningglories
- Prickly sida
- Sicklepod

Imazethapyr 2 SL may be applied in a tank mixture with the following products:

- Bravo®
- Bravo® S
- Orthere®
- Solubor®.

SOYBEANS

(See below for additional directions for soybeans in North Dakota and Minnesota north of Highway 210.)

Apply Imazethapyr 2 SL in soybeans as one of the following treatments:

- early preplant
- preplant incorporated
- preemergence
- postemergence (including in minimum tillage and no-tillage areas)

For all applications, use a rate of 4 fluid ounces of Imazethapyr 2 SL per acre.

Restrictions:

- Do not exceed a maximum of 4 fluid ounces per acre of Imazethapyr 2 SL per year (0.063 pounds of imazethapyr acid equivalent per year).
- Do not exceed a maximum of one application of Imazethapyr 2 SL per season.
- Soybeans must not be harvested for a minimum of 85 days following treatment with Imazethapyr 2 SL.
- Applications of Imazethapyr 2 SL must be made prior to soybean bloom.
- Treated soybean hay, straw or forage must not be used for grazing or feed.
- Do not tank mix Imazethapyr 2 SL with products containing clomazone (e.g. Command®). Where Command® has been applied as a soil application, Imazethapyr 2 SL may be applied as a postemergence treatment.

For furrow irrigated soybeans, ensure that soil is tilled prior to planting winter wheat or barley. Break up the beds and mix the soil with tillage equipment calibrated to cut to a depth of 4-6 inches.

Weeds Controlled

The table below provides a list of weeds controlled by Imazethapyr 2 SL. For optimal postemergence applications, suppression/control, apply Imazethapyr 2 SL before target species reach the specified height below. In assessing the growth stage of weed species (leaf stage), do not count cotyledon leaves.

Imazethapyr 2 SL will provide control of many grasses, sedges and broadleaf weed species. When heavy pressure from grass or common lambsquarters is likely, tank mix Imazethapyr 2 SL with a registered soil-applied grass herbicide such as Prowl®.

Weeds Controlled: Broadleaf Weeds

Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)
Alligator weed	4	1 to 3	Anoda, Spurred†	4	1 to 3	Artichoke, Jerusalem	8	6 to 10	Buffalo-bur††	*	1 to 3
Carpet-weed†			Cocklebur, Common†	8	1 to 8	Galinsoga†			Jimson-weed††	4	1 to 3
Kochia (non-ALS resistant)†	4	1 to 3	Lambs-quarters††	*	1 to 2	Mallow, Veince∞			Marshelder†	4	1 to 3

Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)
Morning-glory, entireleaf [∞]	2	1 to 2	Morning-glory, ivyleaf [∞]	2	1 to 2	Morning-glory, pitted [∞]	2	1 to 2	Morning-glory, small-flower†	4	1 to 3
Morning-glory, tall [∞]	2	1 to 2	Mustard, Species†	4	1 to 3	Nightshade, Black†	4	1 to 3	Nightshade, Eastern Black†	4	1 to 3
Nightshade, Hairy†	4	1 to 3	Pigweed, Redroot†	8	1 to 8	Pigweed, Smooth†	8	1 to 8	Pigweed, Spiny†	8	1 to 8
Poinsettia, Wild†			Puncturevine†			Purslane, Common†			Pusley, Florida†		
Sida, Prickly††			Ragweed, Common [∞]	*	1 to 3	Ragweed, Giant [∞]	*	1 to 3	Sage, Barnyard	1 to 3	
Smartweed, Lady-thumb†	4	1 to 3	Smartweed, Pennsylvania†	4	1 to 3	Spurge, Prostrate†	4	1 to 3	Spurge, Spotted†	4	1 to 3
Starbur, Bristly	2	1 to 2	Sunflower, Common††	4	1 to 3	Thisite, Canada	*	1 to 3	Velvetleaf††	4	1 to 3

* Suppression only.

† Imazethapyr 2 SL will provide control when soil applied.

∞ Imazethapyr 2 SL will provide suppression when soil applied.

‡ For optimal and consistent control, apply Imazethapyr 2 SL as a preplant incorporated treatment.

Ω For optimal effectiveness, application must be preplant incorporated.

Weeds Controlled: Grass Weeds and Sedges

More consistent weed species control can be achieved when Imazethapyr 2 SL is soil applied to grasses through preplant incorporated application.

Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)
Barnyard-grass [∞]	3	1 to 3	Crabgrass, Large [∞]	3	1 to 3	Crabgrass, Smooth [∞]	3	1 to 3	Cupgrass, Woolly (emerged)	3	1 to 3
Foxtail, Giant†	6	1 to 6	Foxtail, Green†	3	1 to 3	Foxtail, Yellow†	3	1 to 3	Goose-grass [∞]		
Johnson-grass, Seeding†	6	1 to 8	Johnson-grass, Rhizome	*	6 to 12	Millet, Wild Proso [∞]	*	1 to 3	Panicum, Fall [∞]		

Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)	Weed	Max Leaf Stage	Size (inches)
Panicum, Texas [∞]			Red Rice	3	1 to 3	Shatter-cane [∞]	6	1 to 8	Signalgrass, Broadleaf [∞]	4	1 to 8
Nutsedge, Purple [∞]	*	1 to 3	Nutsedge, Yellow [∞]	*	1 to 3						

* Suppression only.

† Imazethapyr 2 SL will provide control when soil applied.

∞ Imazethapyr 2 SL will provide suppression when soil applied.

‡ For optimal and consistent control, apply Imazethapyr 2 SL as a preplant incorporated treatment.

Ω For optimal effectiveness, application must be preplant incorporated.

Tank Mixtures for Grass Weeds

Imazethapyr 2 SL can be tank mixed with a registered grass herbicide in order to provide control of grass weed species not listed in this label. Tank mixtures may also increase control of some broadleaf target species (e.g. pigweeds and common lambsquarters).

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply. Imazethapyr may be tank mixed with specified registered products/registered product containing specified active ingredients in order to control weeds listed below that are not specified for control by applications of Imazethapyr 2 SL alone:

Restrictions

- Tank mixtures containing registered products with Trifluralin as an active ingredient must be applied as a preplant incorporated treatment only.
- Do not tank mix Imazethapyr 2 SL with products containing clomazone (e.g., Command[®]). Where Command[®] has been applied as a soil application, Imazethapyr 2 SL may be applied as a postemergence treatment.

Barnyardgrass: Tank mix Imazethapyr with Prowl[®] 3.3 EC, Trifluralin, Lasso[®], Metolachlor or Outlook[®]

Crabgrass, Smooth: Tank mix Imazethapyr with Prowl[®] 3.3 EC, Trifluralin, Lasso[®], Metolachlor, Balan[™] or Sonalan[™]

Crabgrass, Smooth: Tank mix Imazethapyr with Prowl[®] 3.3 EC, Trifluralin, Lasso[®], Metolachlor or Outlook[®].

Crabgrass, Large: Tank mix Imazethapyr with Prowl[®] 3.3 EC, Trifluralin, Lasso[®], Metolachlor or Outlook[®]

Crowfootgrass: Tank mix Imazethapyr with Prowl[®] 3.3 EC or Trifluralin.

Goosegrass: Tank mix Imazethapyr with Prowl[®] 3.3 EC, Trifluralin, Lasso[®], Metolachlor or Outlook[®].

Millet, Wild Pros: Tank mix Imazethapyr with Prowl[®] 3.3 EC or Trifluralin.

Panicum, Fall: Tank mix Imazethapyr with Prowl[®] 3.3 EC, Trifluralin, Lasso[®], Metolachlor or Outlook[®].

Panicum, Texas: Tank mix Imazethapyr with Prowl[®] 3.3 EC or Trifluralin.

Sandbur, Field: Tank mix Imazethapyr with Prowl[®] 3.3 EC or Trifluralin.

Signalgrass, broadleaf: Tank mix Imazethapyr with Prowl[®] 3.3 EC (preplant incorporated treatment only), Trifluralin, Lasso[®], Metolachlor or Outlook[®].

Witchgrass: Tank mix Imazethapyr with Prowl[®] 3.3 EC, Trifluralin, Lasso[®], Metolachlor or Outlook[®].

Tank mix Imazethapyr 2 SL with Prowl[®] and apply as a preplant incorporated treatment in order to suppress rhizome Johnsongrass and Itchgrass.

Grasses not listed in this label and volunteer corn may be controlled using a tank mixture of Imazethapyr 2 SL and a grass herbicide (e.g. Poast Plus®). For optimal results, combine the application solution with a liquid fertilizer and crop oil concentrate.

Tank Mixing Imazethapyr 2 SL with Poast Plus®

Imazethapyr 2 SL may be tank mixed with Poast Plus® at the following rates in order to increase control of grass weed species, particularly where infestation of grass is heavy and for grass species not specified in this label. Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Use the higher specified rate of Poast Plus® where there is a mixture of grass weed species present. See the Poast Plus® label for specified additional weed species controlled. For optimal effectiveness, make applications when the target species are in the size range indicated in the "Application Size" column.

Grass Species	Application size (inches)	Poast Plus® Application Rate* (fluid ounces per acre)	Imazethapyr 2 SL Application Rate (fluid ounces per acre)
Barnyardgrass	3-8	24	4
Crabgrass, Large	3-6	24	4
Crabgrass, Smooth	3-6	24	4
Cupgrass, Woolly	3-8	24	4
Foxtail, Giant	3-8	16	4
Foxtail, Green	3-8	24	4
Foxtail, Yellow	3-8	24	4
Goosegrass	3-6	24	4
Johnsongrass, Seedling	3-8	24	4
Junglerice	3-8	16	4
Panicum, Fall	3-8	16	4
Panicum, Texas	3-8	16	4
Shattercane	3-12	12	4
Signalgrass, Broadleaf	3-8	16	4
Sprangletop, Red	3-8	24	4
Volunteer Corn	4-10	20	4
Wild Proso Millet	4-10	12	4
Witchgrass	3-8	24	4

The effectiveness of Poast Plus® can be reduced by mixing it with Imazethapyr 2 SL. In order to avoid reduced effectiveness, apply Poast Plus® 7 days after treatment with Imazethapyr 2 SL. Alternatively, apply Imazethapyr 2 SL 3 days after treatment with Poast Plus®. When making sequential applications, see the Poast Plus® directions for application instructions.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Tank Mixtures for Broadleaf Weeds

Imazethapyr 2 SL can be tank mixed with the following registered products in order to enhance control of broadleaf weed species:

- Basagran®
- Cobra®
- FirstRate™
- Flexstar®
- Gramoxone® Extra
- Reflex®
- Storm®
- Ultra Blazer®

Imazethapyr 2 SL can be tank mixed with Roundup Ultra® for control of broadleaf species in Roundup Ready® soybeans.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Tank Mixture with Ultra Blazer®

A tank mix of Ultra Blazer® and Imazethapyr 2 SL will provide enhanced control of pigweeds, common ragweed, giant ragweed and waterhemp. Use the following rates according to weed height. Use the higher rate in the specified rate range if weed species are dense or if common ragweed is present:

Broadleaf Species	Ultra Blazer® Application Rate (fluid ounces per acre)			Imazethapyr 2 SL Application Rate (fluid ounces per acre)
	8-10 fl. oz.	12-14 fl. oz.	16-20 fl. oz.	
	Application size (inches)			
Common ragweed	1-4	4-6	6-8	4
Giant Ragweed		1-6	6-8*	4
Pigweed species	1-4	4-6	6-8	4
Waterhemp, common	1-4	4-6	6-8	4
Waterhemp, tall	1-4	4-6	6-8	4

*Use the higher specified rate if giant ragweed is 8 inches in height.

Sequential Applications with Ultra Blazer®

Imazethapyr 2 SL may be applied sequentially with Ultra Blazer® at the following rates. Use the higher rate in the specified rate range if weed species are dense or if common ragweed is present:

Broadleaf Species	Ultra Blazer® Application Rate (ounces per acre)		
	8-10 fl. oz.	12-14 fl. oz.	16-20 fl. oz.
	Application size (inches)		
Common ragweed	1-4	4-6	6-8
Giant Ragweed		1-6	6-8*
Pigweed species	1-4	4-6	6-8

Broadleaf Species	Ultra Blazer® Application Rate (ounces per acre)		
	8-10 fl. oz.	12-14 fl. oz.	16-20 fl. oz.
	Application size (inches)		
Waterhemp, common	1-4	4-6	6-8
Waterhemp, tall	1-4	4-6	6-8

*Use the higher specified rate if giant ragweed is 8 inches in height.

Tank Mixture with FirstRate™

Imazethapyr 2 SL may be tank mixed with FirstRate™ in order to enhance control of ragweed weed species (giant and common). Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Tank Mixtures with Products Containing Sulfentrazone

This product may be tank mixed for application with registered products containing sulfentrazone in order to enhance control of weed species in soybeans (e.g. Authority® and Canopy® XL).

Alternatively, Imazethapyr 2 SL may be applied postemergence, sequentially to soybeans that have been treated with a registered product containing sulfentrazone.

Restriction

- Registered products containing sulfentrazone are only specified for use in soil applications in soybeans.

Tank Mixture with Harmony® GT

Imazethapyr 2 SL may be tank mixed with Harmony® GT in order to enhance control of common lambsquarters.

In order to achieve the most effective target species control, treat sequentially as follows:

1. Treat the area with a grass herbicide that is soil applied (e.g. such as TRI-4®, Prowl® or a registered product containing trifluralin).
2. Make a postemergence application with Imazethapyr 2 SL.

In the event that control of common lambsquarters is inadequate with the soil applied application, Harmony® GT may be tank mixed with Imazethapyr 2 SL in order to enhance control.

A tank mixture of Imazethapyr 2 SL plus Harmony® GT will cause stunting and/or severe damage to soybean crops, particularly where conditions are hot and humid. The applicator assumes sole responsibility for any injury that results from application of this tank mixture in soybeans. Apply the tank mix in soybeans at the 1-3 trifoliolate growth stage.

When tank mixing Imazethapyr 2 SL and Harmony® GT, mix as follows:

Imazethapyr 2 SL	Harmony® GT	Non-ionic Surfactant	Liquid Nitrogen Based Fertilizer ^{1,2}
4 fl. oz. per acre	1/24 oz. per acre	1 qt. per 100 gallons of spray mixture	1.25-2.5 gallons per 100 gallons of spray solution

¹ e.g. 28%N, 32%N, or 10-34-0

² Spray grade ammonium sulfate may be used as an alternative to liquid nitrogen based fertilizer at a rate of 12-15 lbs. per 100 gallons of spray solution.

Tank Mixture with Scepter® DG (Regions 2 and 3)

Imazethapyr 2 SL may be tank mixed with Scepter® DG in Region 2 and 3 in order to enhance control of common sunflower and volunteer corn. For details of Regions 2 and 3, refer to the Scepter® DG label. Refer to the Scepter® DG label for additional weeds controlled/suppressed. This tank mixture may also be applied in South Dakota in the following counties: Bon Homme, Davison, Hanson, Hutchinson, Kingsbury, Lake, McCook, Miner and Yankton.

This tank mixture will provide suppression of volunteer corn when applied to volunteer corn no greater than 10 inches in height.

This tank mixture will enhance control of common sunflowers when applied to sunflowers no greater than 3 inches in height.

Read and follow the label instructions of all tank mix partners. Ensure the product(s) used are labeled for the intended use and mixture. The most restrictive directions must apply.

Restrictions

- This tank mixture must not be applied in North Dakota or in Minnesota north of Highway 210.

Tank mix as follows:

Imazethapyr 2 SL	Scepter® DG
4 fl. oz. per acre	0.53 fl. oz. per acre

At the above rate, one soluble bag of Scepter® DG (14 oz.) will treat 26.4 acres.

SOYBEANS

(North Dakota and Minnesota north of Highway 210.)

Apply Imazethapyr 2 SL in North Dakota and Minnesota north of Highway 210, as a postemergence treatment only at a rate of 3 fluid ounces per acre.

Weed Species	Application Size (in inches)	Max. Leaf Growth Stage (in leaves)
Cocklebur, Common*	4	1-4
Pigweed, Redroot		
Kochia (non-ALS resistant)	4	1-3
Mustard, Species		
Nightshade, Black		
Nightshade, Eastern black		
Nightshade, Hairy	3	1-4
Wild oats (reduced competition)		

Imazethapyr 2 SL may be tank mixed with Ultra Blazer® in order to enhance control of common cocklebur. Mix 3 fluid ounces of Imazethapyr 2 SL per acre with 12 fluid ounces Ultra Blazer® per acre.

ROTATIONAL CROPS: POST APPLICATION PLANTING INTERVALS

In most cases, normal growth of rotational crops can be expected when Imazethapyr 2 SL is used in accordance with this label. However, crop damage is always a risk due to possible variations in and extremes of agronomic and environmental factors.

The following crops can be planted following an application of Imazethapyr 2 SL at the specified interval:

Restriction

- Do not plant crops prior to the end of the specified interval or crop injury may result.
- Refer to the Rotational Crops: Exceptions section for additional intervals.

Crop	Post Application Interval
Lima beans	Crops may be planted immediately after application of Imazethapyr 2 SL
Peanuts	
Southern peas	
Soybeans	
Alfalfa	4 Months after application of Imazethapyr 2 SL
Edible beans and peas ^{1*}	
Clover	
Rye ^{**}	
Wheat ¹	
Field corn ¹	8 1/2 Months after application of Imazethapyr 2 SL
Field corn grown for seed ¹	
Barley ¹	9 1/2 Months after application of Imazethapyr 2 SL
Tobacco	
Cotton ^{1***}	18 Months after application of Imazethapyr 2 SL
Lettuce	
Oats	
Popcorn	
Rye ^{****}	
Safflower	
Sorghum	
Sunflower	
Sweet Corn	
Flax	26 Months after application of Imazethapyr 2 SL
Potatoes	
All crops not listed elsewhere in this table ^{*****}	40 Months after application of Imazethapyr 2 SL

* Not Southern peas and lima beans

**Except in North Dakota and Minnesota north of Highway 210

***See below for details on a Cotton Rotation Interval after treatment with Imazethapyr 2 SL to clover or alfalfa grown for seed.

****In North Dakota and Minnesota north of Highway 210

*****Prior to planting crops not listed in the above table, 40 months after treatment with Imazethapyr 2 SL, a field bioassay must be completed consisting of the following: In the previously treated area, a test strip (which must contain variation in soil pH and soil type, as well as knolls and low areas) must be planted with the intended rotational crop and grown to maturity. If there is no crop injury in the test strip after the crops have reached maturity, the intended rotational crop may be planted in the previously treated area the next year.

¹ Refer to the Rotation Crops: Exceptions section for additional intervals.

The yield of some crops (e.g. sugar beets) may be reduced when soil pH is less than 6.5. If lime is applied to the area in order to adjust soil pH before rotational crops not listed in this section are planted, the lime treatment must take place a minimum of 12 months before the rotational crop is planted.

Rotational Crops: Exceptions

Crop	Use Directions
Barley	In Delaware, Indiana, Kentucky, Maryland, New Jersey, Ohio, Pennsylvania, and Virginia: Plant barley a minimum of 4 months after treatment with Imazethapyr 2 SL. In North Dakota: Plant barley a minimum of 18 months after treatment with Imazethapyr 2 SL. When Imazethapyr is applied to edible legumes at a rate not exceeding 3 fluid ounces per acre, do not plant barley for a minimum of 4 months after treatment.
Chickpeas, Lentils, Peas	When Imazethapyr is applied to edible legumes at a rate not exceeding 3 fluid ounces per acre, chickpeas, lentils and peas may be planted immediately after treatment.
Corn Inbred Lines	Plant corn inbred seed lines in the season following application of Imazethapyr 2 SL.*
Sweetcorn** Popcorn**	In Illinois, Indiana, Iowa, Minnesota, Ohio, Tennessee and Wisconsin: Plant sweetcorn and popcorn varieties in the season following application of Imazethapyr 2 SL. Some crop damage may occur in popcorn and sweetcorn planted within 18 months of treatment with this product. Restriction • Fresh market sweetcorn varieties must be planted a minimum of 18 months after an application of Imazethapyr 2 SL.
Bahiagrass, Cabbage, Cantaloupe, Cucumber, Irish potato, Onion, Sweet potato transplants, Sweet pepper transplants, Tomato transplants, Watermelon	In Alabama, Delaware, Florida, Georgia, Indiana, Kentucky, Maryland, New Jersey, North Carolina, Pennsylvania, South Carolina, and Virginia: Do not plant until a minimum of 18 months following an application with Imazethapyr 2 SL.

Crop	Use Directions
Cotton	<p>In North Carolina, South Carolina and Virginia: In areas where Imazethapyr 2 SL has been applied to peanuts in the previous season in sandy loam/loamy sand and where there has been more than 16 inches of irrigation/rain between treatment and October in the same year of application, cotton can be planted a minimum of 9 1/2 months following treatment.</p> <p>Cotton Rotation Interval Based on Precipitation/Irrigation</p> <p>The following are the crop rotation intervals following application of Imazethapyr 2 SL in clover or alfalfa grown for seed. The following guidance does not apply to clover/alfalfa grown for forage or hay.</p> <p>Where precipitation/irrigation requirements are less than 36" or 3 acre feet of water, crops may be planted 40 months after application of Imazethapyr 2 SL.</p> <p>Where precipitation/irrigation requirements are equal to or more than 36" or 3 acre feet of water, crops may be planted 18 months after application of Imazethapyr 2 SL.</p>
Field Corn, Field Corn Grown for Seed	<p>In Arizona, Hawaii, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming: Do not plant until a minimum of 9 1/2 months following an application with Imazethapyr 2 SL.</p>
Snap Beans	<p>When Imazethapyr is applied at a rate of 1.5 fluid ounces per acre or less, snap beans may be planted immediately after treatment.</p> <p>When Imazethapyr is applied to edible legumes at a rate not exceeding 3 fluid ounces per acre, do not plant snap beans for a minimum of 3 months after treatment.</p>
Wheat	<p>East of highway I-35: Do not plant wheat for a minimum of 3 months following application of Imazethapyr 2 SL.</p>

*Applicators must contact the seed company for directions on planting corn grown for seed in areas which have been treated with Imazethapyr 2 SL in the previous season. Crop damage is always a risk due to possible variations in and extremes of agronomic and environmental factors, as well as variations in agronomic practices. Planting seed corn inbreds into areas previously treated with Imazethapyr 2 SL is at the sole risk of the applicator.

**Contact the processor company/popcorn company for directions on crop tolerance when planning to plant popcorn or sweetcorn the season following an application of Imazethapyr 2 SL. Crop damage (delayed maturity/stunting) is always a risk due to possible variations in and extremes of agronomic and environmental factors, as well as variations in agronomic practices. Planting popcorn or sweetcorn varieties into areas previously treated with Imazethapyr 2 SL is at the sole risk of the applicator.

RESTRICTIONS (ALL CROPS)

Applications of the following products/products containing the following active ingredients at their full specified rates and made in the same year as an application of Imazethapyr 2 SL will increase the likelihood of crop damage to sensitive fallow crops. Consult all labels of product(s) used in combination/sequence with Imazethapyr 2 SL.

Product	Active Ingredient
Classic®	chlorimuron ethyl
Canopy® XL	
Synchrony®	
FirstRate™	chloransulam-methyl
Hornet™	flumetsulam
Python™	
Squadron®	imazaquin
Scepter® 70DG	
Pursuit® DG	imazethapyr
Pursuit® Plus EC	

- Only rotational crops that have been harvested at maturity can be used for food or feed.
- Soybeans, and peanuts can be replanted in the event of crop loss due to weather.
- Soil must not be worked to a depth greater than 2 inches.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store above 32°F in original containers only. If product freezes, return to room temperature and agitate to reconstitute. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (greater than five gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable Container: Refillable container. Refill this container with Imazethapyr 2 SL. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES and LIMITATIONS OF LIABILITY.**

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ADAMA. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ADAMA makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of ADAMA is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ADAMA disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at ADAMA's election, the replacement of product.

Counter® and Thimet® are a registered trademarks of AMVAC Chemical Corporation.

Banvel®, Basagran®, Clarity®, Marksman®, Outlook®, Poast®, Poast Plus®, Prowl®, Pursuit®, Scepter®, Squadron®, and TRI-4® are registered trademarks of BASF Ag Products.

Buctril®, Sencor® and Whip® are registered trademarks of Bayer CropScience.

Balan™, FirstRate™, Hornet™, Lorsban™, Python™ and Sonalan™ are trademarks of Dow AgroSciences LLC.

Accent®, Basis®, Canopy®, Classic®, Harmony®, Staple® and Synchrony® are registered trademarks of E.I. du Pont de Nemours and Company.

Command® and Authority®, are registered trademarks of FMC Corporation.

Lasso®, Roundup Ready® and Roundup Ultra® are registered trademarks of the Monsanto Company.

Bravo®, Gramoxone®, Flexstar® and Reflex® are registered trademarks and Spirit™ is a trademark of Syngenta Crop Protection, Inc.

Orthene® is a registered trademark of OMS Investments, Inc.

Ultra Blazer® and Storm® are registered trademarks of United Phosphorous, Inc.

Solubor® is a registered trademark of U.S. Borax & Chemical Corp.

Cobra®, Prism® and Select® are registered trademarks of Valent U.S.A. Corporation.

Manufactured for:

Makhteshim Agan of North America, Inc.

(d/b/a ADAMA)

3120 Highwoods Blvd., Suite 100

Raleigh, NC 27604

092516-1.0

Imazethapyr 2 SL

GROUP 2 HERBICIDE

Herbicide For use on Alfalfa, Clover, Peas, Beans, Peanuts and Soybeans

ACTIVE INGREDIENT: % BY WT.

Ammonium salt of imazethapyr (±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid* 22.87%

OTHER INGREDIENTS: 77.13%

TOTAL: 100.0%

*Equivalent to 21.6% or 2 pounds per U.S. gallon or 240 grams per liter of imazethapyr acid.

EPA Reg. No. 66222-248

EPA Est. No. 37429-GA-001⁹¹; 37429-GA-002⁹⁰; 37429-GA-003⁹¹
Letter(s) in lot number correspond(s) to superscript in EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

**How can we help?
1-866-406-6262**

Manufactured for:
Makhteshim Agan of North America, Inc.
(d/b/a ADAMA)
3120 Highwoods Blvd., Suite 100
Raleigh, NC 27604

Net Contents
1 gallon



HERBICIDE
ADAMA
ESSENTIALS

FIRST AID

IF ON SKIN OR CLOTHING Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF INHALED Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.

In case of spills, fire, leaks or accidents call 1-800-535-5053.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store above 32°F in original containers only. If product freezes, return to room temperature and agitate to reconstitute. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PEEL BACK BOOK HERE

15697-A
092516-1.0