

ETHABOXAM	GROUP	22	FUNGICIDE
IPCONAZOLE	GROUP	3	FUNGICIDE
METALAXYL	GROUP	4	FUNGICIDE



A FUNGICIDE SEED TREATMENT PRODUCT PROVIDING SYSTEMIC SEED AND SEEDLING PROTECTION AGAINST LISTED EARLY SEASON SEEDLING DISEASES OF SOYBEAN.

Active Ingredients	By Wt
Ethaboxam ¹	5.11%
Ipconazole ²	1.70%
Metalaxyl ³	1.36%
Other Ingredients	91.83%
Total	100.00%

- ¹ (RS)-N-(α-cyano-2-thenyl)-4-ethyl-2-(ethylamino)-1,3-thiazole-5-carboxamide
- ² (1RS,2SR,5RS;1RS,2SR,5SR)-2-(4-chlorobenzyl)-5-isopropyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol
- ³ methyl N-(methoxyacetyl)-N-2,6-xylyl-DL-alaninate

Intego® Fungicide Soybeans contains 0.452 lb ethaboxam per gallon, 0.150 lb ipconazole per gallon and 0.121 lb metalaxyl per gallon.

EPA Reg. No. 59639-238
EPA Est. 89332-GA-2

**KEEP OUT OF REACH OF CHILDREN
SEE BELOW FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.**

NET CONTENTS 14.98 GALLONS

HOT LINE NUMBER
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 800-892-0099 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Workers loading product, treating seed and conducting maintenance and calibration of the treater must wear: long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves made of any waterproof material.

Workers involved with handling treated seed (such as baggers, sewers and bag stackers) must wear: long-sleeved shirt, long pants, socks and shoes.

Workers who load and plant treated seed must wear: long-sleeved shirt, long pants, socks and shoes and chemical-resistant gloves made of any waterproof material when handling treated seed.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates, oysters and shrimp. Treated seed exposed on soil surface may be hazardous to birds, wildlife, fish and aquatic invertebrates. Cover or collect seed spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not apply directly to water or areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply where runoff is likely to occur. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on label.

Ipconazole may persist in soil, and long-term use in the same field may result in accumulation of active ingredient.

Groundwater Advisory

Ethaboxam has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Therefore, contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several days after application. A level, well maintained vegetative

buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential loading of ethaboxam from runoff water and sediment. Run-off of this product will be reduced by avoiding applications where rainfall or irrigation is expected to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

For use only in commercial seed treatment facilities. Do not use in hopper box, planter box, seed box, slurry box or other on-farm seed treatment applications. Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

APPLICATION INSTRUCTIONS

RESTRICTIONS

- *Intego* Fungicide Soybeans must be used only in accordance with use instructions on this label and only in commercial seed treatment facilities. Always mix product thoroughly before use. Do not use in hopper box, planter box, seed box, slurry box or other on-farm seed treatment applications.
- Do not use on soybeans in California.
- Do not use treated seed for food, feed or oil purposes.
- Care must be exercised in the handling of treated seed.
- Do not use augers used for handling treated seed to move seed for feed, food or oil processing.
- Do not re-use packaging from treated seed to handle food or feed products.
- Dispose of seed packaging or containers in accordance with local requirements.
- Do not apply more than the maximum seasonal total for the active ingredient as stated on the label containing the lowest seasonal total on that crop.

As stated by regulations specified in 40 CFR 153.155, all seed treated with *Intego* Fungicide Soybeans must be colored with an EPA-approved dye or colorant to visually distinguish seed that is treated with a pesticide. This product contains such dye or colorant. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.

MODE OF ACTION

Intego Fungicide Soybeans contains

Ethaboxam: a Group 22 fungicide as classified by the Fungicide Resistance Action Committee (FRAC). Its mode of action is the inhibition of cell division.

Ipconazole: a Group 3 fungicide as classified by FRAC. Its mode of action is the inhibition of sterol biosynthesis.

Metalaxyl: a Group 4 fungicide as classified by FRAC. Its mode of action is the inhibition of rRNA biosynthesis.

SEED QUALITY AND STORAGE OF TREATED SEED

Seed treatments applied to low vigor or poor quality seed or to mechanically damaged seed may result in loss of seed germination or reduced seedling vigor. To verify seed safety, treat a small portion of seed with the desired slurry mixture (*Intego* Fungicide Soybeans alone or in combination with other commercial seed treatment products) prior to treatment of an entire seed lot. Utilize standard seed germination methodologies to determine and verify seed safety. Use proper storage conditions for treated seed. Due to seed quality and seed storage conditions beyond the control of Valent U.S.A. LLC, no claims are made to guarantee the germination of carry-over treated seed.

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with *Intego* Fungicide Soybeans must be labeled with the following statements:

- This seed has been treated with *Intego* Fungicide Soybeans which contains ethaboxam, ipconazole and metalaxyl.
- Do not use treated seed for feed, food or oil processing.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

In addition, the U.S. Environmental Protection Agency requires the following statements on the container of seed treated with *Intego* Fungicide Soybeans:

- When handling treated seed (including opening the package, loading, or pouring) wear long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves made of any waterproof material.
- Care must be exercised in the handling of treated seed.
- Store treated seed away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seed.
- Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Treated seed exposed on soil surface may be hazardous to birds, wildlife, fish and aquatic invertebrates. Cover or collect seed spilled during loading and planting.

- Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging or containers in accordance with local requirements.
- This product contains metalaxyl which is known to leach through soil into groundwater under certain conditions as result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Ethaboxam has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- Do not apply more than the maximum seasonal total for the active ingredients as stated on the label containing the lowest seasonal total on that crop.
- Do not re-use packaging from treated seed to handle food or feed products.
- Do not use augers used for handling treated seed to move seed for feed, food or oil processing.
- Treated seed must be planted into the soil at a minimum depth of 0.5 inch.
- Ipconazole may persist in soil, and long-term use in the same field may result in accumulation of active ingredient.
- After seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: Once the seeds are planted in soil, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area without restriction if there will be no worker contact with the soil sub-surface or treated seed. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: long-sleeved shirt and long pants; chemical-resistant gloves made of any waterproof material; shoes plus socks.

PRODUCT INFORMATION

Intego Fungicide Soybeans (containing ethaboxam, ipconazole and metalaxyl) is a fungicide seed treatment product offering both contact and systemic activity in providing broad-spectrum seed and seedling protection against seedling diseases. *Intego* Fungicide Soybeans is effective for protection against *Pythium*, *Phytophthora sojae*, *Fusarium* and *Rhizoctonia solani*. This product is also effective against seed-borne *Phomopsis*, seed-borne *Sclerotinia* and seed rot fungi such as *Aspergillus* and *Penicillium*.

USE DIRECTIONS

Seed is to be sound and well-cured before treatment.

Accurate application of *Intego* Fungicide Soybeans to seed is required for best protection against listed diseases. Application to seed must be done only in commercial seed treatment facilities utilizing standard liquid or slurry treaters. Always mix product thoroughly before use.

Pre-test tank mixtures with other seed treatment products to verify formulation compatibility and to assure proper physical compatibility of products. Follow the most restrictive limitations and precautions on labels of all products used in a seed treatment tank mixture. Do not mix with any products which contain a prohibition on tank mixing.

SOYBEAN (Do not use on soybeans in California.)		
Seed/Seedling Diseases	<i>Intego</i> Fungicide Soybeans Use Rates	
	fl oz/ 100 lb seed	mg active ingredient per seed¹
Seed Rots (including seed-borne <i>Phomopsis</i> , seed-borne <i>Sclerotinia</i> , seed-borne and soil-borne <i>Fusarium</i> , <i>Penicillium</i> and <i>Aspergillus</i>) Damping Off and Seedling Blights <i>Pythium</i> seed decay and seedling dieback Early season <i>Phytophthora sojae</i> root rot Soil-borne <i>Fusarium</i> and <i>Rhizoctonia solani</i>	2.11	ethaboxam 0.012 mg ipconazole 0.004 mg metalaxyl 0.0032 mg

¹ mg active ingredient per seed is based on 2800 seeds per pound, thus delivering 0.0192 mg ai/seed total active ingredients.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE

Store in a dry place away from excessive heat.

Do not store near food or feed.

Store in original container only.

To close package, replace and tighten cap to form an airtight seal.

Do not freeze.

PESTICIDE DISPOSAL

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING

Empty the remaining contents into application equipment or a mix tank. Add water – at least 2% of the container volume, up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

RECYCLING

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 877-952-2272 (toll free) or www.acrecycle.org.

DISCLAIMER, RISKS OF USING THIS PRODUCT, LIMITED WARRANTY AND LIMITATION OF LIABILITY

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

(continued)

(continued)

ASSUMPTION OF RISK

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests to this product. Such risks of crop injury, non-performance, or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from

(continued)

(continued)

the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

COMMERCIAL TANK MIXES

NOTICE: Commercial tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the commercial tank mix with this product.

©2019 Valent U.S.A. LLC

Manufactured for:

Valent U.S.A. LLC

P.O. Box 8025

Walnut Creek CA 94596-8025

Made in U.S.A.

Form 2251-A

EPA Reg. No. 59639-238

EPA Est. 89332-GA-2

059639-00238.20190924.V10442_0.72FS_NewProd.FINAL
SAL20190925

Information contained in this booklet is accurate at the time of printing. Since product testing is a continuous process, please read and follow the directions on the product label for the most current directions and precautionary statements.

Always check with your state to verify state registration status or call 800-6-VALENT (682-5368).



For state registration and/or supplemental labels, please call or visit us online.

Products That Work, From People Who Care® | valent.com | 800-6-VALENT (682-5368)

Always read and follow label instructions.

©2019 Valent U.S.A. LLC. All rights reserved. Printed in the U.S.A.