

# IVC 415 SUPREME SPRAY OIL

## Insecticide - Miticide 415 Type Oil



## INSECTICIDE

<b>ACTIVE INGREDIENT:</b>	<b>BY WT.</b>
Mineral Oil .....	99.70%
<b>INERT INGREDIENTS</b> .....	<b>0.30%</b>
<b>TOTAL</b> .....	<b>100.00%</b>

Contains petroleum distillates  
Petroleum Oil Unsulfonated Residue - 92%  
Petroleum Oil Classification - Light Medium

## KEEP OUT OF REACH OF CHILDREN CAUTION

See Below For Additional Precautionary Statements

FIRST AID	
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>Immediately call a poison control center or doctor.</li><li><b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li><b>DO NOT</b> give ANY liquid to a person.</li><li><b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>Take off contaminated clothing.</li><li>Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>Move person to fresh air.</li><li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>Call a poison control center or doctor for further treatment advice.</li></ul>
<b>HOTLINE NUMBER:</b> Have the product container or label with you when calling a poison control center or doctor or going for treatment. Call Chemtrec, 1-800-424-9300.	
<b>NOTE TO PHYSICIAN:</b> Contains petroleum distillates. Vomiting may pose aspiration pneumonia.	

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, nitrile rubber, neoprene rubber or viton. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

**Mixers, loaders, applicators, flaggers, and other handlers must wear:** long-sleeved shirt and long pants, shoes plus socks, and chemical resistant gloves. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. **DO NOT** reuse them.

### ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, or enclosed cabs in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

**Users should:**  
Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

**DO NOT** discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. **DO NOT** discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas

### PHYSICAL OR CHEMICAL HAZARDS

**DO NOT** weld, heat or drill this container. Residues may ignite with explosive violence if heated sufficiently.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. **DO NOT** enter or allow workers entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as soil, or water, is coveralls, shoes plus socks, and chemical resistant gloves made of any waterproof material.

### MIXING DIRECTIONS

Check to make sure this product will emulsify. If the mixture is not milky after agitation, **DO NOT** use this product. With the agitator running, start filling the spray tank with water. Add wettable powders or other products when the tank is one third full and always before adding **IVC 415 SUPREME SPRAY OIL**. When the tank is one half full and with good agitation, add the proper amount of **IVC 415 SUPREME SPRAY OIL**, then fill to the desired volume. Maintain agitation during mixing and application.

### PRECAUTIONS AND RESTRICTIONS

- DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift.
- Only protected handlers may be in the area during application.
- DO NOT** apply this product through any type of irrigation system.
- DO NOT** apply by aerial application.
- DO NOT** apply if this product will drift to food forage or other crops that might be damaged.
- For applications, the spray tank must be free of any sulfur residue before using **IVC 415 SUPREME SPRAY OIL**. **DO NOT** combine this product with any type of sulfur spray and **DO NOT** follow a previous application of sulfur or apply sulfur after **IVC 415 SUPREME SPRAY OIL** until sufficient time has passed for safety (at least 60 days). If doubts exist, spray a **IVC 415 SUPREME SPRAY OIL** mixture on a portion of one tree that was treated with sulfur. If no burn is noted in several days, enough time has passed. **DO NOT** spray when temperatures are above 90 °F or when temperatures drop below 35 °F. **DO NOT** use on foliage closer than 60 days after or 90 days before using Captan or Folpet products.
- IVC 415 SUPREME SPRAY OIL** must be diluted before an application is made
- Follow all precautions and instructions on all pesticide labels being tank mixed.
- All rates are for mature trees; rates must be proportionately adjusted downward for smaller or immature trees.
- Apply by ground application from a sprayer that provides thorough coverage for best efficacy. For dilute sprays, spray for thorough coverage without significant runoff.
- Application to trees weakened by disease, drought, drying winds or high nitrogen applications may result in injury to the trees.
- See your Innvictis Fieldman of Agricultural Extension Service for recommendations on timing and combinations with **IVC 415 SUPREME SPRAY OIL**.

### SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind directions, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product.

#### Ground Application

- Release Height: Apply using a nozzle height of no more than 4 feet above the ground or crop canopy.
- Wind Speed: **DO NOT** apply at wind speeds greater than 15 mph at the application site.
- Droplet Size: Apply as a medium or coarser spray (ASAE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

#### Airblast:

For airblast applications, turn off outward pointing nozzles at row ends and when spraying outer row. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

#### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. **DO NOT** make applications into areas of temperature inversions or stable atmospheric conditions.

### FOLIAR APPLICATION - DILUTE SPRAY

**ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES:** Brown Apricot Scale, Black Scale, San Jose Scale, Red Spider Mites - Use 1 to 1½ gallons per 100 gallons of water at 300 to 600 gallons per acre.

**CAUTION:** Certain varieties of Plums and Prunes may be injured by oil sprays, check for tolerance to oil before Treatment.

**APPLES:** Rust Mites, Red Spider Mites, San Jose Scale - Use 1 gallon per 100 gallons of water at 400 to 800 gallons per acre.

**AVOCADOS:** Avocado Brown Mite, Persea Mite - Use 1 gallon per 100 gallons of water with thorough coverage through late fall. **DO NOT** exceed 30 gallons of oil per acre per application.

**GRAPES:** Mealybug - Post Harvest spray only. Use 4 to 6 gallons in 200 to 300 gallons of water per acre.

**ORANGES, LEMONS, GRAPEFRUIT, AND OTHER CITRUS VARIETIES:** Red Scale, Yellow Scale, Citricola Scale, Brown Soft Scale, Purple Scale, Black Scale, Citrus Red Mite, Pacific Spider Mite, Two Spotted Spider Mite - Use 1 to 1½ gallons per 100 gallons of water at 400 to 2000 gallons per acre. **DO NOT** use within 8 weeks of harvest for lemons. Maximum of 159 pounds active ingredient per acre (maximum of 1500 gallons of spray mix per acre) in Texas and Florida, and 212 pounds of active ingredient per acre (maximum of 2000 gallons of spray mix per acre) in California.

**OLIVES:** Olive Scale, Spider Mites, and Oleander Scale - Use 1½ gal. per 100 gal. of water at 400 to 1200 gallons per acre. Apply prebloom or postbloom up to buckshot. Black Scale (Calif. Only) - Use 1 to 1½ gal. per 100 gal. of water at 400 to 1200 gal. per acre. Make applications between July 1<sup>st</sup> and August 31<sup>st</sup>. Phytotoxicity can occur at high concentrations on stressed trees or at high temperatures.

**PEARS:** Red Spider Mites, San Jose Scale, Italian Pear Scale, Pear Psylla - Use 1 to 1½ gallons per 100 gallons water at 400 to 800 gallons per acre. Blister Mite - Use 1 to 1½ gallons per 100 gallons water at 400 to 800 gallons per acre as a fall spray after harvest only. **WALNUTS:** European Fruit Scale, Frosted Scale, San Jose Scale, Soft Shelled Scales, Red Spider Mites - Use 4 to 6 gallons in 200 gallons of water per acre. Apply at late spring to mid summer. **DO NOT** apply after husk split.

### FOLIAR APPLICATION - CONCENTRATE SPRAY

**NOTE-** Concentrate sprays using 50 to 100 gallons of Spray Solution per acre is preferred but not required depending on the performance of the sprayer. **IVC 415 SUPREME SPRAY OIL** must be diluted before an application is made.

**ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES:** Red Spider Mite, Brown Spider Mite, Pacific Spider Mite, Two Spotted Spider Mite- Use 4 to 6 gallons per acre in a minimum of 20 gallons of water. PEACHES, NECTARINES, ALMONDS and PLUMS - At least 2 days before harvest, use the lower concentrate rate of 4 gallons per acre. For PRUNES and APRICOTS - At least one week before harvest, use the lower concentrate rate of 2 gallons per acre in a minimum of 20 gallons per acre.

**APPLES, PEARS:** Two Spotted Spider Mites, Red Spider Mites - Use not more than 4 gallons per acre in a minimum of 20 gallons of water per acre. At least one week before harvest, use the lower concentrate rate of 2 gallons per acre in a minimum of 20 gallons per acre.

**AVOCADOS:** Avocado Brown Mite, Persea Mite - Use 1 to 18 gallons per 100 gallons of water per acre.

**ORANGES, LEMONS, GRAPEFRUIT, AND OTHER CITRUS VARIETIES:** Red Scale, Yellow Scale, Citricola Scale, Brown Soft Scale, Purple Scale, Black Scale, Citrus Red Mite, Pacific Spider Mite, Two Spotted Spider Mite – Use 5 to 15 gallons per acre in a minimum of 50 gallons of water per acre. **DO NOT** use within 8 weeks of harvest for lemons. Maximum of 159 pounds active ingredient per acre (maximum of 1500 gallons of spray mix per acre) in Texas and Florida, and 212 pounds of active ingredient per acre (maximum of 2000 gallons of spray mix per acre) in California.

### DORMANT OR DELAYED DORMANT - DILUTE SPRAY

**ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS PRUNES:** Brown Apricot Scale, Black Scale, San Jose Scale, Olive Scale, Oleander Scale, Frosted Scale, Italian Pear Scale, European Fruit Lecanium, Putnam Scale, Soft Brown Scale, Eggs of Aphids, and Spider Mites - Use 1½ to 2 gallons per 100 gallons of water at 300 to 600 gallons per acre. **CAUTION:** Spray Plums and Prunes during the delayed dormant period only. Oil sensitive varieties may develop dead limbs and scaffolds that will reduce yields.

**APPLES:** San Jose Scale, Eggs of Aphids, Red Spider Mites - Use 1½ to 2 gallons per 100 gallons of water at 400 to 800 gallons per acre. Rust Mites - Use 1½ to 2 gallons per 100 gallons of water at 400 to 800 gallons per acre for delay dormant spray only.

**FIGS:** Fig Scale - Use 3 gallons per 100 gallons of water at 300 to 600 gallons per acre.

**GRAPES:** Mealybug - Use 4 to 6 gallons in 200 to 300 gallons of water per acre as a dormant spray only.

**PEARS:** San Jose Scale, Italian Pear Scale, Eggs of Aphids, and Red Spider Mites - Use 1½ to 2 gal. in 100 gallons of water at 400 to 800 gallons per acre. Pear Psylla and Blister Mite - Use 1½ to 2 gal. in 100 gallons of water at 400 to 800 per acre as a delayed dormant spray only.

### DORMANT OR DELAYED DORMANT - CONCENTRATE SPRAY

**NOTE -** Concentrate sprays using 50 to 100 gal. of spray solution per acre is preferred but not required depending on the performance of the sprayer. **IVC 415 SUPREME SPRAY OIL** must be diluted before an application is made.

**ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES:** Brown Apricot Scale, Black Scale, San Jose Scale, Olive Scale, Oleander Scale, Frosted Scale, Italian Pear Scale, European Fruit Lecanium, Putnam Scale, Soft Brown Scale, Eggs of Aphids, and Spider Mites - Use 6 to 8 gallons per acre in a minimum of 20 gallons of water.

**PLUMS, PRUNES:** Black Scale, Brown Apricot Scale, Olive Scale, San Jose Scale, Brown Mite Eggs, Eggs of European Red Mites, Eggs of Mealy Plum Aphids, Plum Aphid Eggs - Use 3 to 5 gallons per acre in a minimum of 20 gallons of water. **CAUTION:** Spray Plums and Prunes during the delayed dormant period only. Oil sensitive varieties may develop dead limbs and scaffolds that will reduce yields.

**APPLES, PEARS:** San Jose Scale, Red Spider Mite Eggs, Pear Psylla - Use 4 to 8 gallons per acre in minimum of 20 gallons of water. **DO NOT** apply to oil sensitive varieties as phytotoxicity to buds and new wood may result.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep pesticide in original container. **DO NOT** put concentrate into food or drink containers. Protect from water contamination during storage and transfer. Contamination with water may make this product unfit for its intended use. Always check emulsification before using. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

#### CONTAINER HANDLING:

Nonrefillable (5 and 30 gallon containers). **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on to its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If recycling is not available, puncture container and dispose of it in a sanitary landfill or by other procedures approved by State and local authorities.

(cont'd)

## STORAGE AND DISPOSAL (cont'd)

**Refillable (Bulk and trailer tank containers greater than 30 gallons):** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of INNICTIS ORGANICS ADVANCEMENTS or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold INNICTIS ORGANICS ADVANCEMENTS and Seller harmless for any claims relating to such factors.

INNICTIS ORGANICS ADVANCEMENTS warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or INNICTIS ORGANICS ADVANCEMENTS, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, INNICTIS ORGANICS ADVANCEMENTS MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither INNICTIS ORGANICS ADVANCEMENTS nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF INNICTIS ORGANICS ADVANCEMENTS AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF INNICTIS ORGANICS ADVANCEMENTS OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

INNICTIS ORGANICS ADVANCEMENTS and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of INNICTIS ORGANICS ADVANCEMENTS.



EPA Reg. No.: 89168-143-96463

EPA Est. No.:

NET CONTENTS: 5 Gal (18.9 L)



8 10100 40096 5



10810100400962



Distributed By:  
INNICTIS ORGANIC ADVANCEMENTS  
P.O. Box 9296  
Boise, ID 83707

021821R0061522