KRISS™

5-8-4

GUARANTEED ANALYSIS

Total Nitrogen (N)	5.0%
1.7% Ammoniacal Nitrogen	
1.0% Nitrate Nitrogen	
1.7% Urea Nitrogen	
0.6% Water Insoluble Nitrogen	
Available Phosphate (P ₂ O ₅)	8.0%
Soluble Potash (K ₂ O)	4.0%

Derived from: Urea, Monoammonium phosphate, Potassium nitrate, Yeast Extract

SHAKE WELL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

CAUTION

While this product is not anticipated to be hazardous for use, good industrial hygiene practices should be used in handling the product. Wash hands thoroughly after handling and take appropriate precautions to avoid eye and skin contact with the product. Keep away from flames in a cool, well-ventilated place. Avoid prolonged exposure to low temperatures. Check container periodically. In case of accidental spill contain the leak, avoid water contamination and, whenever necessary, dispose according to federal, state and local laws and regulations.

DIRECTIONS FOR USE

Foliar application:

- Alfalfa: Apply 16 ounces/acre at establishment, and after 1st and 2nd cutting for a total of 3 applications.
- **Blueberries:** Apply 10 30 ounces/acre. Apply 2 3 times per season.
- Citrus: Apply 2.6 pints/acre every 12 -15 days beginning after fruit set for a total of 3 applications.
- Cucumber, zucchini: Apply 1.5 2.0 mL per liter of water every 5 to 7 days throughout the entire crop cycle.
- Corn: Apply 1.0 pint/acre between V4 and V8. DO NOT apply this product during or after VT.
- Dry Beans, Lentils and Peas: Apply 16 ounces/acre at first initiation of bloom.
- Eggplant, pepper, tomato: Apply 1.5 2.0 mL per liter of water every 5 to 7 days after fruit set of the first truss.
- Grape (Table and Wine): Apply 2.6 pints/acre every 8 to 10 days beginning after fruit set for a total of 2 applications.
- **Melon**, **watermelon**: Apply 1.7 pints/acre every 6 to 7 days beginning after fruit set for a total of 3 applications.
- **Mint:** Apply 16 ounces/acre at establishment, and after 1st and 2nd cutting for a total of 3 applications.
- Pome and stone fruit: Apply 2.6 pints/acre every 6 to 7 days beginning at complete petal fall for a total of 2 applications.
- **Processing tomato:** Apply 1.7 pints/acre every 6 to 7 days beginning after fruit set for a total of 3 applications.
- **Strawberries:** Apply 10 30 ounces/acre. Apply 2 3 times per season.
- Soybeans: Apply 1.0 pint/acre between R1 and R3.

Dosages are calculated for normal water distribution volumes.

F1683

Distributed and Guaranteed By: Winfield Solutions, LLC P.O. Box 64589 St. Paul, MN 55164 2/0210/0 Weight Per Gallon: 10.2 lbs (4.63 kg) NET WEIGHT: 25.50 lbs. (11.57 kg) NET CONTENTS: 2.5 Gallons (9.46 Liters)



Please consult your regional Winfield agronomist for additional information on application rates, crops not listed and use rates. The product can be mixed with all common formulations, except with products with products based on Copper. A simple mixture test to check compatibility is recommended.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html

WARRANTY DISCLAIMER

Winfield Solutions, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labelled use when applied according to the Directions for Use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.

Kriss™ is a trademark of Biolchim S.p.A