


# Specimen Label



®Trademark of Dow AgroSciences LLC

## A systemic, protectant and curative fungicide for disease control in the commercial production of almonds

### Active Ingredient

myclobutanil:  $\alpha$ -butyl- $\alpha$ -(4-chlorophenyl)-

1H-1,2,4-triazole-1-propanenitrile ..... 19.7%

Other Ingredients ..... 80.3%

Total..... 100.0%

Contains petroleum distillates

Contains 1.67 lb of active ingredient per gallon

EPA Reg. No. 62719-493

### Keep Out of Reach of Children

## CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

**Causes Moderate Eye Irritation • Harmful If Swallowed Or Absorbed Through The Skin**

**Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.**

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made from any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or on clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**Note to Physician:** This product may pose an aspiration pneumonia hazard. Contains petroleum distillates.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

### Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

**Notice:** Read the entire label. Use only according to label directions.

**Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

### Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made from any waterproof material
- Shoes plus socks

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal

**Pesticide Storage:** Store in a cool, dry area at temperatures above freezing.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### Nonrefillable containers 5 gallons or less:

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

### Storage and Disposal (Cont.)

#### Refillable containers 5 gallons or larger:

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

#### Nonrefillable containers 5 gallons or larger:

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times.

Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

### General Information

Laredo® EW fungicide is a systemic, protectant and curative fungicide for disease control in the commercial production of almonds. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program.

### Use Rate Determination

Carefully read and follow label directions, including recommended use rates and restrictions. For proper application, determine the size of the area to be treated, the recommended label application rate and the required gallonage for the area to be treated. Prepare only the amount of spray solution needed for the treatment area. Careful calibration of spray equipment is recommended prior to use.

### Mixing

Be sure sprayer is clean and not contaminated with other materials prior to use. Fill the spray tank 1/4 to 1/2 full with clean water and start agitation. Be certain the agitation system is working properly and creates a rolling or rippling effect on the liquid surface. With the agitator running, add the required amount of Laredo EW to the tank and finish filling tank. Always add Laredo EW to the spray tank before adding other materials.

**Compatibility:** Laredo EW is compatible with most commonly used fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. When preparing tank mixtures, spray compatibility charts or State Cooperative Extension Service Specialist should be consulted prior to use.

**Application**

**Ground:** Thorough coverage sprays generally result in optimum disease control. To achieve good coverage use proper spray pressure, gallonage per acre, nozzles, nozzle spacing and tractor speed. Consult manufacturer’s spray nozzle and accessory catalogues for specific information on proper equipment calibration.

**Aerial:** Apply in a minimum of 10 gallons of water per acre. Avoid application under conditions when uniform coverage cannot be obtained or when spray drift may occur.

**Hand Pressurized Sprayers:** For best control of labeled diseases, achieve thorough coverage of all plant parts on a protectant application schedule.

**Almond**

Best disease control is achieved in thorough coverage sprays applied on a protectant schedule that does not exceed 10 days.

Diseases	Laredo EW		Specific Use Instructions	Restrictions
	fl oz/100 gal <sup>1</sup>	fl oz/acre		
blossom blight ( <i>Monilinia</i> spp.) shothole ( <i>Stigmina</i> spp.)	2.4 - 3.8 (0.5 - 0.8 oz active)	9.6 - 15.3 (2 - 3.2 oz active)	Begin applications at pink bud stage (about 5% bloom). If conditions are favorable for disease development, apply again at full bloom and at petal fall.	<ul style="list-style-type: none"> <li>Do not apply within 90 days of harvest.</li> <li>Do not make more than three applications or apply more than 46.0 fl oz of Laredo EW (0.6 lb active) per acre per season.</li> </ul>
rust ( <i>Tranzschelia</i> spp.)			Apply 11.5 fl oz (2.4 fl oz active) per acre. Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development. For optimum disease control, do not apply on an application schedule exceeding 21 days.	
anthracnose ( <i>Colletotrichum</i> spp.)			The application of Laredo EW for control of blossom blight and shothole will suppress anthracnose.	

<sup>1</sup>Rate is based on standard dilute spray volume of 400 gallons per acre, or an equivalent amount of product per acre.

**Terms and Conditions of Use**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

**Warranty Disclaimer**

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

**Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

---

## Limitation of Remedies

---

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC

**Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.**

Label Code: D02-196-003

Replaces Label: D02-196-002

LOES Number: 010-02074

EPA accepted 07/09/04

### Revisions:

1. Updated storage and disposal instructions.