



**Dow AgroSciences**

# Lontrel™ 72

## Herbicide

GROUP	4	HERBICIDE
-------	---	-----------

For control of perennial and annual broadleaved weeds in field crops, Christmas tree plantations, pasture, rangeland, vegetable and fruit crops, and non-cropland.

COMMERCIAL

READ THE LABEL AND BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

GUARANTEE: clopyralid, present as monoethanolamine salt: 72.0 %  
Soluble granules

REGISTRATION NO. 31039 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 kg - bulk

**Dow AgroSciences Canada Inc.**  
Suite 2100, 450 - 1 Street S.W.  
Calgary, Alberta  
T2P 5H1  
1-800-667-3852

®™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

## **PRECAUTIONS**

### **KEEP OUT OF REACH OF CHILDREN**

Avoid contact with eyes, skin and clothing. Avoid breathing dust or spray mist.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

Workers must wear long pants, a long-sleeved shirt and chemical-resistant gloves. Goggles or a face shield are required during mixing and loading. Gloves are not required to be worn during groundboom applications but are required for mixing/loading, clean-up and repair.

Do not enter or allow workers entry into treated areas for 12 hours following application to all crops.

Do not use in greenhouses.

Apply only when the potential for drift to areas of human habitation or areas of human activity (such as houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

**AT COMPLETION OF SPRAYING OR END OF THE DAY:** Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.

## **FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

## **TOXICOLOGICAL INFORMATION**

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

## **AGRICULTURAL CHEMICAL**

Do not ship or store with food, feeds, drugs or clothing.

## **ENVIRONMENTAL HAZARDS**

TOXIC to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE. The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted or fine textured such as clay).

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

## **STORAGE**

Store away from food, feedstuff, fertilizer, seeds, insecticides, fungicides or other pesticides or herbicides intended to be used on crops sensitive to Lontrel 72 Herbicide.

## **DISPOSAL**

### **Recyclable Containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

### **Returnable Containers:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## **GENERAL INFORMATION**

Lontrel 72 Herbicide is a soluble granular herbicide intended for dilution with water and for use on canola, sugar beets, rutabagas, summerfallow, flax (including low linolenic acid varieties), wheat (spring), barley (spring), oats, strawberry, seedling and established grasses grown for forage and seed production, non-crop farmland, balsam fir Christmas trees and highbush blueberry. It is readily absorbed by both foliage and roots and translocates both upwards and downwards in plants. The product controls Canada thistle, wild buckwheat, scentless chamomile, common groundsel and volunteer alfalfa. It suppresses growth of perennial sow-thistle through control of top growth.

## **DIRECTION FOR USE**

### **VEGETATION AND CROP PRECAUTIONS**

#### **Sensitive Plants**

Do not apply Lontrel 72 Herbicide directly to, or otherwise permit it to come into contact with sunflowers, legumes (such as peas, beans, lentils or alfalfa), fruit or vegetable crops, flowers or other desirable broadleaved plants. Take precautions to prevent spray mists containing it to drift onto them. Residues of Lontrel 72 Herbicide can remain in the soil following the year of use, thereby affecting growth of sensitive crops.

Special precautions should be taken during application to non cropland areas such as roadsides, pipelines and railways where sensitive desirable vegetation may be present. Do not apply to or allow drift to come into contact with sensitive desirable vegetation such as vetch or clover which may be found on embankments.

### **Non-Target Sites**

Avoid contamination of non-target land, water or irrigation ditches. Do not use Lontrel 72 Herbicide in the following areas: standing or flowing water; the inner banks or bottoms of irrigation ditches; in areas where surface water can run off to adjacent croplands either planted or to be planted to sensitive crops.

### **Crop Rotation**

Fields previously treated with Lontrel 72 Herbicide can be seeded the following year to wheat, oats, barley, rye (not underseeded with legumes, clover or alfalfa), forage grasses, flax, canola, mustard or it can be summerfallowed.

Do not seed to crops other than those listed above in the calendar year following treatment.

### **Tank Mixtures**

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Dow AgroSciences Canada Inc at 1-800-667-3852 or [www.dowagro.ca](http://www.dowagro.ca) for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by Dow AgroSciences Canada Inc.

### **Grazing and Harvesting for Feed**

There are no restrictions on the grazing of crops or forages treated with Lontrel 72 Herbicide. If necessary, treated areas may be grazed immediately following application.

### **Manure and Straw**

Residues of the herbicide occurring in the straw may be harmful to susceptible plants; therefore, do not use straw or crop residue from treated crops for composting or mulching susceptible broadleaved crops. If the straw or crop residue is used for animal bedding or feed, return the manure to fields to be planted to clopyralid tolerant crops such as wheat, barley, oats, rye, forage grasses, canola or flax. Do not grow susceptible crops such as peas, beans, lentils, potatoes, sunflowers or other sensitive crops on land which has been mulched with straw containing Lontrel 72 Herbicide residues within the last 12 months.

### **SPRAY EQUIPMENT AND CONTAINER PRECAUTIONS**

**Field sprayer application:** **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

### **DO NOT APPLY BY AIR.**

To Reduce Spray Drift:

1. Use nozzles that deliver higher volumes and coarser droplets.
2. Use low pressures (200 to 275 kPa).
3. Use 100 to 200 L/ha of spray solution.
4. Spray when the wind velocity is 15 km/hr or less.

### **Equipment Clean-Up**

Equipment used to apply Lontrel 72 Herbicide should not be used to apply other pesticides to sensitive crops without thorough cleaning. Contact your Lontrel 72 Herbicide dealer for a detailed equipment cleaning procedure.

## APPLICATION DIRECTIONS

### Spray Preparation

To prepare the spray solution add about half the desired amount of water to the spray tank, then with mechanical or bypass agitation, add the recommended amount of Lontrel 72 Herbicide. Mix thoroughly in the tank. Second, add the recommended tank-mix herbicide. Finally with continued agitation, add the rest of the water.

### Spray Application Volume

Apply Lontrel 72 Herbicide at 140 to 415 g/ha plus any other herbicide approved as a tank-mix at the recommended rate in sufficient water to ensure thorough coverage (100 to 200 L/ha of spray solution) by ground equipment only at pressures of 200 to 275 kPa. Treat when weeds are young and actively growing, when the Canada thistle is in the rosette to pre-bud stage and before the purple bud stage and volunteer alfalfa is 5-50 cm in height.

For spot spraying of weed patches, mix the required amount of Lontrel 72 Herbicide in 200 L of water and apply to 1000 m<sup>2</sup> of weeds. Refer to the following table for the correct amount of Lontrel 72 Herbicide to use:

Recommended Rate of Lontrel 72 Herbicide Required Per Hectare	Amount of Lontrel 72 Herbicide Required To Treat 1000 m <sup>2</sup>
140 g	14 g
210 g	21 g
280 g	28 g
415 g	41.5 g

#### Approximate Conversions:

200 to 275 kPa = 30 to 40 PSI

100 to 200 L/ha = 10 to 20 gallons/acre

1 sq. metre = 1.2 sq. yards

100 g/ha = 1.43 oz/acre

## DIRECTIONS FOR USE – Lontrel 72 Herbicide Applied Alone

### CROPLAND AND NON-CROP FARMLAND AREAS

Weeds Controlled	Rate <sup>♦</sup>
Canada thistle (top growth) vetch ( <i>Vicia</i> spp.) alsike clover	210 g/ha
Canada thistle scentless chamomile wild buckwheat perennial sow-thistle (top growth) common groundsel volunteer alfalfa common ragweed sheep sorrel (suppression) ox-eye daisy (suppression)	280 g/ha
Canada thistle (season-long control of top growth with a reduction in population in the following year)	415 g/ha

<sup>♦</sup> Refer to individual crop sections below for appropriate use rate.

### Weed Stages at Application

Applications of Lontrel 72 Herbicide should be made when Canada thistle, perennial sow-thistle and scentless chamomile are in the rosette to pre-bud stage of growth. Best results are obtained when Canada thistle is actively growing and soil moisture is adequate for rapid growth. Under dry soil conditions and poor growing conditions, control of Canada thistle may be severely reduced. Applications of Lontrel 72 Herbicide made after the Canada thistle flower has reached the purple bud stage will not provide satisfactory control.

### Control of Canada Thistle

For in crop control of top growth of Canada thistle apply Lontrel 72 Herbicide at the rate of 210 g/ha. This will suppress top growth of Canada thistle for 6 to 8 weeks. Some regrowth may occur by the end of the season but this will not interfere with the harvesting of the crop.

For season long control of top growth of Canada thistle apply Lontrel 72 Herbicide at the rate of 280 g/ha. This rate will generally provide season long control of Canada thistle. Not all rhizomes will be killed and some regrowth may occur by the end of the growing season.

For season long control of top growth, with a reduction of Canada thistle population in the following year, apply Lontrel 72 Herbicide at the rate of 415 g/ha. This rate will provide season long control of Canada thistle and suppression into the following season, resulting in a reduction of the total number of Canada thistle shoots in the treated area.

### CANOLA (Western Canada Only)

For use on Polish and Argentine varieties, including canola. Lontrel 72 Herbicide should be diluted with water and applied at the 2 to 6 leaf stage of the crop to effectively control Canada thistle, scentless chamomile, common groundsel, wild buckwheat, the top growth of perennial sow-thistle and volunteer alfalfa. For specific directions for control of Canada thistle only refer to the section entitled Control of Canada Thistle.

### Tank-mix Combinations in Canola

REFER TO THE PRODUCT LABELS OF THESE HERBICIDES FOR A LIST OF OTHER WEEDS CONTROLLED, RATES (IF NOT LISTED IN THE TABLE BELOW) AND TIMINGS OF APPLICATION, WATER VOLUMES AND USE PRECAUTIONS.

Herbicide Tank-Mix Partner	Rate Lontrel 72 Herbicide	Rate Tank-Mix Partner	Additional Weeds Controlled
Poast Ultra	210 – 415 g/ha	0.32 – 0.47 L/ha + Merge Adjuvant 0.75-1.0 L/ha	annual grass weeds, Canada thistle
Venture L Herbicide	210 – 415 g/ha	0.6 – 2.0 L/ha	annual grass weeds, Canada thistle
Select	210 – 415 g/ha	0.19 L/ha + Amigo Adjuvant at 0.5% v/v	Canada thistle <sup>***</sup> , wild buckwheat <sup>****</sup> , wild oats, green foxtail, volunteer barley, volunteer wheat, and volunteer oats
Odyssey <sup>®</sup> WDG <sup>*</sup>	210 – 280 g/ha	29 – 43 g/ha	Canada thistle (Lontrel 72 Herbicide at 210 g/ha will provide top growth control of Canada thistle for 6-8 weeks, while the 280 g/ha rate will provide season long control of top growth)

Vantage™ Plus MAX II**	140 g/ha	0.94 L/ha	Annual Grasses: wild oats, green foxtail, volunteer barley, volunteer wheat
Roundup Transorb HC Liquid **		0.83 L/ha	
Roundup WeatherMAX With Transorb 2 Technology**		0.83 L/ha	Annual Broadleaves: wild buckwheat
			For rates and timing of application for annual grass and broadleaf weeds please see tank- mix partner label.
			wild Buckwheat and Perennial Weeds (season long control): quackgrass, Canada thistle (top growth), perennial sowthistle (season- long top growth), dandelion <15 cm diameter (season- long top growth), dandelion >15 cm diameter (suppression)

\*Clearfield canola varieties only - apply to Clearfield canola when in the 2 to 6 leaf stage and Canada thistle is actively growing.

\*\*Glyphosate-tolerant canola varieties only - apply to canola when in the 2 to 6 leaf stage. Use 100 L/ha water.

\*\*\*Canada thistle – 210 g Lontrel 72 Herbicide/ha top growth control to 6 - 8 weeks. 280 g Lontrel 72 Herbicide/ha for season-long control, 415 g Lontrel 72 Herbicide/ha control into following year.

\*\*\*\*Wild buckwheat - 280 g Lontrel 72 Herbicide/ha for season-long control.

### Tank-Mix Instructions

**Note 1:** When tank mixing water soluble formulations such as Lontrel 72 Herbicide with emulsifiable concentrates such as Poast Ultra and Select herbicides, first add the Lontrel 72 Herbicide to the spray tank. Once it is half filled with water, add the emulsifiable concentrate as the remaining water is put into the spray tank.

**Note 2:** If the sprayer has been previously used to apply herbicides which contain 2,4-D or MCPA herbicides, it is imperative that the spray equipment be thoroughly cleaned before Lontrel 72 Herbicide is mixed in the spray tank. Trace contamination of the spray solution with these herbicides will result in damage to the canola.

**Note 3:** Use 100 L/ha of water. Use a 50 mesh (or coarser) filter screen. Fill the spray tank three-quarters full with water. Add the required amount of Odyssey WDG herbicide soluble bag(s) directly into the sprayer through the tank opening. Agitate for at least ten minutes to dissolve the herbicide. After the herbicide is dissolved, use a separate calibrated measuring device to add the required amount of Lontrel 72 Herbicide while agitating the spray solution. After the Lontrel 72 Herbicide is dissolved, continue agitation and add the required amount of Merge adjuvant or non-ionic surfactant plus fertilizer. If excess foaming occurs, a silicone anti-foaming agent may be added (e.g. Halt). Complete filling the tank to the desired level with water. If agitation is stopped for more than 5 minutes, re-suspend spray solution by full agitation prior to commencing spraying again. Between loads of Odyssey WDG herbicide, check in-line and nozzle screens and rinse and clean if necessary. Upon completion of spraying Odyssey WDG herbicide, thoroughly flush tank, boom, hoses and in-line and nozzle screens with clean water to avoid possible injury to other crops.

**FLAX, Including Low Linolenic Acid Varieties  
(Western Canada Only)**

For use in flax, Lontrel 72 Herbicide should be applied when the flax is 5 to 10 cm high and the weeds are actively growing. Use Lontrel 72 Herbicide at 280 to 415 g/ha to control Canada thistle, common groundsel, scentless chamomile, wild buckwheat, perennial sow-thistle (top growth) and volunteer alfalfa.

The 415 g/ha rate will extend control of Canada thistle into the following year.

**Tank-Mix Combinations in Flax**

REFER TO THE PRODUCT LABELS OF THESE HERBICIDES FOR A LIST OF OTHER WEEDS CONTROLLED, RATES (IF NOT LISTED IN THE TABLE BELOW) AND TIMINGS OF APPLICATION, WATER VOLUMES AND USE PRECAUTIONS.

Herbicide Tank-Mix Partner	Rate Lontrel 72 Herbicide	Rate Tank-Mix Partner	Additional Weeds Controlled
MCPA Ester or MCPA Amine  MCPA Amine (500 g ae/L) MCPA Ester (500 g ae/L)  MCPA Ester (600 g ae/L)	210 g/ha	420 – 560 g ae/ha  0.84 – 1.12 L/ha  0.7 – 0.93 L/ha	Canada thistle (top growth control), shepherd's-purse, common groundsel, common ragweed, cocklebur, dandelion, stinkweed, lamb's-quarters, tartary buckwheat, scentless chamomile, wild buckwheat, wild mustard, volunteer canola, redroot pigweed*, perennial sow-thistle (top growth), volunteer alfalfa
Poast Ultra	210 – 415 g/ha	0.32 – 0.47 L/ha + Merge Adjuvant 0.75-1.0 L/ha	annual grass weeds, Canada thistle
Poast Ultra + MCPA Ester	210 – 415 g/ha	0.32 – 0.47 L/ha + Merge Adjuvant 0.75-1.0 L/ha + 420 – 560 g ai/ha	broadleaved, annual grasses, and certain perennial broadleaved weeds
Select	280 – 415 g/ha	0.19 L/ha + Amigo Adjuvant at 0.5% v/v	Canada thistle**, wild oats, green foxtail, volunteer barley, volunteer wheat, volunteer oats, and wild buckwheat



Select + MCPA Ester	105– 140 g/ha	0.19 L/ha + Amigo Adjuvant at 0.5% v/v + 420 – 560 g ae/ha	Low rate: 105 g Lontrel 72 Herbicide + 420 g ae/ha MCPA Ester: Canada thistle (low infestation), wild oats, green foxtail, volunteer cereals (wheat, barley, oats).  High rate: 140 g Lontrel 72 Herbicide + 560 g ae/ha MCPA Ester: Canada thistle (medium to high infestation), wild oats, green foxtail, redroot pigweed, smartweed, sow-thistle (annual and perennial (top growth)), volunteer cereals (wheat, barley, oats), volunteer canola, and wild buckwheat
---------------------------	---------------	---	---

\*Refer to MCPA herbicide label for rates and control rating.

\*\*Canada thistle – 280 g Lontrel 72 Herbicide/ha for season-long control, 415 g Lontrel 72 Herbicide/ha control extended into following year.

#### Tank-Mix Instructions

**Note 1:** Rates of MCPA ester herbicide of 420 g active ingredient/ha or higher, or MCPA amine herbicide of 490 g active ingredient/ha or higher may cause some delay in maturity with resulting yield reduction.

**Note 2:** Where contact herbicides such as bromoxynil herbicide are used (which damage the leaves of the Canada thistle) Lontrel 72 Herbicide should be applied 7 to 14 days prior or after an interval of 14 days. This allows the Canada thistle to recover and resume growth.

**Note 3:** Add the correct amount of Lontrel 72 Herbicide to spray tank half filled with water and agitate. Add the correct amount of Poast Ultra herbicide and continue to agitate. Add the correct amount of Merge adjuvant along with the remaining amount of water necessary to fill the spray tank. Continue agitation.

**Note 4:** Add the correct amount of Lontrel 72 Herbicide, then MCPA herbicide to half filled sprayer and agitate for 2 to 3 minutes. Next, add Poast Ultra herbicide, and follow with the addition of Merge adjuvant with the remaining water to the required spray volume. Continuously agitate at all times.

#### OATS (Western Canada Only), WHEAT (SPRING) AND BARLEY (SPRING)

Lontrel 72 Herbicide may be used on wheat (spring), barley (spring) and oats to control Canada thistle, common groundsel, perennial sow-thistle (top growth control), wild buckwheat, scentless chamomile and volunteer alfalfa. Lontrel 72 Herbicide should be applied when the wheat, barley or oats are between the 3 leaf to flag leaf emergence stages of growth and weeds are actively growing. Since Lontrel 72 Herbicide damages legumes such as clover and alfalfa, these should not be undersown into the cereals. See Grazing and Harvesting for Feed Section of label for grazing/harvesting intervals for immature crops.

#### Rates of Use

Lontrel 72 Herbicide may be used alone in cereals for Canada thistle control.

Use 210 g/ha of Lontrel 72 Herbicide for the control of top growth of Canada thistle. This rate will suppress top growth of Canada thistle for 6 to 8 weeks. Some regrowth may occur by the end of the season but will not interfere with the harvesting of the crop.

Use 280 g/ha of Lontrel 72 Herbicide for season long control of Canada thistle.

**Tank-Mix Combinations Oats (Western Canada Only), Spring Wheat and Barley**

REFER TO THE PRODUCT LABELS OF THESE HERBICIDES FOR A LIST OF OTHER WEEDS CONTROLLED, RATES (IF NOT LISTED IN THE TABLE BELOW) AND TIMINGS OF APPLICATION, WATER VOLUMES AND USE PRECAUTIONS.

**The following tank mixtures will control both annual and perennial broadleaved weeds listed on the tank-mix partner labels and in addition the weeds named in the Comments column below**

Herbicide Tank-Mix Partner	Crops Registered	Lontrel 72 Herbicide	Tank-Mix Partner	Additional Weeds Controlled
2,4-D Ester or 2,4-D Amine  2,4-D Amine (470 g ae/L)  2,4-D Amine (564 g ae/L) 2,4-D Ester (564 g ae/L)  2,4-D Ester (660 g ae/L)	spring wheat durum wheat spring barley	140 - 210 g/ha	420 - 560 g ae/ha  0.9 - 1.2 L/ha  0.75 – 1.0 L/ha “  0.64 – 0.85 L/ha	Canada thistle (Lontrel 72 Herbicide at 140 g/ha will control Canada thistle for 6 to 8 weeks and at 210 g/ha rate will provide season long control)  DO NOT USE ON OATS
MCPA Ester or MCPA Amine  MCPA Amine (500 g ae/L) MCPA Ester (500 g ae/L)  MCPA Ester (600 g ae/L)	spring wheat durum wheat spring barley oats	140 - 210 g/ha	420 – 560 g ae/ha  0.84 – 1.12 L/ha “  0.7 – 0.93 L/ha	Canada thistle (Lontrel 72 Herbicide at 140 g/ha will control Canada thistle for 6 to 8 weeks and at 210 g/ha rate will provide season long control)
MCPA Ester + Achieve Liquid	spring wheat durum wheat spring barley	105- 140 g/ha	420 - 560 g ae/ha + 0.5 L/ha + Turbocharge 0.5% v/v	green foxtail, yellow foxtail, barnyard grass, Persian dandel, wild oats  DO NOT USE ON OATS
MCPA Ester + Assert 300 SC	spring wheat durum wheat spring barley	105- 140 g/ha	420 - 560 g ae/ha + 1.3 - 1.6 L/ha	wild oats: use 1.3 L/ha of Assert 300 SC on 1-3 leaf wild oats and 1.6 L/ha of Assert 300 SC on 4 leaf wild oats  DO NOT USE ON OATS
MCPA Ester + Axial 100 EC	spring wheat spring barley	105- 140 g/ha	420 - 560 g ae/ha + 600 mL/ha + Adigor 700 mL/ha	wild oats, green foxtail, yellow foxtail, volunteer oats, volunteer canary seed, proso millet  DO NOT USE ON OATS
MCPA Ester + Everest 2.0 Herbicide	spring wheat	105- 140 g/ha	420 - 560 g ae/ha + 72 mL/ha	green foxtail, wild oats  DO NOT USE ON OATS

MCPA Ester + Horizon 240 EC	spring wheat durum wheat	105- 140 g/ha	420 - 560 g ae/ha + 230 - 290 mL/ha + Score Adjuvant 0.8 - 1.0% v/v	wild oats, green & yellow foxtail, volunteer (tame) oats, barnyard grass, volunteer canary seed at 230 mL/ha Horizon 240 EC + 0.8% v/v Score Adjuvant.  above weeds + Persian darnel at 290mL/ha Horizon 240 EC + 1.0% v/v Score Adjuvant  DO NOT USE ON OATS
MCPA Ester + Puma 120 Super	spring wheat durum wheat spring barley	105- 140 g/ha	420 - 560 g ae/ha + 385 - 770 mL/ha	green foxtail at 385 mL/ha of Puma 120 Super; green foxtail, wild oats, barnyard grass at 770 mL/ha of Puma 120 Super  DO NOT USE ON OATS
Florasulam SC + MCPA Ester	spring wheat durum wheat spring barley oats	105 g/ha	0.1 L/ha + 420 g ae/ha	Canada thistle, volunteer canola*, common chickweed, cleavers, dandelions (seedlings; over-wintered rosettes <15 cm), flixweed (spring rosettes only), hempnettle, lamb's quarters, redroot pigweed, shepherd's purse, smartweed, perennial sowthistle (top growth only)**, annual sowthistle, stinkweed, stork's-bill, wild buckwheat, and wild mustard, and suppression of dandelion (over-wintered rosettes >15 cm; mature plants)  *Including Roundup Ready, Liberty-Link and Smart herbicide-tolerant canola varieties **Control not observed until a minimum of 40 days after treatment

Florasulam SC + MCPA Ester + Assert 300 SC	spring wheat durum wheat spring barley	105 g/ha	0.1 L/ha + 420 g ae/ha + 1.6 L/ha	wild oats Use 1.3 L/ha of Assert 300 SC on 1-3 leaf wild oats and 1.6 L/ha of Assert 300 SC on 4 leaf wild oats  DO NOT USE ON OATS
Florasulam SC + MCPA Ester + Everest 2.0 Herbicide	spring wheat durum wheat	105 g/ha	0.1 L/ha + 420 g ae/ha + 72 mL/ha + Agral 90 or Ag-Surf 0.25%v/v	green foxtail, wild oats  DO NOT USE ON OATS
Florasulam SC + MCPA Ester + Axial 100 EC	spring wheat spring barley	105 g/ha	0.1 L/ha + 420 g ae/ha + 600 mL/ha + ADIGOR 700 mL/ha	wild oats, green foxtail* yellow foxtail, volunteer oats, volunteer canary seed, proso millet and barnyard grass**  *Suppression only of green foxtail. **A reduction in barnyard grass control may be observed with this tank-mix combination  DO NOT USE ON OATS

Starane™ II + MCPA Ester	spring wheat durum wheat spring barley	105 g/ha	0.31 L/ha + 420 g ae/ha	<p>Low Rate: cleavers (1-4 whorls), Canada thistle (low infestations), volunteer flax (1-12 cm), flixweed (spring seedling 2-4 lf), kochia<sup>♦♦♦</sup>, lamb's-quarters, wild mustard, shepherd's purse, stinkweed, volunteer sunflower and suppression of stork's bill (1-8 leaf)</p> <p>High Rate: cleavers (1-4 whorls), flixweed<sup>♦♦</sup>, kochia<sup>♦♦♦</sup>, lamb's quarters, shepherd's purse, stinkweed, stork's bill (1-8 leaf), sunflower (volunteer), volunteer flax (1-12, cm), wild mustard, tartary buckwheat, wild buckwheat (1-4 leaf), Canada thistle<sup>♦</sup> (medium to high infestations), volunteer canola, dandelion<sup>♦♦</sup>, common groundsel, round-leaved mallow (1-6 leaf), red-root pigweed, Russian pigweed, scentless chamomile, smartweed, annual sowthistle, perennial sowthistle<sup>♦</sup>, and suppression of common chickweed<sup>♦♦♦</sup>, hemp-nettle (2-6 leaf stage)</p> <p>♦ Season long control, with some regrowth in the fall (top growth control). ♦♦ spring rosettes only. ♦♦♦ Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.</p> <p>DO NOT USE ON OATS</p>
		or	or	
		140 g/ha	0.41 L/ha + 560 g ae/ha	

<p>Starane II + MCPA Ester + Achieve Liquid</p>	<p>spring wheat durum wheat spring barley</p>	<p>105 g/ha  or  140 g/ha</p>	<p>0.31 L/ha + 420 g ae/ha + 0.5 L/ha + Turbocharge 0.5%v/v  or  0.41 L/ha + 560 g ae/ha + 0.5 L/ha + Turbocharge 0.5%v/v</p>	<p>green foxtail, wild oats  DO NOT USE ON OATS</p>
<p>Starane II + MCPA Ester + Assert 300 SC</p>	<p>spring wheat durum wheat spring barley</p>	<p>105 g/ha  or  140 g/ha</p>	<p>0.31 L/ha + 420 g ae/ha + 1.3 - 1.6 L/ha  or  0.41 L/ha + 560 g ae/ha + 1.3 - 1.6 L/ha</p>	<p>wild oats Use 1.3 L/ha of Assert 300 SC on 3-4 leaf wild oats and 1.6 L/ha of Assert 300 SC on 4 leaf wild oats  DO NOT USE ON OATS</p>
<p>Starane II + MCPA Ester + Everest 2.0 Herbicide</p>	<p>spring wheat durum wheat</p>	<p>105 g/ha  or  140 g/ha</p>	<p>0.31 L/ha + 420 g ae/ha + 72 mL/ha  or  0.41 L/ha + 560 g ae/ha + 72 mL/ha</p>	<p>green foxtail, wild oats  DO NOT USE ON OATS</p>

Starane II + MCPA Ester + Horizon 240 EC	spring wheat durum wheat	105 g/ha  or 140 g/ha	0.31 L/ha + 420 g ae/ha + 230 mL/ha + Score Adjuvant 0.8% v/v  or 0.41 L/ha + 560 g ae/ha + 230 mL/ha + Score Adjuvant 0.8% v/v	green foxtail, wild oats  DO NOT USE ON OATS
Starane II + MCPA Ester + Puma 120 Super	spring wheat durum wheat spring barley	105 g/ha  or 140 g/ha	0.31 L/ha + 420 g ae/ha + 385 - 770 mL/ha  or 0.41 L/ha + 560 g ae/ha + 385 - 770 mL/ha	green foxtail at 385 mL/ha of Puma 120 Super,  green foxtail, wild oats, and barnyard grass at 770 mL/ha of Puma 120 Super  DO NOT USE ON OATS
Refine Extra + 2,4-D Ester or MCPA Ester	spring wheat spring barley (Western Canada Only)	105 g/ha	20 g/ha + Agral 90 Adjuvant 0.2% v/v + 420 g ae/ha	perennial sow thistle, wild buckwheat, volunteer canola, wild mustard, lady's thumb and stinkweed, while providing seasonal control of Canada thistle and suppression of cleavers  DO NOT USE ON OATS

### Tank-Mix Instructions

**Note 1:** If a tank-mix partner requires the addition of an adjuvant add the recommended adjuvant and dilution rate to the tank-mix.

**Note 2:** When tank mixing with Refine Extra, mix Refine Extra herbicide first in fresh water, then add the required amount of Lontrel 72 Herbicide followed by MCPA or 2,4-D herbicides. Add the surfactant last.

### SUMMERFALLOW AND NON-CROP FARMLAND

Lontrel 72 Herbicide may be used on summerfallow (one application per year) and non-crop farmland (around farm buildings, storage areas, fence rows, etc.) at 415 g/ha for the control of Canada thistle, scentless chamomile, common groundsel, wild buckwheat, the top growth of perennial sow-thistle and volunteer alfalfa. The Canada thistle plants should be between the rosette and the early bud stage and actively growing at the time of spraying.

## **SEEDLING AND ESTABLISHED GRASSES FOR SEED PRODUCTION AND FORAGE (WESTERN CANADA ONLY)**

Including Kentucky bluegrass, smooth brome grass, reed canary grass, creeping red fescue, meadow fescue, tall fescue, meadow foxtail, orchard grass, altai wild ryegrass, Russian wild ryegrass, timothy, crested wheatgrass, intermediate wheatgrass, slender wheatgrass and streambank wheatgrass for forage and seed production and tall wheatgrass for forage only: For control of the weeds listed on the label plus alsike clover, apply Lontrel 72 Herbicide at the rate of 210 to 415 g/ha in 110 to 220 L/ha of water. Make one application per season by ground sprayer. For seedling grasses, apply at the 2 to 4 leaf stage. For established grasses, apply at the shot-blade stage, or in the fall after harvest or in early spring. See Grazing and Harvesting for Feed Section.

## **BALSAM FIR CHRISTMAS TREE PLANTATIONS**

For the control of vetch (*Vicia* spp.) apply Lontrel 72 Herbicide at 210 g/ha in 150 to 200 L/ha of water as a directed foliar application using a hydraulic sprayer. Best control is obtained when vetch stems are 10 to 15 cm long and prior to the vetch climbing into a tree crown. Avoid contact with the upper two thirds of the tree crown. Do not use on seedbeds, transplants or any over-the-top applications.

## **SUGAR BEETS**

For Canada thistle control apply Lontrel 72 Herbicide at 280 to 415 g/ha with ground equipment as a foliar spray either broadcast or in a band over the row. When applied in the band, the amount of Lontrel 72 Herbicide should be reduced proportional to the band width. Lontrel 72 Herbicide should be applied when sugarbeets are in the cotyledon to 8 leaf stage. For the most effective control of Canada thistle, apply Lontrel 72 Herbicide as a broadcast treatment to the entire infested area. Do not apply within 90 days of harvest.

## **RUTABAGA**

For control of common ragweed, apply Lontrel 72 Herbicide with a boom sprayer at the rate of 280 g/ha in approximately 200 to 300 L/ha of water. Apply as a postemergent spray when ragweed plants are 5 to 10 cm tall. Application to larger ragweed plants will result in reduced weed control. Make only one application per season. Preharvest interval is 83 days

**NOTE TO USER:** READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS: The DIRECTIONS FOR USE for this product for the use(s) described below were developed by persons other than Dow AgroSciences Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Dow AgroSciences Canada Inc. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crop(s) listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Dow AgroSciences Canada Inc. harmless from any claims based on efficacy and/or phytotoxicity in connection with the use(s) described on this label.

## **DIRECTIONS FOR USE**

**HIGHBUSH BLUEBERRY:** Make one application per year, post emergent for the control of vetch and red and white clover. Apply as a broadcast treatment. Apply at a rate of 210 g/ha for vetch and 415 g/ha for red and white clover. The Pre-harvest interval is 45 days.

**CANOLA - ONTARIO ONLY:** To be used on Canola in Ontario on the following NATIONALLY REGISTERED spring canola cultivars: Cyclone, Ebony, Jewel, 46A65 and Hyola 401.



Weeds Controlled	Rate	4.45 L Bottle Will Treat	For Optimum Results
Canada thistle, scentless chamomile, wild buckwheat, perennial sow-thistle (top growth), common groundsel	210 g/ha to 280 g/ha	10.6 ha to 7.9ha	1. Treat when weeds are actively growing 2. Use 100 to 200 L/ha spray solution for uniform coverage

Make one application per season; post emergent. Apply at the 2-6 leaf stage of canola, when weeds are actively growing. Apply to Canada thistle at the rosette to pre-bud stage.

**BEARING AND NON-BEARING APPLES:** To be used on bearing and non-bearing apples for control of perennial vetch species. Apply at a rate of 280 g/ha on vetch at the early flowering stage as a spot treatment. Apply as a ground application only. Avoid contact of the spray with the tree limbs. For best results apply in early spring. Allow at least 30 days between weed treatment application and harvest.

**CABBAGE, CAULIFLOWER, BROCCOLI AND KOHLRABI (ALL TRANSPLANTED), NAPPA CABBAGE (TRANSPLANTED AND SEEDED), CHINESE RADISH, MUSTARD CABBAGE AND CHINESE BROCCOLI (ALL SEEDED).** Make one application per year to control ragweed, vetch, common groundsel, Canada thistle and for suppression of sheep sorrel. Apply post planting as a ground application only.

**Application rate:** Apply at a rate of 280 g/ha in 300 L water/ha. The pre-harvest interval (PHI) is 30 days.

**CRANBERRY: Make one-two applications per year,** for the control of vetch. Apply with wiper-type application equipment.

Wipe treatments may be applied as a spot application. The treatment may be applied using equipment such as a hockey stick applicator. **The treatment solution should be wiped onto weed foliage which extends above the cranberry canopy.** Wiper applications may be made in the fall at least 2 weeks after harvest and after the vines have attained their winter dormancy colour, and in the spring prior to bud-break. Wiper application treatments may also be applied following cranberry bud-break (first emergence (1 to 2 mm) of terminal meristem) to control late emerging weeds or weeds which escaped earlier control measures. **Contact of the treatment solution with cranberry foliage after bud break should be avoided since it will result in plant injury.**

**Application rate:** Apply a solution of 10 g Lontrel 72 Herbicide in 1 L water. The preharvest interval is 60 days.

**WEED CONTROL IN SHELTERBELTS:** For control of Canada thistle in shelterbelts of villosa lilac, acute willow, Colorado spruce, white spruce, buffaloberry and chokecherry. Make one application per year. Apply to Canada thistle at the rosette to pre-bud stage, post emergent, ground application only.

**Application rate:** Apply at a rate of 415 g/ha.

### **STRAWBERRY (Renovation)**

For control of tufted vetch and Canada thistle and suppression of sheep sorrel and ox-eye daisy, apply Lontrel 72 Herbicide at the rate of 280 to 415 g/ha. Apply as a broadcast application with a boom sprayer calibrated to deliver a total volume of 150 to 200 L/ha. For Canada thistle control, refer to the Control of Canada Thistle section for rate selection. For control of tufted vetch apply at the 280 g/ha rate. For sheep sorrel and ox-eye daisy, apply at the 415 g/ha rate. Apply as the single treatment immediately after harvest but before mowing. Wait at least 7 to 10 days after Lontrel 72 Herbicide application before mowing. Do not apply Lontrel 72 Herbicide after mid-August because of its possible effects on runner development and flower bud formation. Later applications of Lontrel 72 Herbicide may cause crop damage resulting in reduced yields in the season following treatment.

Apply Lontrel 72 Herbicide only as a summer renovation treatment.

Do not apply Lontrel 72 Herbicide more than once per year.

Early strawberry varieties such as Annapolis or Veestar may be more susceptible to injury. Certain environmental stresses such as drought, flooding or severe overwintering conditions may increase the risk of injury from Lontrel 72 Herbicide.

### **CONTROL OF TUFTED VETCH IN LOWBUSH BLUEBERRY IN EASTERN CANADA ONLY**

Apply Lontrel 72 Herbicide to control tufted vetch in lowbush blueberry. **FOR SPOT APPLICATION ONLY.** When using a hand gun or backpack sprayer to treat small infestations, apply Lontrel 72 Herbicide at a rate of 21 g per 1000 square metre area in 200 L of water. When applying with a boom sprayer to treat larger infestations, apply 210 g of Lontrel 72 Herbicide per hectare in 150-200 L water. Make one application per year, in the non-bearing year (prune year). Apply in June OR when tufted vetch is in early bloom. Applications of Lontrel 72 Herbicide may cause crop damage resulting in reduced yields in the season following application. The Pre-harvest interval is 10 months.

### **CONTROL OF CANADA THISTLE AND OTHER LABELLED WEEDS IN POPLAR SPECIES AND THEIR HYBRIDS**

Apply Lontrel 72 Herbicide at a rate of 415 g/ha to control Canada thistle and other labelled weeds in new and established short rotation intensive culture crops of poplar (*Populus*) species and their hybrids. Make one application per year. Apply to Canada thistle in the rosette to pre-bud stage. Apply by ground application only using an overall spray or as a directed spray to the base of the tree. Some leaf cupping and stem twisting may occur, but will not adversely affect growth.

**WARNING:** Poplar clones/hybrids vary in their tolerance to Lontrel 72 Herbicide. Injury observed includes leaf injury, leaf cupping, stem twisting, height reduction and diameter reduction. As not all clones/hybrids have been tested for tolerance to Lontrel 72 Herbicide, use of this product should be limited to a small area of each clone/hybrid to confirm tolerance prior to adoption as a general field practice.

REFER TO THE MAIN LONTREL 72 HERBICIDE LABEL FOR ADDITIONAL DETAILS AND INSTRUCTIONS, INCLUDING ROTATIONAL CROPPING RESTRICTIONS, BEFORE USING THIS PRODUCT

## **TRANSLINE™ IVM SYSTEM (NON-CROP USES)**

### **FOR USE IN BRITISH COLUMBIA INTERIOR, PRAIRIE (including the Peace River Region of B.C.), CENTRAL AND ATLANTIC REGIONS OF CANADA ONLY.**

Lontrel 72 Herbicide may be used on the following non-crop areas: rights-of-way (hydro, railroad, communication lines, pipelines) and associated stations, industrial manufacturing sites, storage sites, vacant lots and roadsides, military bases and low maintenance rough turf areas (grass areas where the dominant species are those listed in the Rangeland and Grass Pasture Section of this label, and where little or no maintenance is applied). This product is not registered for use on fine turf lawns or turf grass receiving high maintenance. Apply between 210 to 415 g/ha depending on weeds present and level of Canada thistle control required. Refer to the Weeds Controlled table for appropriate application rate.

### **TANK-MIX COMBINATIONS**

Lontrel 72 Herbicide may be tank mixed with 2,4-D Amine or Ester or MCPA Amine or Ester for control of additional broadleaf weeds on roadsides and vacant lots. Lontrel 72 Herbicide may also be tank mixed with 2,4-D Amine for additional broadleaf use control on rights-of-way (hydro, railroad, communication lines, pipelines) and associated stations, industrial manufacturing sites and storage sites. Read and follow the label of each tank-mix product used for precautionary statements, directions for use, weeds controlled and any other restrictions. When tank mixing adhere to the most restrictive label limitations and precautions.

Lontrel 72 Herbicide at 210 to 415 g/ha may be tank mixed with 2,4-D or MCPA Herbicides at the rate of 420 to 560 g active ingredient/ha. The tank-mix will control many weeds, including: Canada thistle, cocklebur, common ragweed, dandelion, lamb's-quarters, scentless chamomile, perennial sow-thistle, shepherd's-purse, stinkweed, tartary buckwheat, wild buckwheat and wild mustard. Apply up to the 15 cm height of annual broadleaf weeds.

### **RANGELAND AND GRASS PASTURE**

Including Kentucky bluegrass, smooth brome grass, reed canary grass, creeping red fescue, meadow fescue, tall fescue, meadow foxtail, orchard grass, alтай wild ryegrass, Russian wild ryegrass, timothy, crested wheatgrass, intermediate wheatgrass, slender wheatgrass, streambank wheatgrass and tall wheatgrass.

For control of the weeds on the label plus alsike clover, apply Lontrel 72 Herbicide at the rate of 210 to 415 g/ha in 110 to 120 L/ha of water. Make one application per season by ground sprayer. For seedling grasses, apply at the 2 to 4 leaf stage. For established grasses, apply at the shot-blade stage or in the fall after harvest or early spring. Do not apply tank mixtures containing 2,4-D or MCPA.

**NOTE TO USER:** READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS: The DIRECTIONS FOR USE for this product for the use(s) described below were developed by persons other than Dow AgroSciences Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Dow AgroSciences Canada Inc. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crop(s) listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Dow AgroSciences Canada Inc. harmless from any claims based on efficacy and/or phytotoxicity in connection with the use(s) described on this label.

### **DIRECTIONS FOR USE**

**FOR USE IN BRITISH COLUMBIA INTERIOR, PRAIRIE (including the Peace River Region of B.C.), CENTRAL AND ATLANTIC REGIONS OF CANADA ONLY.**

**CONTROL OF SPOTTED AND DIFFUSE KNAPWEED IN NON-CROPS AREAS (rights-of-way [hydro, railroad, communication lines, pipelines] and associated stations, industrial manufacturing sites, storage sites, roadsides, airports, military bases and low maintenance rough turf areas) AND IN RANGELAND, PASTURE AND BALSAM FIR CHRISTMAS TREE STANDS OR PLANTATIONS.**

**Weeds Controlled**

Spotted and Diffuse Knapweed

**Application Rate**

350 g/ha

Make one application per year for the control of spotted and diffuse knapweed. Apply in the spring prior to the bud stage of the weeds. Apply in 100 - 200 L water/ha. Apply to both seedling or established plants.

REFER TO THE MAIN LONTREL 72 HERBICIDE LABEL FOR ADDITIONAL DETAILS AND INSTRUCTIONS, INCLUDING ROTATIONAL CROPPING RESTRICTIONS, BEFORE USING THIS PRODUCT

**BUFFER ZONES**

Uses of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment.

For application to rights-of-way, buffer zones for protection of sensitive terrestrial habitats are not required; however, the best available application strategies which minimize off-site drift, including meteorological conditions (e.g., wind direction, low wind speed) and spray equipment (e.g., coarse droplet sizes, minimizing height above canopy), should be used.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, woodlots, hedgerows, riparian areas and shrublands).

Method of application	Crop	Buffer Zones (metres) Required for the Protection of Terrestrial Habitat:
Field sprayer	Wheat, barley, oats, flax, canola, forage grasses, high-bush blueberry, low-bush blueberry, strawberry, sugar beet, rutabaga, cabbage, broccoli, cauliflower, field corn (hybrid), canary seed, balsam fir, Christmas tree plantations, shelterbelts, poplar and their hybrids, non-crop uses, rangeland and grass pasture	2*
	Apple	3

\*Buffer zones for the protection of terrestrial habitats are not required for use on rights-of-way including railroad ballast, rail and hydro rights-of-way, utility easements, roads, and training grounds and firing ranges on military bases.

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank-mix partners.

**NOTE:** Applicators may recalculate a site-specific buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: 'DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) [Fine or Medium or Coarse] classification.' To access the Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, Lontrel 72 Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Lontrel 72 Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Lontrel 72 Herbicide or other Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Dow AgroSciences Canada Inc. at 1-800-667-3852 or at [www.dowagro.ca](http://www.dowagro.ca).

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

<sup>TM</sup>Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow  
All other products are registered trademarks of their respective companies

080713

Label Code: CN-31039-001-E

Specimen label notes:  
Initial registration