

Specimen Label

CLOTHIANIDIN	GROUP	4A	INSECTICIDE
--------------	-------	----	-------------



™Trademarks of Corteva Agriscience and its affiliated companies

AN INSECTICIDE SEED TREATMENT PRODUCT PROVIDING PROTECTION AGAINST LISTED INSECT PESTS

Active Ingredient	By Wt.
Clothianidin*.....	47.8%
Other Ingredients.....	52.2%
Total.....	100.0%

*(E)-1-[(2-chlorothiazol-5-yl)methyl]-3-methyl-2-nitroguanidine

Lumisure contains 5 lb clothianidin per gallon.

FIRST AID	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN	
Treatment is supportive and symptomatic.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

EPA Reg. No. 59639-151-352

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators, and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks and chemical resistant gloves made out of: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils or Viton ≥14 mils.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to birds, mammals and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not apply where runoff is likely to occur. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on label.

Treated seed exposed on soil surface maybe hazardous to birds and mammals. Cover or collect spilled seed.

This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE

Store in a cool, dry secured storage area.

Do not store near food or feed.

Store in original container only.

To close package, replace and tighten cap to form an airtight seal.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

[Statement for containers equal to or less than 5 gallons]

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

[Statement for rigid nonrefillable containers greater than 5 gallons]

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

[Statement for all high-density polyethylene (HDPE) refillable containers]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 12 hours.

Exception: Once the seeds are planted in soil, the WPS, under certain circumstances, allows workers to enter the treated area without restriction if there will be no worker contact with the soil subsurface or treated seed. PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water, is: long-sleeved shirt and long pants; shoes plus socks, and chemical resistant gloves.

RESISTANCE MANAGEMENT

For insecticide resistance management, please note that Lumisure contains a Group 4A insecticide: clothianidin. Any insect population may contain individuals naturally resistant to Lumisure and other Group 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay insecticide/acaricide resistance, take the following steps:

- Rotate the use of Lumisure or other Group 4A insecticides/acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides/acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance contact 1-800-258-3033.

Use Instructions

This product is for commercial seed treatment facilities as specified below. This product may only be used for on-farm application for cereal grains (except corn, rice, sorghum, and wild rice). Use Lumisure only in accordance with specifications on this label utilizing mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. Always mix product thoroughly before use.

As stated by regulations specified in 40 CFR 153.155, all seed treated with Lumisure must be colored with an EPA-approved dye or colorant to visually distinguish seed that is treated with a pesticide. It is mandatory that such dye or colorant be added into the slurry mixture containing Lumisure prior to application. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.

RESTRICTIONS

- Regardless of application method, do not apply more than 0.2 lb active ingredient clothianidin/A/calendar year.
- Do not use treated seed for feed, food, or oil processing.
- Store treated seed away from feeds and other foodstuffs.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- Do not re-use bags from treated seed to handle food or feed products.
- The compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.
- Do not apply this product in hopper box or planter box at planting time.
- Do not use for on-farm seed treatment applications except for cereal grains (except corn, rice, sorghum and wild rice).
- All treated seeds must be planted at a minimum depth of 1 inch unless otherwise specified.
- Not for use on rice, sorghum, or soybean in the state of California.

Seed Quality And Storage Of Treated Seed

Seed treatments applied to low vigor or poor quality seed or to mechanically damaged seed may result in loss of seed germination or reduced seedling vigor. To verify seed safety, treat a small portion of seed with the desired slurry mixture (Lumisure alone or in combination with other commercial seed treatment products) prior to treatment of an entire seed lot. Utilize standard seed germination methodologies to determine and verify seed safety. Proper storage conditions of treated seed are to be followed. Due to seed quality and seed storage conditions beyond the control of Corteva Agriscience, no claims are made to guarantee the germination of carry-over treated seed.

Tank Mixing

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Accurate application of Lumisure to seed is required for control of pests. Pre-test tank mixtures with other seed treatment products to verify formulation compatibility and to assure proper physical compatibility of products.

PRODUCT INFORMATION

Lumisure (clothianidin) is a systemic insecticide seed treatment providing early season protection against listed insect pests of barley, buckwheat, field corn, pearl millet, popcorn, proso millet, oats, rice**, rye, sorghum**, soybean**, sugar beet, sweet corn, teosinte, triticale, and wheat. This product provides protection under low to moderate corn rootworm pressure.

**Not for use in California.

ROTATIONAL CROP PLANT-BACK INTERVALS

Crops Planted	Immediate Plant-back	30-Day Plant-back	4-Month Plant-back	8-Month Plant-back	12-Month Plant-back
Barley Buckwheat Field Corn Pearl Millet Popcorn Proso Millet Oats Rice** Rye Soybean** Sugar Beet Teosinte Triticale Wheat	Bulb Vegetables Canola Cereal Grains Cotton Fruiting Vegetables (including Cucurbits) Leafy Vegetables (including Brassica Vegetables) Rapeseed Root and Tuber Vegetables Soybean	Dry Beans Grasses Non-grass Animal Feeds		Sugarcane	Any crops without an earlier plant- back interval
Sorghum** Sweet Corn	Bulb Vegetables Canola Cereal Grains Cotton Fruiting Vegetables (including Cucurbits) Leafy Vegetables (including Brassica Vegetables) Rapeseed Root and Tuber Vegetables Soybean	Dry Beans Grasses Non-grass Animal Feeds	Any crop without an earlier plant- back interval		

**Not for use in California.

ON-FARM APPLICATION

CEREAL GRAINS (Except Corn, Rice, Sorghum, and Wild Rice)			
Crops	Insect Pests	Lumisure Use Rates fl oz/100 lb seed	Rates/Instructions
Barley Buckwheat Pearl Millet Proso Millet Oats Rye Teosinte Triticale	Wireworm	0.25 – 1.79 (0.0098-0.07 lb ai/100 lb seed)	For wireworm seed protection, use a 0.25 fl oz/100 lb seed rate or higher, depending on field history and site populations. Use the higher rate to provide protection from heavy insect pressure.
	Aphid	0.75 – 1.79 (0.029-0.07 lb ai/100 lb seed)	Use the higher rate to provide increased length of protection and for heavy insect pressure.
Wheat	Wireworm	0.25 – 1.79 (0.0098-0.07 lb ai/100 lb seed)	For wireworm seed protection, use a 0.25 fl oz/100 lb seed rate or higher, depending on field history and site populations. Use the higher rate to provide protection from heavy insect pressure.
	Aphid	0.75 – 1.79 (0.029-0.07 lb ai/100 lb seed)	Use the higher rate to provide increased length of protection and for heavy insect pressure.
	Hessian Fly	1.79 (0.07 lb ai/100 lb seed)	For Hessian Fly protection to wheat seedlings in the Fall season, use 1.79 fl oz/100 lb seed.
Restriction Regardless of application method, do not apply more than 0.2 lb active ingredient clothianidin/A/calendar year.			

COMMERCIAL SEED TREATMENT FACILITIES ONLY

CEREAL GRAINS (Except Corn, Rice, Sorghum, and Wild Rice)			
Crops	Insect Pests	Lumisure Use Rates fl oz/100 lb seed	Rates/Instructions
Barley Buckwheat Pearl Millet Proso Millet Oats Rye Teosinte Triticale	Wireworm	0.25 – 1.79 (0.0098-0.07 lb ai/100 lb seed)	For wireworm seed protection, use a 0.25 fl oz/100 lb seed rate or higher, depending on field history and site populations. Use the higher rate to provide protection from heavy insect pressure.
	Aphid	0.75 – 1.79 (0.029-0.07 lb ai/100 lb seed)	Use the higher rate to provide increased length of protection and for heavy insect pressure.

COMMERCIAL SEED TREATMENT FACILITIES ONLY (Cont.)

CEREAL GRAINS (Except Corn, Rice, Sorghum, and Wild Rice)			
Crops	Insect Pests	Lumisure Use Rates fl oz/100 lb seed	Rates/Instructions
Wheat	Wireworm	0.25 – 1.79 (0.0098-0.07 lb ai/100 lb seed)	For wireworm seed protection, use a 0.25 fl oz/100 lb seed rate or higher, depending on field history and site populations. Use the higher rate to provide protection from heavy insect pressure.
	Aphid	0.75 – 1.79 (0.029-0.07 lb ai/100 lb seed)	Use the higher rate to provide increased length of protection and for heavy insect pressure.
	Hessian Fly	1.79 (0.07 lb ai/100 lb seed)	
Restriction Regardless of application method, do not apply more than 0.2 lb active ingredient clothianidin/A/calendar year.			

CORN				
Crops	Insect Pests	Lumisure Use Rates		Rates/Instructions
		fl oz per 80,000 seeds	mg ai/kernel	
Corn: Seed Field Pop Sweet	Chinch Bug Corn Flea Beetle Corn Leaf Aphid Cutworm (black) Grape Colaspis Seedcorn Maggot Southern Corn Leaf Beetle Southern Corn Rootworm Southern Green Stinkbug Storage Pests, including: Indian Meal Moth Lesser Grain Borer Red Flour Beetle Rice Weevil Rusty Grain Beetle Sugarcane Beetle White Grub, including: European Chafer Larvae May/June Beetle Larvae Japanese Beetle Larvae Thrips Wireworm	1.13 to 2.26	0.25 to 0.5	Provide early seed protection against listed insect pests. For additional insect protection commercially treated seed with 0.25 mg ai/kernel rate of clothianidin may be over-treated with up to 0.25 mg/ai/kernel to equal total dose rate of 0.5 mg ai/kernel. For Storage Pests: Follow guidelines for control of stored grain insects, it is advised that seed with existing populations of storage pests be fumigated prior to treating and bagging of seed.
Field Corn Popcorn	Corn Rootworm, including: Mexican Northern Southern Western Southern Corn Billbug	5.64	1.25	
Restriction Regardless of application method, do not apply more than 0.2 lb active ingredient clothianidin/A/calendar year.				

RICE (Not for Use in California)		
Insect Pests	Lumisure Use Rates fl oz/100 lb seed	Rates/Instructions
Aphid Black Bug Chinch Bug Grape Colaspis Rice Water Weevil Thrips	1.92 (0.075 lb ai/100 lb seed)	Lumisure provides early season protection against listed insect pests.
Restrictions <ul style="list-style-type: none"> • Lumisure is to be used only as a dry-seeded rice seed treatment. Cover the planted seed thoroughly by soil. • Lumisure is not to be used in areas where rice/crawfish aquaculture practices are in place or in or near fish farms, shrimp, prawn or crab pond or nursery operations. • Lumisure is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert. • Do not apply to Wild Rice. • Regardless of application method, do not apply more than a total of 0.2 lb active ingredient clothianidin/A/calendar year. 		

SORGHUM (Not for Use in California)		
Insect Pests	Lumisure Use Rates fl oz/100 lb seed	Rates/Instructions
Black Granulate Cutworm Chinch Bug Corn Leaf Aphid English Grain Aphid Greenbug Seed Corn Maggot White Grub Wireworm Yellow Sugarcane Aphid	5.1 to 6.4 (0.2 to 0.25 lb ai/100 lb seed)	Lumisure provides protection against listed insect pests. Use the higher rate to provide increased length of protection and for heavy insect pressure.
Restrictions Regardless of application method, do not apply more than 0.2 lb active ingredient clothianidin/A/calendar year.		

SOYBEAN (Not for Use in California)			
Insect Pests	Lumisure Use Rates fl oz/140,000 seeds	Lumisure Use Rates mg ai/seed	Rates/Instructions
Aphids Bean Leaf Beetle Grape Colaspis Leafhoppers Pea Leaf Weevil Seed Corn Maggot Three Cornered Alfalfa Hopper Thrips White Grubs Wireworm	0.87	0.11	Lumisure provides protection against listed insect pests.
Restrictions <ul style="list-style-type: none"> • Regardless of application method, do not apply more than 0.2 lb active ingredient clothianidin/A/calendar year. • Do not graze or feed soybean forage and hay to livestock. 			

SUGAR BEET		
Insect Pests	Lumisure Use Rates fl oz per 100,000 pelleted seeds	Rates/Instructions
Black Bean Aphid Cutworms Flea Beetle Garden Springtail Sugar Beet Leafhopper Sugar Beet Leafminer Sugar Beet Root Maggot Wireworm	3.4 (0.6 mg ai/seed)	Lumisure provides protection against listed insect pests. Protection against beet leafhopper injury aids in reducing potential spread of beet curly top virus, vectored by beet leafhopper. Apply Lumisure to seed with a coating system giving a minimum weight gain build-up of approximately 30%.
Restrictions Regardless of application method, do not apply more than 0.2 lb active ingredient clothianidin/A/calendar year.		

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with Lumisure must be labeled with the following information:

- This seed has been treated with Lumisure which contains clothianidin.
- Do not use treated seed for feed, food, or oil processing.

In addition, the U.S. Environmental Protection Agency requires the following statements on the container of seed treated with Lumisure:

- Regardless of application method, do not apply more 0.2 lb active ingredient clothianidin/A/calendar year.
- Lumisure is to be used only as a dry-seeded rice seed treatment. Cover the planted seed thoroughly by soil.
- Do not plant Lumisure-treated rice in areas where rice/crawfish aquaculture practices are in place or in or near fish farms, shrimp, prawn or crab pond or nursery operations.
- Store treated seed away from feeds and other foodstuffs.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Wear long sleeved shirt, long pants, and chemical resistant gloves when handling treated seed.
- Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

- Dispose of seed packaging or containers in accordance with local requirements. Do not use empty seed bags for any other purpose.
- The compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.
- All seeds must be planted at a minimum depth of 1 inch.
- Do not graze or feed soybean forage and hay to livestock.
- Not for use on rice, sorghum and soybean in California.
- **Rotational Crop Plant-Back Intervals:** Areas planted with treated seed of barley, buckwheat, field corn, pearl millet, popcorn, proso millet, oats, rice**, rye, sugar beet, soybean**, teosinte, triticale, and wheat may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables, and soybean. These areas may also be replanted after 30 days with dry beans, grasses, and non-grass animal feeds. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- **Rotational Crop Plant-Back Intervals:** Areas planted with treated seed of sorghum** and sweet corn may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables, and soybean. These areas may also be replanted after 30 days with dry beans, grasses, and non-grass animal feeds. Do not plant any other crops in the treated area for at least 4 months after these treated seeds are planted.

** Not for use on rice, sorghum, and soybean in California.

- For barley, buckwheat, pearl millet, proso millet, oats, rye, teosinte, triticale wheat, this seed has been treated with a rate of XX [0.0098 -0.07] fl oz/100 lb seed which will result in XX lb ai clothianidin per 100 lb seed.
- For field corn, popcorn, seed corn, sweetcorn, this seed has been treated with XX [0.25 - 1.25] mg clothianidin per seed.
- For rice, this seed has been treated with 0.075 lb ai clothianidin per 100 lb seed.
- For sorghum, this seed has been treated with a rate of XX [5.1-6.4] fl oz/ 100 lb seed which will result in XX lb ai clothianidin per 100 lb seed.
- For soybean, this been treated with 0.11 mg clothianidin per seed.
- For sugar beet, this seed has been treated with 0.6 mg clothianidin per seed.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation, of Remedies.

Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. Corteva Agriscience will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by Corteva Agriscience. To the extent permitted by law, all such risks associated with non-directed use shall be assumed by buyer and/or user.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Terms and Conditions of Use (Cont.)

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Corteva Agriscience be liable for consequential, incidental or special damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator, and/or application advisor. **Read and follow the entire label of each product to be used in the tank mix with this product.**

™Trademarks of Corteva Agriscience and its affiliated companies

Manufactured For:
Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268

Label Code: CD02-648-021
 Initial publication

Revisions:

1. Updated First Aid and PPE
2. Updated Storage and Disposal
3. Updated Use Restrictions and Resistance Management
4. Updated Rotational Crop Plant-Back Intervals
5. Updated instructions, restrictions and rates for cereal grains, corn, sorghum, soybean, sugar beet
6. Updated Seed Bag Tag
7. Updated legal entity, phone numbers, and warranty disclaimers