

MAX-IN[®] Copper

The Next Generation of Foliar Nutrition

Guaranteed Analysis

Copper5.0%

Derived from: Copper EDTA

* U.S. Patent No. 7,666,242

Store Above 40° F

KEEP OUT OF REACH OF CHILDREN



WARNING

Harmful if swallowed. Causes skin irritation. Causes eye irritation. May cause respiratory irritation.

PRECAUTIONARY STATEMENTS

Wash hands thoroughly after handling. Wear protective gloves. Do not eat, drink or smoke when using this product. Avoid breathing mist or vapors. Use only outdoors in a well-ventilated area. Store in a well-ventilated and secured place. Keep container tightly closed. MAX-IN[®] COPPER contains EDTA which can be corrosive to metals. Recommended materials of construction for handling and storage are fiberglass, CPVC or PVC and polyethylene. Dispose of contents/container in accordance with Federal, state and local regulations.

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor for treatment advice. Rinse mouth.

IF ON SKIN: Wash with plenty of water. If skin irritation occurs: Get medical attention. Take off contaminated clothing and wash before reuse.

IF IN EYES: Rinse cautiously with water for 15 minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical attention.

IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a poison control center or doctor for treatment advice if you feel unwell.

For Chemical Spill, Leak, Fire or Exposure, Call Chemtrec (800) 424-9300.

GENERAL INFORMATION

For use in a balanced fertility program for all crops requiring copper. User should seek the advice of a certified crop advisor or trained agronomist. For specific crop recommendations in your area, consult your state's recommendations.

CAUTION: Over-application or excessive use of copper, or use on crops other than those listed as having a copper response, can result in copper toxicity and may cause serious crop injury.

MAX-IN[®] COPPER is not to be used as a replacement for any surfactant in recommendations tank-mixed with herbicides.

F1683

Distributed and Guaranteed By:
Winfield Solutions, LLC
PO Box 64589
St. Paul, Minnesota 55164-0589

1/0105/7

Weight Per Gallon: 9.92 pounds (4.50 kg)
NET WEIGHT: 24.80 lbs. (28.75 kg)
NET CONTENTS: 2.5 Gallons (9.464 Liters)

WINFIELD™

MAX-IN® COPPER is specifically designed for use on crops that have displayed deficiencies to copper. Use in a minimum of 10 gallons of water per acre.

DIRECTIONS FOR USE

MAX-IN® COPPER includes SureTank™ technology for compatibility with many agricultural pesticides. However, if local information is not available, a jar test for compatibility should be conducted prior to large scale mixing. When using **MAX-IN® COPPER** with pesticides read and follow the pesticide label instructions. Do not mix **MAX-IN® COPPER** where prohibited by the pesticide label. Mixing **MAX-IN® COPPER** with certain pesticides or adjuvants, especially oil based products can increase the risk of phytotoxicity to sensitive crops in conditions that may increase the activity of the spray solution. Do not mix **MAX-IN® COPPER** with pesticides unless potential phytotoxicity can be tolerated. If uncertain, spray a small test area with the spray mixture before making large scale applications.

Foliar Application:

Alfalfa: Apply 8 to 10 ounces per acre of **MAX-IN® COPPER** after each cutting. Applications should be made when sufficient re-growth has occurred.

Canola: Apply 8 to 10 ounces per acre when Canola is in the earlier vegetative stages. Later application may be needed; tissue samples should be used to determine later usages.

Cereal Crops: Apply 8 to 10 ounces per acre when plants are between tillering and early joint stages. Additional application at boot may be needed; tissue samples should be used to determine later usages.

Corn: Apply 8 to 10 ounces per acre when corn is in the V3 to V8 stage. Later application may be needed; tissue samples should be used to determine later usages.

Cotton: Apply 8 to 10 ounces per acre between the first true leaf and pin head square and/or 8 to 10 ounces per acre from first bloom to boll fill. Later application may be needed; tissue samples should be used to determine later usages.

MAX-IN® COPPER can be tank mixed with MAX-IN Cotton.

Peanuts: Use 8 to 10 ounces per acre of **MAX-IN® COPPER** during the vegetative stage in a minimum of 10 gallons of water by ground. A second application may be made at pegging.

Peppers and Chilies: Apply 8 to 10 ounces per acre when plants are in the earlier reproductive stage. Repeat applications at 10 to 14 days as long as fruiting continues.

Potatoes: Apply 8 to 10 ounces per acre from emergence to tuber initiation. Later application may be needed; tissue samples should be used to determine later usages.

Soybeans: Apply 8 to 10 ounces per acre when soybeans are in the vegetative stage (V3 to V5) or earlier reproductive stage. Later application may be needed; tissue samples should be used to determine later usages.

Sugar Beets: Apply 8 to 10 ounces per acre between the second and 10 true leaf stages. Later application may be needed; tissue samples should be used to determine later usages. **MAX-IN® COPPER** can be tank mixed with glyphosate herbicides.

Sunflowers: Apply 8 to 10 ounces per acre when Sunflowers are in the earlier flower initiation period through pollination. Later application may be needed; tissue samples should be used to determine later usages.

Turfgrass (golf courses, sports fields, lawns, and other turf areas): Apply 0.25 to 2 fl. oz. per 1000 ft² in 1 to 3 gallons of water during the growing season as needed.

Other Crops: Apply according to directions of your regional Winfield Plant Nutrition Specialist or local agricultural professional to crops where the addition of copper would be beneficial.

For Turfgrass Applications: DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN, AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK ONTO YOUR LAWN/GARDEN.

Aerial Application:

Above use rates should be used in a minimum of 3 gallons of water by air. Repeat applications 10-21 days apart may be necessary.

A number of variables influence the rate of application necessary to prevent or correct a specific deficiency. For best results, follow good agronomic practices and tissue test. For specific recommendations for crops in your area, consult your state's recommendations. An ammonium sulfate source is recommended with herbicides where ammonium sulfate or a nitrogen source is recommended.

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.aapfco.org/metals.htm>

WARRANTY DISCLAIMER

Winfield Solutions, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions

for Use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.

SureTank™ is a trademark and Max-IN® is a registered trademark of Winfield Solutions, LLC.