

## Maximum N-PACT 12 - 0 - 0

GUARANTEED ANALYSIS Total Nitrogen (N)	
8.04% Urea Nitrogen	
3.96% Other Water Soluble Nitrogen*	
Boron (B)	5.00%

Derived from: Urea-Triazone Solution, Boric Acid

\*3.96% Slowly Available Nitrogen from Urea-Triazone Solution

WARNING: This product contains boron (B) and may be harmful to certain crops. Use only according to manufacturer's directions

### **KEEP OUT OF REACH OF CHILDREN**

# WARNING

May be harmful if swallowed. Causes skin irritation. Causes eye irritation. May cause respiratory irritation.

F1548

### FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL 1-866-944-8565.

### FOR FOLIAR / SOIL APPLICATION

### DIRECTIONS FOR USE Apply as needed based on soil tests or according to recommendations in your nutrient management plan.

APPLICATION INSTRUCTIONS				
CROPS (EPA GROUPING)	FOLIER APPLIED RATE	APPLICATION TIMING	COMMENTS	
Vegatable, Root and Tuber (01) Potato, Sugarbeet etc	1.0 to 6.0 pts/A	Apply prior to row canopy closing		
Leafy Veg (04) Celery Lettuce (Head, Leaf) etc	1.0 to 6.0 pts/A	Apply starting at thinning		
Brassica Veg (05) Broccoli, Cabbage, Cauliflower, Collards, Brussels Sprouts etc	1.0 to 6.0 pts/A	Apply at bottom establishment		
Legume Veg (06) All beans, Soybeans, etc	1.0 to 8.0 pts/A	Apply at early bloom		
Vegetable, Fruiting (08) Peppers, Tomato, Eggplant etc	1.0 to 6.0 pts/A	Apply starting at fruit formation		
Cucurbits (09) All melons, Cucumber, Squash etc	1.0 to 6.0 pts/A	Apply at bud formation to fruit establishment		
Citrus Fruits (10) Oranges, Lemons, Limes, etc	1.0 to 6.0 pts/A	Apply starting at fruit set		
Pome Fruit (11) Apple, Pear, etc	1.0 to 6.0 pts/A	Apply starting at fruit set		
Stone Fruit (12) Peach, Plum, etc	1.0 to 6.0 pts/A	Apply starting at fruit set		
Berry and Small Fruit (13) Strawberry, Blueberry, Caneberries (Blackberry, raspberry) etc	1.0 to 6.0 pts/A	Apply starting at fruit set		
Berry and Small Fruit (13) Grape	1.0 to 6.0 pts/A	Apply starting at fruit set through verasion		
Treenuts (14) Almond, Filbert, Pistachio, Pecan, Walnut etc	1.0 to 6.0 pts/A			
Cereal Grains (15) Corn, Wheat, Grain Sorghum, Rice, etc	1.0 to 6.0 pts/A	Apply from V5 to VT growth stage		
Grass forage Fodder and Hay (16) Grasses, etc.	1.0 to 6.0 pts/A			
Grass forage Fodder and Hay (17) Grass Grown for Seed	1.0 to 4.0 pts/A			
Animal Feed, Non Grass (18) All Alfalfa, Clover, etc	1.0 to 6.0 pts/A	Apply at crown green up or on regrowth after cutting		
Oilseed (20) Cotton, Sunflower, Canola etc	1.0 to 6.0 pts/A	Apply at bud to bolting stage		
Misc (99) Avocado, Artichoke, Hops, Flax, Mint, Olive, Peanut etc	1.0 to 4.0 pts/A			
Misc (99) Tobacco*	1.0 to 4.0 pts/A	Apply just prior to tasseling	*Do not use with alcohol based sucker control products	
			· · · · · · · · · · · · · · · · · · ·	

This product has been tested on a number of crop varieties with no phytotoxicity observed at labeled rates; however, this product has not been tested on all varieties and cultivars. Due to the large number of plant species and their associated varieties or cultivars, and due to variable growing conditions, it is impossible to test every plant for tolerance to this product. It is the responsibility of the user to determine if this product can be used safely prior to commercial use. In a small test area, test the specified rates on plants for phytotoxicity prior to general use. Before using any tank mix, test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. Treat a small group of target crop including plant cultivars and evaluate for 3 to 7 days prior to making an application to your entire crop.

This product is compatible with many agricultural pesticides. However, if specific information is not available, a jar test for compatibility should be conducted prior to large scale mixing.

thoroughly after handling this product.

## For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary. LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the lobal and is macroached if for the purpose of the lobal integration for the purpose the product is used in according to the product is used in according to the purpose. the label and is reasonably fit for the purposes stated in the Directions for Use, when the product is used in accordance with such Directions for Use under normal conditions of use. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

California Proposition 65: WARNING: This product contains chemicals known to the State of California to cause cancer or birth defects or other reproductive harm.

info-lpi.com

the goods are compatible with all other chemicals or under all use conditions. Maximum N-Pact is a registered trademark of Loveland Products, Inc.

Density: 10.51 lbs/gallon @ 68 °F

Liquid Measure: 250 gallons (946.35 liters) Net Weight: 2,627.50 lbs (1,191.81 kg) Product Code: 1000798813

Density: 10.51 lbs/gallon @	≥ 68 °F		
Liquid Measure:	gallons (		liters)
Net Weight:	lbs (	kg)	
Product Code: 1000798814	4	•	

### Crop Safety

### Tank Mix Compatibility

### STORAGE AND DISPOSAL

**STORAGE:** Store in original containers only. Keep containers tightly closed when not in use. Store in a cool, dry well-ventilated area, preferably in a locked storage area away from children, feed and food products, and seed. Store at ambient temperatures. Do not contaminate water, food, or feed by storage or disposal.

DISPOSAL: Do not reuse container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. Do not contaminate water, food, or feed by storage or disposal.

HANDLING: Avoid breathing fertilizer dusts. Avoid prolonged or repeated skin contact. Avoid eye contact. Wash hands

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that