

GROUP

4

HERBICIDE

MCPA Amine

ACTIVE INGREDIENT:

Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid *	48.72%
INERT INGREDIENTS:	51.28%
Total	100.00%

* Equivalent to 39.72% 2-methyl-4-chlorophenoxyacetic acid. Contains 3.70 lbs. of MCPA acid per gallon.
Isomer specific by AOAC Method No. 6.A18-22 (13th Edition)

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle
(If you do not understand the label, find some one to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eyelids open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

For additional information in case of emergency call toll free 1-877-424-7452.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage. Wear protective eyewear (goggles, face shield, safety glasses). Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin, or clothing. Avoid contact with skin or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are any waterproof material.

Mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves,
- Shoes plus socks, and
- Protective eyewear (goggles, face shield, safety glasses).

EPA Reg. No. 1381-104

EPA Est. No. 70989-IA-1

Distributed by
Winfield Solutions, LLC
P.O. Box 64589, St. Paul, MN 55164-0589

NET CONTENTS 2-1/2 GALS.

1/0821/7

Additional PPE requirements for mixers and loaders supporting aerial application to rangelands, pasture lands, or noncropland. These mixers/loaders also must wear:

- A chemical-resistant apron and
- A NIOSH approved particulate filter with any N, R, P filter with NIOSH approval number prefix TC-84A; or a NIOSH-approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C. It is recommended that the respirator wearer be fit tested and trained in the use, maintenance, and limitations of the respirator.

See Engineering Controls for additional requirements.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning or maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering controls statement:

When handlers use closed systems, enclosed cabs, or aircraft, in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates and aquatic plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated area. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA Amine have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA Amine pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Use care to avoid spray contact or drift to susceptible plants such as beans and other legumes, cotton, flowers, grapes, ornamental, vegetables, and other plants. Do not permit spray mist containing this product to drift onto them, since even very small quantities of the spray, which may not be visible, can cause severe injury during both growing and dormant periods.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements of this box apply only to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls,
- Chemical resistant gloves made of any waterproof material,
- Shoes plus socks, and
- Protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

ANNUAL AND BIENNIAL WEEDS

Beggarticks, bull thistle, burdock (common), burhead, cocklebur, *goatsbeard, *hempnettle, jimsonweed, kochia, *ladysthumb, lambsquarter, *marshelder, mustards (except blue), pennycress (fanweed), pepperweed (field), pigweed (tumble, prostrate, smooth), poison hemlock, prickly lettuce, *puncturevine, ragweed (common, giant), redstem, *shepherdspurse, sicklepod, stinkweed, *sunflower, *velvetleaf, Venice mallow, *wild carrot, wild radish, witchweed.

PERENNIAL WEEDS*

Arrowhead, bulrush, bindweed (hedge, field), buttercup (corn, creeping, hairy), Canada thistle, curly indigo, dandelion, dragonhead mint, **hempnettle, hoary cress (white top), honeysuckle, plantain, rush (soft), smartweed, **Texas blueweed, water plantain, **whitebrush, wild garlic, wild sage (Sonoma), **wild onion, yellow rocket.

*Use the higher rate for control. Repeat treatments may be necessary. Refer to specific crop Restrictions and Limitations.

**Regrowth may occur after repeat application.

FOR AERIAL APPLICATION: Unless noted otherwise under individual **DIRECTIONS** section, for aerial application, apply the specified amount in a minimum of 2 gallons of water per acre.

FOR GROUND APPLICATION: Apply the specified amount in a minimum of 10 gallons of water per acre. Use more water for both methods when adverse growing conditions are present. **DO NOT** apply with high spray pressures, hollow cone or other nozzle types that produce small spray droplets which may drift. Avoid spray drift by making applications when conditions such as wind, air stability and temperature inversions are not a factor. The use of a suitable drift control agent at the proper rate will aid in the reduction of spray drift.

Do not apply this product through any type of irrigation system.

PLANTBACK INTERVAL

Wheat, barley, oats, rye, flax, and peas treated with MCPA may be replanted with any crop specified on an MCPA label or any crop for which a residue tolerance exists for MCPA. For crops not listed on an MCPA label, or on crops for which no residue tolerances for MCPA have been established, a 60 day plantback interval must be observed.

RESISTANCE MANAGEMENT RECOMMENDATIONS

MCPA Amine is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to MCPA Amine or other Group 4 herbicides. Weed species with acquired resistance to Group 4 may eventually dominate the weed population if Group 4 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by MCPA Amine or other Group 4 herbicides.

To delay herbicide resistance consider:

- Avoiding the consecutive use of MCPA Amine or other target site of action Group 4 herbicides that have a similar target site of action, on the same weed species.
- Using tank-mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive IPM program.
- Monitoring treated weed populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or Winfield Solutions, LLC representative for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Additional requirements for aerial application:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

MIXING INSTRUCTIONS

WATER BASED SPRAY -- Fill the equipment half full of water, agitate while adding this product, then add the rest of the water.

NITROGEN FERTILIZER: Weed and feed applications for small grains, grasses grown for seed or grass pastures according to label use rates. Add half the fertilizer to the tank, then add ½ to ¾ pint of **MCPA Amine** per acre. Agitate constantly and vigorously and finish filling the spray tank with fertilizer. Apply as soon as possible, agitating constantly. Do not hold spray mixture overnight. If incompatibility is a problem, the use of a compatibility agent at the specified label rate may correct the problem. Fertilize according to the recommendations of your supplier-or your Extension specialist. Herbicide foliage contact burning may occur as a result of fertilizer use. Lower use rates and concentrations will reduce this problem.

Wash spray equipment thoroughly with a tank cleaner after using this product. When cleaning equipment, do not pour washwater on the ground: spray or drain over a large area away from wells or other water sources.

Note: When stored at temperatures below freezing, it may be necessary to warm contents to 70° F and mix thoroughly before using.

SPRING PLANTED SMALL GRAINS - Oats, Barley, Rye, and Wheat Not Underseeded With Legumes

WEEDS	Amount of MCPA Amine Per Acre	DIRECTIONS
Annual weeds Annual and biennial weeds Perennial weeds	½ pint 1 to 1-5/8 pints* 1½ to 1-5/8 pints*	Apply when weeds are small. Apply to grain at 3 to 5 leaf stage or up to early boot stage (Zadoks 1 ₃ to 4 ₀). Do not apply during boot to dough stage (Zadoks 4 ₀ to 8 ₃).

FALL PLANTED SMALL GRAINS - Oats, Barley, Rye, and Wheat Not Underseeded With Legumes

Annual and biennial broadleaf weeds Perennial weeds	½ pint ½ to 1 pint*	Apply in early spring when weeds are small and grain is fully tillered, but before boot stage (Zadoks 2 ₅ to 4 ₀).
--	------------------------	---

RESCUE TREATMENT - Wheat

Perennial broadleaf weeds	1-5/8 pints**	For weed suppression, treat in spring after grain is tillered and at least 8 to 10" tall and when weeds are approaching bud stage, but do not spray when grain is in boot to dough stage (Zadoks 4 ₀ to 8 ₃).
---------------------------	---------------	--

* The higher rate may cause grain injury.

** The 1-5/8 pint rate can produce injury to wheat. Use good judgment to determine the severity of weed problem against the possibility of crop damage. When weeds are scattered, spot treatment will lessen damage.

RESTRICTIONS AND LIMITATIONS FOR SMALL GRAINS

Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not apply more than 1-5/8 pints (3/4 lb. ae) MCPA Amine per acre per year.

ESTABLISHED ALFALFA

WEEDS	Amount of MCPA Amine Per Acre	DIRECTIONS
Yellow rocket, pennycress (fanweed) and other weeds	1 pint	Apply in late fall following frost or in early spring when crop is dormant. The temperature at the time of spraying should be above 40 ° F.

RESTRICTIONS AND LIMITATIONS FOR ALFALFA

Do not forage or graze treated areas for 7 days after treatment.

Do not apply more than 1 pint (1/2 lb. ae) per acre per year.

SEEDLING CLOVER ALONE, OR UNDERSEEDED WITH GRAIN AND ESTABLISHED STANDS OF ARROWLEAF, CRIMSON, RED, LADINO, AND WHITE CLOVER (EXCEPT DUTCH)

WEEDS	Amount of MCPA Amine Per Acre	DIRECTIONS
Seedling clovers (except Dutch) Established stands	½ pint ¾ to 1 pint	Treat seedling stands after they have 2 or more true leaves (trifoliate stage). Apply in late fall after frost or early spring when weeds are small. Temperature at spraying should be above 40° F.

FOR USE IN THE PACIFIC NORTHWEST ONLY. Do not apply to sweet clover. Observe drift precautions. Stem twisting will show up for a short time after application. Some growth reduction may occur.

RESTRICTIONS AND LIMITATIONS FOR CLOVERS

Do not forage or graze treated areas for 7 days after treatment.

Do not apply more than 1 pint (1/2 lb. ae) per acre per year.

PASTURES, RANGELAND, CONSERVATION RESERVE PROGRAMS AND SET-ASIDE ACRES

WEEDS	Amount of MCPA Amine Per Acre	DIRECTIONS
Newly seeded clover and grass pastures	½ pint	Treat seedling stands after they have 2 or more true leaves (trifoliate stage).
Established pastures Annual, biennial and perennial weeds	1 to 3¼ pints*	Apply when weeds are small and growing. Spray in spring and fall. Spray perennial in early bud to full bloom and regrowth in fall. Use higher rate for hoary cress (white top), Canada thistle, buttercup, and other hard to control weeds.
Whitebrush	3 pints*	Apply in 1 gallon of diesel oil and 6 to 10 gallons of water per acre by aircraft. Apply in spring or fall when foliage is fully developed and growing. Spray during bloom, but not after blossom shed. Retreatment may be necessary. NOTE: Add MCPA Amine to the water, then add the oil while agitating. Continue to agitate while spraying.

* Do not spray legumes, if present, unless injury can be tolerated.

RESTRICTIONS AND LIMITATIONS FOR PASTURES, RANGELAND - Do not forage or graze treated areas for 7 days after treatment.

CONSERVATION RESERVE PROGRAMS AND SET-ASIDE ACRES - Do not forage or graze at any time.

RESTRICTIONS AND LIMITATIONS FOR PASTURES, RANGELAND, CONSERVATION RESERVE PROGRAMS AND SET-ASIDE ACRES - Do not apply more than 3¼ pints (1½ lbs. ae) MCPA Amine per acre per application. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

FALLOW LAND AND CROP STUBBLE

WEEDS	Amount of MCPA Amine Per Acre	DIRECTIONS
Canada thistle, hoary cress(white top) and Texas blueweed	3¼ pints	Apply when weeds are young and continue treatment for 2 to 3 years for the best results.

NON-CROP AREAS - Airfields, Drainage Ditches, Fence rows, Rights of Way (Roadways, Utility, Pipeline, Railroad, Highway) Industrial Sites, Vacant Lots

WEEDS	Amount of MCPA Amine Per Acre	DIRECTIONS
Canada thistle, meadow buttercup, hoary cress (white top), and other hard-to-control weeds.	3¼ pints	Apply in a minimum of 2 gallons of water per acre for aerial application and 20 to 100 gallons of water for ground application. Spray the weeds thoroughly when they are in bud to early bloom and again during fall.
Spot treatment	1/4 pint	Apply in 3 to 4 gallons of water.

RESTRICTIONS AND LIMITATIONS FOR NON-CROP AREAS - Do not forage or graze treated areas for 7 days after treatment. Broadcast Treatment: Do not apply more than 3¼ pints (1½ lbs. ae) MCPA Amine per acre per application. Do not apply more than 2 applications per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days. Spot Treatment: Do not apply more than 6.48 pints (3 lbs. ae) MCPA Amine per acre per year.

GRASSES GROWN FOR SEED - Do not use on alfalfa, clover, other legumes or newly seeded areas.

WEEDS	Amount of MCPA Amine Per Acre	DIRECTIONS
Annual, biennial and perennial weeds	1 to 2 pints*	Apply to established grasses before the head comes into boot. Apply to seedling grasses after tillering. Use the higher rate for heavier weed stands.

* Legumes present may be injured or killed. Bent, Buffalo, Carpet and St. Augustine grasses may be injured by this treatment. NOTE: Repeat treatment may be needed.

RESTRICTIONS AND LIMITATIONS FOR SEED GRASSES - Do not forage or graze treated areas for 7 days after treatment. Do not apply more than 3 ¼ pints (1½ lbs. ae) MCPA Amine per acre per application. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

FLAX

WEEDS	Amount Per Acre	DIRECTIONS
Annual and biennial weeds	1/3 to ½ pint	Apply when flax is 2 to 8 inches tall. Do not spray once flax has reached the bud stage. Use at least 8 to 10 gallons of water per acre for ground application and at least 3 to 5 gallons of water per acre for aerial application.

RESTRICTIONS AND LIMITATIONS FOR USE ON FLAX – Do not allow livestock animals to forage or graze treated areas within 7 days of slaughter. Do not apply more than ½ pint (1/4 lb. ae) MCPA Amine per acre per year.

PEAS

WEEDS	Amount Per Acre	DIRECTIONS
Annual and biennial weeds	½ to ¾ pint	Use only in the WA, OR and ID. Treat when peas are 4 to 6 inches tall. Do not spray while peas are in blossom. Peas may show slight injury, but they usually recover a few days after treatment. Use the higher specified rate before buds form on Canada thistle to prevent from going to seed. Do not spray peas that are stressed from lack of moisture or when temperatures are over 90° F. Apply in 15 to 30 gallons of water per acre for ground applications and 5 gallons of water per acre for aerial application.

RESTRICTIONS AND LIMITATIONS FOR USE ON PEAS – Do not allow livestock to forage or graze treated pea fields. Do not feed treated vines to livestock. Do not treat bean-type peas such as crowder, blackeye, etc. Do not apply more than ¾ pint (3/8 lb. ae) per acre per year.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a secure area, in original container only. Do not store near feed or foodstuffs.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Use label language appropriate for container size and type.

Nonrefillable containers. Do not reuse or refill this container. Clean container promptly after emptying.

Nonrefillable container equal to or less than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities such as burning of plastic containers. If burned, stay out of smoke.

Nonrefillable container greater than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other

procedures approved by state and local authorities such as burning of plastic containers. If burned, stay out of smoke. **Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities such as burning of plastic containers. If burned, stay out of smoke.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call
CHEMTREC 1-800-424-9300.**

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF

MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.