

METOLACHLOR	GROUP 15	HERBICIDE
METRIBUZIN	GROUP 5	HERBICIDE

Drexel®

Me-Too-Lachlor™ MTZ

Herbicide

For control of certain grasses and broadleaf weeds in Potatoes and Soybeans.*

ACTIVE INGREDIENTS:

Metolachlor	58.2%
Metribuzin	13.8%

OTHER INGREDIENTS:	28.0%
TOTAL:	100.0%

This product contains 5.25 pounds of Metolachlor and 1.25 pounds of Metribuzin per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

[See FIRST AID Below]

[See Side (Back) Panel for FIRST AID];

[See Page ____ for FIRST AID]

[See Container Labeling for (FIRST AID and Complete Directions for Use]

[See (Attached) Booklet (Container Labeling) for Complete Directions for Use]

EPA Reg. No. 19713-704

Net Content:

EPA Est. No. 19713-MS-1

____ Gals. (____ L)

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give any liquid to the person.
- Do not give anything to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN:

Contains petroleum distillates. May pose an aspiration pneumonia hazard. Vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call CHEMTREC at 800-424-9300 for emergency medical treatment information.

* Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING: Causes temporary but substantial eye injury. Harmful if swallowed. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: Protective eyewear, coveralls over short-sleeved shirt and short pants, chemical-resistant gloves made of barrier laminate, butyl rubber, nitrile rubber or Viton (all 2: 14 mils), chemical-resistant footwear plus socks, chemical-resistant headgear for overhead exposure and chemical-resistant apron when mixing/loading and cleaning equipment. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Reporting Ecological Incidents:

To report ecological incidents, including mortality, injury, or harm to plants and animals, call 1-901-774-4370.

Groundwater Advisory

The active ingredient, Metolachlor, is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Manufactured By:

Drexel Chemical Company

P.O. Box 13327, Memphis, TN 38113-0327

SINCE 1972

ME-TOO-LACHLOR, DE-ESTER, FOMA, FOMASATE, IMITATOR, MIX, PEPTOIL, PIN-DEE, QUIK-QUAT, SURF-AC and the DREXEL logo are either trademarks or registered trademarks of Drexel Chemical Company. All other brand names, product names or trademarks belong to their respective holders.

704SP-0123*

ME-TOO-LACHLOR MTZ Page 1 of 13

Metribuzin is a chemical which can travel (seep or leach) through soil and can contaminate groundwater which may be used as drinking water. Metribuzin has been found in groundwater as a result of agricultural use. Users are advised not to apply Metribuzin where the water table (groundwater) is close to the surface and where the soils are very permeable, i.e., well-drained soils such as *Loamy sands*. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several weeks or months after application. A level, well maintained, vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential loading of Metolachlor from runoff water and sediment. Runoff of this product will be reduced by avoiding application when rainfall or irrigation is expected to occur within 48 hours.

Mixing and Loading Instructions

Take care when using this product to prevent back-siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

Check-valves or anti-siphoning devices must be used on mixing equipment.

This product may not be mixed/loaded or used within 50 feet of wells, including abandoned wells, drainage wells and sink holes. Operations that involve mixing, loading, rinsing or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

NON-TARGET ORGANISMS ADVISORY STATEMENT

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL AND CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

ENDANGERED SPECIES PROTECTION REQUIREMENTS:

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g. kill or otherwise harm) of an endangered species and certain threatened species, under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <https://www.epa.gov/endangered-species>, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 12 hours. Exception: If the product is soil-injected or soil incorporated, the WPS, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is: Protective eyewear, coveralls over short-sleeved shirt and short pants, chemical-resistant gloves made of barrier laminate, butyl rubber, nitrile rubber or viton (all 2:14 mils), chemical-resistant footwear plus socks and chemical-resistant headgear for overhead exposure.

New York State: Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY OR ILLEGAL RESIDUES.

HERBICIDE RESISTANCE MANAGEMENT

METOLACHLOR	GROUP	15	HERBICIDE
METRIBUZIN	GROUP	5	HERBICIDE

For resistance management, this product is a Group 15 and 5 modes of action herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 15 and 5 modes of action herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 15 and 5 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank-mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance management and/or integrated weed management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Drexel Chemical Company representatives at (901)774-4370.

PRODUCT INFORMATION

Observe all precautions and limitations on the labels of each product used in tank-mixtures. Tank-mixture partners must be registered in states where they are used. Refer to and follow the label for each tank-mix product used. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

Do not apply under conditions which favor runoff or wind erosion of soil containing this product to non-target areas.

To prevent off-site movement due to runoff or wind erosion:

- Avoid treating powdery dry or light *Sand soils* when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
- Do not apply to impervious substrates such as paved or highly compacted surfaces.
- Do not use tail water from the first flood or furrow irrigation of treated fields to treat non-target crops, unless at least 0.5 inch of rainfall has occurred between application and the first irrigation.

MIXING INSTRUCTIONS

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean the spray equipment before using this product. Vigorous agitation is necessary to maintain uniformity of the spray mixture. Maintain maximum agitation throughout the spraying operation. Do not allow spray mixture to stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

APPLICATION IN WATER OR FLUID FERTILIZERS

This Product Alone: Add one-third of the required amount of water or fluid fertilizer to the spray or mixing tank. With the agitator running, add this product into the spray tank. Continue agitation while adding the remainder of the water or fluid fertilizer. Begin application of the spray solution after this product has completely dispersed in the water or fluid fertilizer. Maintain agitation until all of the mixture has been applied.

This Product + Tank-Mixtures: Add one-third of the required amount of water or fluid fertilizer to the mix tank. Start the agitator running before adding any tank-mix partners. In general, tank-mix partners should be added in this order:

- 1) Products packaged in water-soluble packaging
- 2) Wettable powders
- 3) Wettable granules (dry flowables)
- 4) Liquid flowables
- 5) Liquids such as this product, and
- 6) Emulsifiable concentrates

Always allow each tank-mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

(1) When using this product in tank-mixtures, all products in water-soluble packaging should be added to the tank and mixed with plain water before any other tank-mix partner, including this product. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix partner to the tank. (2) Water-soluble packets will not properly dissolve in most spray solutions that contain fluid fertilizers.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

This product is compatible with most common tank-mix partners. However, if not known, the physical compatibility of this product with tank-mix partners should be tested before use. To determine the physical compatibility of this product with other products, use a jar test, as described below.

COMPATIBILITY TEST

A jar test is recommended before tank-mixing to ensure compatibility of this product with other pesticides. The following test assumes a spray volume of 25 gallons per acre. For other spray volumes, make appropriate changes in the ingredients.

Note: Nitrogen solutions or complete fluid fertilizers may replace all or part of the water in the spray. Because liquid fertilizers vary, even within the same analysis, **always check compatibility with pesticide(s) before use.** Incompatibility of tank-mixtures is more common with suspensions of fertilizer and pesticides.

Test Procedure

1. Add 1.0 pint of carrier (fertilizer or water) to each of 2 one quart jars with tight lids. **Note:** Use the same source of water that will be used for the tank-mix and conduct the test at the temperature the tank-mix will be applied.
2. To one of the jars, add 0.25 teaspoon or 1.2 milliliters of a compatibility agent approved for this use such as Mix™, Compex® or Unite® (0.25 tsp. is equivalent to 2.0 pts./100 gal. spray). Shake or stir gently to mix.
3. To both jars, add the appropriate amount of pesticide(s) in their relative proportions based on label rates. If more than one pesticide is used, add them separately with dry pesticides first, flowables next and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix.
4. After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixtures stand 15 to 30 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (a) Slurry the dry pesticide(s) in water before addition, or (b) add one-half of the compatibility agent to the fertilizer or water and the other half to the emulsifiable concentrate or flowable pesticide before addition to the mixture. If incompatibility is still observed, do not use the mixture.
5. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the "STORAGE AND DISPOSAL" section in this label.

GROUND APPLICATION

Apply this product alone or in tank-mixtures by ground equipment in a minimum of 10 gallons of spray mixture per acre, unless otherwise specified. For certain ground application equipment, apply in a minimum of 2 gallons of spray mixture per acre. Contact your local manufacturer's representative for a list of this type of equipment.

Use sprayers that provide accurate and uniform application. Calibrate the sprayer before use at the beginning of the season. For tank-mixtures of this product with wettable powder or dry flowable formulations, screens and strainers should be no finer than 50 mesh. Calculate the amount of herbicide needed for band treatment by the formula:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{broadcast rate}}{\text{per acre}} = \frac{\text{Amount needed}}{\text{per acre}}$$

CENTER PIVOT IRRIGATION APPLICATION

If applying this product by chemigation, apply only through a center pivot irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact your State Extension specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Operating Instructions

- The system must contain a functional check-valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump or piston pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Prepare a mixture with a minimum of 1 part water to 1 part herbicide(s) and inject this mixture into the center pivot system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension.
- Meter into irrigation water during entire period of water application.
- Apply in 1.5 inches of water. Use the lower water volume (0.5 inch) on *Coarse textured soils* and the higher volume (1 inch) on *Fine textured soils*. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil.

Use Precautions for Center Pivot Applications: Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result. Where sprinkler distribution patterns overlap excessively, crop injury may result.

AERIAL APPLICATION

Apply this product in water using a minimum spray volume of 2 gallons per acre. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

MANDATORY SPRAY DRIFT MANAGEMENT

GROUND BOOM APPLICATIONS:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 ft. above the ground or crop canopy.
- Applicators are required to select the nozzles and pressure that deliver medium or coarser droplets (ASABE S572).
- Do not apply when wind speeds exceed 15 mph at the application site.
- Do not apply during temperature inversions.

BOOMLESS GROUND APPLICATIONS:

- Applicators are required to select the nozzle and pressure that deliver medium or coarser droplet size (ASABE S572.3) for all applications.
- Do not apply when wind speeds exceed 15 mph at the application site.
- Do not apply during temperature inversions.

AERIAL APPLICATIONS:

- Do not release spray at a height greater than 10 ft. above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select the nozzle and pressure that deliver medium or coarser droplets (ASABE S641).
- If the wind speed is 10 mph or less, applicators must use one-half swath displacement upwind at the downwind edge of the field. When the wind speed is between 11 to 15 mph, applicators must use three fourths swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 15 miles per hour at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.

Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind.

The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

BOOMLESS GROUND APPLICATIONS

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

HANDHELD TECHNOLOGY APPLICATIONS

Take precautions to minimize spray drift.

CLEANING EQUIPMENT AFTER APPLICATION

Because some non-labeled crops are sensitive to low rates of this product, special attention must be given to cleaning equipment before spraying a crop other than those registered for use and on this label. Mix only as much spray solution as needed. Immediately after spraying, clean equipment thoroughly using the following procedure:

1. Flush tank, hoses, boom and nozzles with clean water.
2. Prepare a cleaning solution of 1 gallon of household ammonia per 50 gallons of water. Many commercial spray tank cleaners may be used as well. Consult your Manufacturer's representative for a partial listing of approved tank cleaners and more information about proper tank cleaning procedures. Do not use chlorine-based cleaners such as Clorox®.
3. When available, use a pressure washer to clean the inside of the spray tank with this solution. Take care to wash all parts of the tank, including the inside top surface. Completely fill the sprayer with the cleaning solution to ensure contact of the cleaning solution with all internal surfaces of the tank and plumbing. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for **at least 15 minutes**. All visible deposits must be removed from the spraying system.
4. Flush hoses, spray lines and nozzles for at least one minute with the cleaning solution.
5. Dispose of rinsate from steps 1 to 3 as described under the "ENVIRONMENTAL HAZARDS" section.
6. Repeat steps 2 to 5.
7. Remove nozzles, screens and strainers and clean separately in the ammonia cleaning solution after completing the above procedures.
8. Rinse the complete spraying system with clean water.

IMPREGNATION ONTO DRY BULK GRANULAR FERTILIZERS

Many dry bulk granular fertilizers may be impregnated or coated with this product and used to control weeds. When applying this product with dry bulk fertilizers, follow all directions for use and precautions on this label regarding target crops, rates per acre, soil texture, application methods and rotational crops.

Complying with all individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling and application is the responsibility of the individual and/or company selling the herbicide/fertilizer mixture.

Prepare the herbicide/fertilizer mixture by using any closed drum, belt, ribbon or other commonly used dry bulk fertilizer blender. Nozzles used to spray this product onto the fertilizer must be spaced

to provide uniform spray coverage. Care should be taken to aim the spray onto the fertilizer only, avoiding the walls of the blender. If the herbicide/fertilizer mixture is too wet, add a highly absorptive material such as Agsorb® FG or Celatom MP-79® or similar granular clay or diatomaceous earth materials, to obtain a dry, free-flowing mixture. Absorptive materials should be added only after the herbicide has been thoroughly blended into the fertilizer mixture. Best application results will be obtained by using a granule of 6/30 particle size or of a size similar to that of the fertilizer materials being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% absorptive material by weight. Calculate the amount of this product to be used by the following formula:

$$\frac{2,000}{\text{Lbs. of Fertilizer per Acre}} \times \frac{\text{Pt. of This Product}}{\text{per Acre}} = \frac{\text{Pt. of This Product}}{\text{per Ton of Fertilizer}}$$

PNEUMATIC (COMPRESSED AIR) APPLICATION

High humidity, high urea concentrations, low fertilizer use rates and dusty fertilizer may cause fertilizer mixtures to build up or plug the distributor head, air tubes or nozzle deflector plates. To minimize buildup, pre-mix this product with Exxon Aromatic 200 at a rate of 2.0 to 2.5 pints per gallon of this product. Aromatic 200 is a noncombustible/nonflammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Drying agents should not be used when using Aromatic 200. Mixtures of this product and Aromatic 200 must be used on dry fertilizer only. Poor results or crop injury may result if these mixtures are used in water or liquid fertilizer solutions for spraying applications. When impregnating this product in a blender before application, a drier mixture can be obtained by substituting a drying agent for Aromatic 200. The use of Agsorb FG or another drying agent of 6/30 particle size is recommended. Drying agents are not recommended for use with On-The-Go impregnation equipment.

Use Restrictions: To avoid potential for explosion, (1) Do not impregnate this product on ammonium nitrate, potassium nitrate or sodium nitrate, either alone or in blends with other fertilizers. (2) Do not combine this product with a single superphosphate (1-20-0) or treble superphosphate (0-46-0). (3) Do not use this product on straight limestone, since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated.

APPLICATION OF IMPREGNATED DRY BULK GRANULAR FERTILIZERS

Apply 200 to 700 pounds of the herbicide/fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential in order to prevent possible crop injury to subsequent rotational crops. Non-uniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil is recommended to obtain satisfactory weed control. On *Fine* or *Medium textured soils* in areas where soil incorporation is not planned, i.e., reduced-tillage situations or in some conventional till situations, make applications approximately 30 days before planting to allow moisture to move the herbicide/fertilizer mixture into the soil. On *Coarse textured soils*, make applications approximately 14 days prior to planting.

Use Precautions: To help avoid rotational crop injury, make applications as early as possible, since this product impregnated onto dry bulk fertilizers can be expected to last longer in the soil than this product applied as a spray in water or fluid fertilizer.

CROP ROTATION INTERVALS^{1,3}

Rotational Interval After Application of This Product ²				
4 Months	4.5 Months	8 Months	12 Months	18 Months
Corn	Alfalfa Winter barley Winter wheat	Peas Rice Spring barley Spring wheat	Asparagus Cotton Forage grasses Lentils Sainfoin Sugarcane Tomatoes Other crops not listed (except Root crops)	Onions Sugar beets and Other Root crops

¹ Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed. Stand reductions may occur in some areas.

² Crop rotation directions do not include restrictions for the tank-mix partner. Refer to the label of the other product for additional restrictions.

³ Refer to the "CROPS" section for additional crop rotation precautions.

Replanting

If replanting is necessary in fields previously treated with this product, the field may be replanted to Potatoes or Soybeans. Before replanting, refer to the "CROPS" sections for directions, precautions and restrictions.

Activation

A small amount of rainfall or irrigation is required to activate this product following application. In areas of low rainfall, a pre-emergence application should be followed by light irrigation of 0.25 to 0.5 inch of water. Do not apply heavy irrigation immediately after application. As with many surface-applied herbicides, weed control and crop tolerance may vary with rainfall and/or soil texture.

CROPS

POTATOES*

This product may be used for pre-emergence weed control prior to or after Potato emergence. This product has some post-emergence activity on weeds, but the consistency and spectrum of weed control is much better pre-emergence to weeds. Pre-plant incorporated applications are not recommended due to an increased risk of crop injury.

PRE-EMERGENCE APPLICATIONS

Apply with ground spray equipment, aerial spray equipment or by center pivot irrigation equipment which is capable of making a uniform broadcast application. Apply after planting but before crop emergence or apply after drag-off if this operation is part of the usual cultural practice.

POST-EMERGENCE APPLICATIONS

Apply this product post-emergence only in center pivot irrigation water, after drag-off if that is a usual cultural practice, but not closer than 60 days before harvest. Refer to the "CENTER PIVOT IRRIGATION APPLICATION" section of this label for application information.

*Not registered for use in California.

WEEDS CONTROLLED BY THIS PRODUCT

Annual Broadleaves*		
Anoda, Spurred Beggarweed, Florida Carpetweed Chickweed, Common Cocklebur** Copperleaf, Hophornbeam <i>Galinsoga</i> spp. Henbit Jimsonweed Knotweed, spp. Kochia** Ladysthumb	Lambsquarters, Common Lettuce, Prickly Mallow, Venice Mustard, spp. Nightshade, Black Nightshade, Hairy** Pennycress, Field Pepperweed, Virginia Pigweed, spp. Purslane, Common Pusley, Florida Ragweed, Common**	Redweed <i>Sesbania</i> spp. Shepherdspurse Sicklepod Sida, Prickly/ Teaweed Smartweed, Pennsylvania Spurge, Spotted Starbur, Bristly Sunflower, Common** Thistle, Russian Velvetleaf** Waterhemp, spp.
Annual Grasses		
Barnyardgrass Bluegrass, Annual Crabgrass, spp. Crowfootgrass Cupgrass, Prairie Cupgrass, Southwestern	Foxtail, spp. Goosegrass Johnsongrass, Seedling** Junglerice Panicum, Fall Panicum, Texas**	Red rice Sandbur, spp.** Shattercane** Signalgrass, Broadleaf Witchgrass
Sedges		
Yellow nutsedge	-	-
Volunteer Crops		
Barley**	Sorghum**	Wheat**

*Except Triazine resistant biotypes other than Black nightshade, *Galinsoga* spp., Pigweed, spp. and Waterhemp, spp.

**Suppression only. Suppression means significant activity, but not always at a level considered acceptable for commercial weed control.

APPLICATION RATES

The application rates of this product for use in Potatoes are provided below. Where a rate range is given, use the lower end of the rate range on the more *Coarse textured soils* listed within that group and/or where weed pressures are known to be light; use the high end of the rate range on the more *Fine textured soils* listed within that group and/or where the weeds pressures are known to be heavy.

Pre-emergence Use Rates in Potatoes - This Product (Pt./Ac.)		
Soil Texture	0.5 to 3% Organic Matter	> 3% Organic Matter
COARSE*: Sand, Loamy sand, Sandy loam	1.5 to 2.0	2.0 to 2.4
MEDIUM or FINE: Loam, Silt loam, Silt, Sandy clay, Sandy clay loam, Silty clay, Silty clay loam, Clay, Clay loam	2.4 to 2.75	2.75 to 2.9
*On soils that classify as a Sand texture, do not use more than 1.5 pt./Ac. of this product or more than 0.5 lb. Metribuzin a.i./Ac. in total or crop injury may occur.		

Post-emergence Use Rates in Potatoes - This Product (Pt./Ac.)*	
Soil Texture	0.5% Organic Matter
COARSE**: Sand, Loamy sand, Sandy loam	1.5
MEDIUM or FINE: Loam, Silt loam, Silt, Sandy clay, Sandy clay loam, Silty clay, Silty clay loam, Clay, Clay loam	1.5 to 2.2
*For application in center pivot irrigation water only.	
**Crop injury may occur on soils that classify as a Sand texture and have less than 0.5% organic matter.	

USE PRECAUTIONS

- To avoid crop injury, post-emergence applications should be made only on russeted or white skinned varieties of Potatoes that are not early maturing. Avoid post-emergence applications on Atlantic, Bellchip, Centennial, Chipbelle, Shepody and Superior varieties. Pre-emergence applications on these varieties may cause crop injury under adverse weather conditions, on *Coarse soils*, under high soil pH and with higher use rates.
- Potato varieties may vary in their response to a given herbicide application. When using this product for the first time on a particular variety, always determine crop tolerance before using on a field scale.
- The planting of sensitive crops such as Cole crops, Cucurbits and Lettuce during the next growing season following application of this product may result in injury to that crop.
- Certain Cereal varieties are sensitive to Metribuzin (e.g., see section on Cereals of the Sencor® 4 or Sencor DF label) and should not be planted during the next growing season unless the following cultural practices occur:
 - Potato vines left in the row as a result of harvest must be uniformly distributed over the soil surface prior to plowing, and
 - Plow with a moldboard plow to a depth sufficient to mix the upper 8 inches of soil.
- Crop injury may occur if this product is applied as pre-plant incorporated in Potatoes.

USE RESTRICTIONS

- Two applications may be applied per year. For Potatoes grown in soils with organic matter between 3% and 10%, do not apply more than 5.1 pints (3.35 lbs. Metolachlor a.i.) per acre per year; and in soils with organic matter between 0.5% and 3.0%, do not apply more than 4.95 pints (3.25 lbs. Metolachlor a.i.) per acre per year. Do not apply more than 1.0 pound Metribuzin a.i. per acre per year. This product is not to be used for application to *Muck* or *Peat soils*.
- Do not apply this product post-emergence if the weather in the next 3 days is predicted to be cool, wet or cloudy, as crop injury may occur.
- Pre-Harvest Interval (PHI): Do not harvest within 60 days of the last application of this product.
- Do not apply after June 30 in Idaho, Oregon or Washington if the treated land will be planted to a crop other than Potatoes in the Fall.
- Do not apply this product to Sweet potatoes or Yams.

THIS PRODUCT IN TANK-MIXTURE FOR USE IN POTATOES

For pre-emergence applications in Potatoes, this product may be tank-mixed with other pesticide products registered for use in this way and timing in Potatoes. If you have no previous experience mixing these products under your conditions, perform a compatibility test before attempting large-scale mixing (see "COMPATIBILITY TEST" section of this label).

For post-emergence applications (center pivot irrigation applications only), i.e., where Potato vines are exposed, there may be increased risk of crop injury from certain product mixtures. At this application timing, tank-mix this product only with pesticide products which allow tank-mixing and post-emergence chemigation on their product label. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

SOYBEANS*

This product may be applied pre-plant surface, pre-plant incorporated, pre-emergence or as a sequential application in Soybeans to control the weeds listed on this label.

*Not registered for use in California.

GRAZING AND FEEDING TREATED SOYBEAN PLANTS

Treated Soybean plants may be grazed or fed to livestock 40 days after the last application of this product.

RATE RANGES

Where a rate range is shown, use a lower rate on soils that are *Coarse textured* and/or low in organic matter. Use a higher rate on soils that are relatively *Fine textured* and/or high in organic matter.

REPLANTING

If replanting is necessary in fields previously treated with this product, the field may be replanted to Soybeans. A minimum of tillage is directed. Do not apply a second treatment as injury to Soybeans may occur.

USE PRECAUTIONS

Injury to Soybeans or reduced weed control may occur when this product is used under the following conditions. Avoid these conditions wherever possible.

- When soils have a calcareous surface area or a pH of 7.5 or higher.
- Due to the sensitivity of certain Soybean varieties, this product is not for use on Altona, AP 55, AP 71, Asgrow 6520, Burlison, Coker 102, Coker 156, Dassel, GL 3202, Govan, Maple Amber, NB 3665, NKS 1884, Paloma 350, Portage, Regal, Semmes, Terra-Vig 505, Terra-Vig 606, Tracy, Vansoy and Vinton 81. If you choose to plant a newly released Soybean variety, consult your seed supplier for information on its tolerance to Metribuzin (an a.i. in this product) before using this product.
- When applied in conjunction with soil-applied organic phosphate pesticides.
- Uneven application or improper incorporation of this product can decrease the level of weed control and/or increase the level of crop injury.
- When applied to any soil with less than 0.5% organic matter.
- Where soil incorporation is deeper than recommended.
- When sprayers were not calibrated accurately.
- When heavy rain occurs soon after application, especially in poorly drained areas where water may stand for several days.
- When Soybeans are planted less than 1.5 inches deep, particularly when this product is applied pre-emergence.
- Where high soil levels of Atrazine are present.
- When using poor quality Soybean seed.

WEEDS CONTROLLED BY THIS PRODUCT

When applied as directed, this product will control or provide suppression of the following weeds.

Annual Broadleaves*		
Anoda, Spurred Beggarweed, Florida Carpetweed Chickweed, Common Cocklebur** Copperleaf, Hophornbeam <i>Galinsoga</i> spp. Henbit Jimsonweed Knotweed, spp. Kochia Ladysthumb	Lambsquarters, Common Lettuce, Prickly Mallow, Venice Mustard, spp. Nightshade, Black Nightshade, Hairy** Pennycress, Field Pepperweed, Virginia Pigweed, spp. Purslane, Common Pusley, Florida Ragweed, Common**	Redweed <i>Sesbania</i> spp. Shepherdspurse Sicklegod Sida, Prickly/ Teaweed Smartweed, Pennsylvania Spurge, Spotted Starbur, Bristly Sunflower, Common** Thistle, Russian Velvetleaf** Waterhemp, spp.
(Continued)		

WEEDS CONTROLLED BY THIS PRODUCT (Cont.)

Annual Grasses		
Barnyardgrass	Foxtail, spp.	Red rice
Bluegrass, Annual	Goosegrass	Sandbur, spp.**
Crabgrass, spp.	Johnsongrass,	Shattercane**
Crowfootgrass	Seedling**	Signalgrass,
Cupgrass, Prairie	Junglerice	Broadleaf
Cupgrass,	Panicum, Fall	Witchgrass
Southwestern	Panicum, Texas**	
Sedge		
Yellow nutsedge**	-	-
Volunteer Crops		
Barley**	Sorghum**	Wheat**
*Except Triazine resistant biotypes other than Black nightshade, <i>Galinsoga</i> spp., Pigweed, spp. and Waterhemp, spp.		
**Suppression only. Suppression means significant activity, but not always at a level considered acceptable for commercial weed control.		

THIS PRODUCT IN FOUNDATION PROGRAM FOR PLANNED TWO-PASS WEED CONTROL SYSTEMS

This product may be applied pre-plant incorporated or pre-emergence at 1.5 to 1.8 pints per acre on all soils to reduce competition from the weeds listed in the above "WEEDS CONTROLLED BY THIS PRODUCT" table for a 30 day period when followed by a planned post-emergence weed control treatment. Post-emergence treatments include any product or combination of products labeled to control the specific weeds remaining in the field including Glyphosate, i.e., Imitator®, Roundup® or Touchdown® (for use only on Roundup Ready® or Glyphosate tolerant Soybean varieties). Follow all application directions for this product used alone, either pre-plant incorporated or pre-emergence. For the post-emergence herbicide application, consult the selected post-emergence herbicide manufacturer's label for weeds controlled, weed size, application rate, additional use directions, precautions and limitations before use.

Use Restriction: On soils with pH above 7.0, use only the 1.5 pints per acre rate of this product.

THIS PRODUCT IN CONVENTIONAL TILLAGE SYSTEMS**Pre-plant Incorporated Application:**

Incorporate this product uniformly into the top 2 inches of soil within 14 days before planting using a disk, field cultivator, rolling cultivator or similar implement. Apply this product pre-plant incorporated if furrow irrigation is used or when a period of dry weather after application is expected. If Soybeans are planted on beds, apply and incorporate the tank-mixture after bed formation.

Pre-emergence Application:

Dry weather following pre-emergence application of this product may reduce effectiveness. If weeds develop, cultivate uniformly with shallow tilling equipment such as a rotary hoe that will not damage Soybeans.

For information on applying this product in fluid or dry fertilizer, refer to "APPLICATION IN WATER OR FLUID FERTILIZERS" or "IMPREGNATION ONTO DRY BULK FERTILIZERS" and "APPLICATION OF IMPREGNATED DRY BULK GRANULAR FERTILIZER" on this label.

Application Rate:

This Product in Conventional Tillage Systems – Broadcast Rate (Pt./Ac.)		
Soil Texture	0.5 to 3% Organic Matter	> 3% Organic Matter**
COARSE* Loamy sand, Sandy loam	1.2 to 1.5***	1.5 to 1.8
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	1.8 to 2.1	2.1 to 2.4
FINE Silty clay, Silty clay loam****, Clay, Clay loam	2.4 to 2.7	2.4 to 3.0
*Do not use on Sandy soils. On Coarse textured soils, do not use on Loamy sand soils with less than 2% organic matter.		
**For pre-plant incorporated application, use the lower specified rate.		
***For southern and southeastern states, see "THIS PRODUCT IN COARSE (LIGHT) SOILS" section below.		
****Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.		

Use Precautions: On soils with pH above 7.0, Soybean injury caused by Metribuzin in this product may occur at rates higher than 1.5 pints per acre. To avoid injury, do not use this product at rates greater than 1.5 pints per acre on soils above pH 7.0.

THIS PRODUCT IN COARSE (LIGHT) SOILS

(Only in AL, AR, FL, GA, LA, MS, MO, NC, OK, SC, TN, TX, VA)

This product may be applied as a pre-plant incorporated or pre-emergence application in *Coarse textured*, low organic matter soils in the states listed above. Refer to the appropriate sections of this label for specific use directions, precautions and restrictions.

Weeds Controlled:

Refer to the "WEEDS CONTROLLED BY THIS PRODUCT" table under the "SOYBEANS" section.

Application Rate:

Pre-emergence Application of This Product – Broadcast Rate (Pt./Ac.)		
Soil Texture	Organic Matter	This Product (Pt./Ac.)*
COARSE Sand**, Loamy sand, Sandy loam	0.5% or above	1.2 to 2.1
*Use the higher specified rate under heavy weed pressures and/or on soils higher in organic matter. For maximum control of Sicklepod, use a pre-emergence application.		
**Not for use on Sand with less than 1% organic matter.		

Use Precautions: On soils with pH above 7.0, Soybean injury caused by Metribuzin in this product may occur at rates higher than 1.5 pints per acre. To avoid injury, do not use this product at rates greater than 1.5 pints per acre on soils above pH 7.0.

THIS PRODUCT IN TANK-MIXTURE FOR USE IN SOYBEANS**This Product Plus Flumetsulam (e.g., Python®) Tank-Mix Application**

This product may be applied with Flumetsulam (e.g., Python) herbicide as a pre-plant surface, pre-plant incorporated or pre-emergence for the control of certain broadleaf weeds and grasses in Soybeans. Consult the Flumetsulam (e.g., Python) label for specific use directions, precautions and restrictions not specified on this label. Where differences arise, the more restrictive language must be followed.

Weeds Controlled:

In addition to weeds controlled by this product alone, this product plus Flumetsulam (e.g., Python) will improve control of Common ragweed, Kochia, Palmer amaranth, Velvetleaf, Waterhemp, spp., Wild sunflower and Triazine resistant Common lambsquarters. (**Note:** Flumetsulam (e.g., Python) will not improve control of ALS resistant weeds.)

Application Rate:

This Product + Flumetsulam (e.g., Python) – Broadcast Rate*		
Soil Texture	This Product (Pt./Ac.)	Python 80 WDG (Oz./Ac.)
COARSE Loamy sand or Sandy loam	1.2 to 1.5**	Labeled rates
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	1.5 to 2.1	Labeled rates
FINE Silty clay, Silty clay loam***, Clay, Clay loam	2.1 to 2.7	Labeled rates
*Use the higher specified rate on soils with more than 3% organic matter.		
**For southern and southeastern states in Coarse soils, see "THIS PRODUCT IN COARSE (LIGHT) SOILS" section for rates of this product.		
***Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.		

Use Precautions: On soils with pH above 7.0, Soybean injury caused by Metribuzin in this product occasionally occurs at rates higher than 1.5 pints per acre. To avoid injury, do not use this product at rates greater than 1.5 pints per acre on soils above pH 7.0.

This product Plus Imazaquin (e.g., Scepter®) Tank-Mix Application

This product may be applied with Imazaquin (e.g., Scepter) herbicide as a pre-plant surface, pre-plant incorporated or pre-emergence for the control of certain broadleaf weeds and grasses in Soybeans. Consult the Imazaquin (e.g., Scepter) label for specific use directions, precautions, restrictions and any additional weeds not specified on this label. Where differences arise, the more restrictive language must be followed.

Weeds Controlled:

In addition to weeds controlled by this product alone, this product plus Imazaquin (e.g., Scepter) improves control or suppress (reduce the competition) the following annual broadleaf weeds:

Buffalobur Cocklebur Ivyleaf* Morningglory, Pitted	Morningglory, Smallflower Morningglory, Tall* Ragweed, Common	Ragweed, Giant* Sicklepod Sunflower
---	--	---

*Suppression only (reduced competition).

Application Rate:

This Product + Imazaquin (e.g., Scepter) – Broadcast Rate		
Soil Texture	This Product* (Pt./Ac.)	Scepter 70 DG** (Oz./Ac.)
COARSE Loamy sand or Sandy loam	1.2 to 1.5***	Labeled rates
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	1.5 to 2.1	Labeled rates
FINE Silty clay, Silty clay loam****, Clay, Clay loam	2.1 to 2.7	Labeled rates

*Use the higher specified rate on soils with more than 3% organic matter.
 **For pre-emergence application, use the higher specified rate. For maximum control of moderate to heavy infestations of Cocklebur, Giant ragweed and Sicklepod, use the higher specified rate and a pre-plant incorporated application.
 ***For southern and southeastern states in Coarse soils, see "THIS PRODUCT IN COARSE (LIGHT) SOILS" section for rates of this product.
 ****Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.

Use Precautions: On soils with pH above 7.0, Soybean injury caused by Metribuzin in this product occasionally occurs at rates higher than 1.5 pints per acre. To avoid injury, do not use this product at rates greater than 1.5 pints per acre on soils above pH 7.0.

This product Plus Metribuzin + Chlorimuron (e.g., Canopy®) Tank-Mix Application

This product may be applied with Metribuzin + Chlorimuron (e.g., Canopy) herbicide as a pre-plant surface, pre-plant incorporated or pre-emergence for the control of certain broadleaf weeds and grasses in Soybeans. Consult the Metribuzin + Chlorimuron (e.g., Canopy) label for specific use directions, precautions, restrictions and any additional weeds not specified on this label. Where differences arise, the more restrictive language must be followed.

Weeds Controlled:

In addition to weeds controlled by this product alone, this product plus Metribuzin + Chlorimuron (e.g., Canopy) will improve control of Cocklebur and Velvetleaf and provide additional suppression (reduce competition) of Common ragweed, Giant ragweed and Morningglory, spp.

Application Rate:

This Product + Chlorimuron (e.g., Canopy) – Broadcast Rate		
Soil Texture*	This Product** (Pt./Ac.)	Canopy 75 DG*** (Oz./Ac.)
COARSE Loamy sand or Sandy loam	1.2 to 1.5****	N/A
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	1.5 to 2.1	Labeled rates
FINE Silty clay, Silty clay loam*****, Clay, Clay loam	2.1 to 2.7	Labeled rates

*Do not use on soils with pH greater than 7.0.
 **Use the higher specified rate on soils with more than 3% organic matter.
 ***Do not use Chlorimuron (e.g., Canopy 75 DG) as a tank-mix partner on soils with pH greater than 6.8.
 ****For southern and southeastern states in Coarse soils, see "THIS PRODUCT IN COARSE (LIGHT) SOILS" section for rates of this product.
 *****Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.

Use Precautions: On soils with pH above 7.0, Soybean injury caused by Metribuzin in this product occasionally occurs at rates higher than 1.5 pints per acre. To avoid injury, do not use this product at rates greater than 1.5 pints per acre on soils above pH 7.0.

This product Plus Cloransulam (e.g., FirstRate®) Tank-Mix Application

This product may be applied with Cloransulam (e.g., FirstRate) herbicide as a pre-plant, pre-plant incorporated or pre-emergence for the control of certain broadleaf weeds and grasses in Soybeans. Consult the Cloransulam (e.g., FirstRate) label for specific use directions, precautions, restrictions and any additional weeds not specified on this label. Where differences arise, the more restrictive language must be followed.

Weeds Controlled:

In addition to weeds controlled by this product alone, this product plus Cloransulam (e.g., FirstRate) will improve control of Cocklebur, Common ragweed, Common sunflower, Giant ragweed and Velvetleaf and provide additional suppression (reduce competition) of Morningglory, spp.

Application Rate:

This Product + Cloransulam (e.g., FirstRate) – Broadcast Rate		
Soil Texture	This Product (Pt./Ac.)	FirstRate 84 WDG* (Oz./Ac.)
COARSE Loamy sand or Sandy loam	1.2 to 1.5**	Labeled rates
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	1.5 to 2.1	Labeled rates
FINE Silty clay, Silty clay loam***, Clay, Clay loam	2.1 to 2.7	Labeled rates

*Use the higher specified rate on soils with more than 3% organic matter.
 **For southern and southeastern states in Coarse soils, see "THIS PRODUCT IN COARSE (LIGHT) SOILS" section for rates of this product.
 ***Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.

Use Precautions: On soils with pH above 7.0, Soybean injury caused by Metribuzin in this product occasionally occurs at rates higher than 1.5 pints per acre. To avoid injury, do not use this product at rates greater than 1.5 pints per acre on soils above pH 7.0.

This product Plus Clomazone (e.g., Command®) Tank-Mix Application

This product may be applied with Clomazone (e.g., Command) herbicide as a pre-plant or shallow incorporated broadcast application for the control of certain broadleaf weeds and grasses in Soybeans. Consult the Clomazone (e.g., Command) label for specific use directions, precautions, restrictions and any additional weeds not specified on this label. Where differences arise, the more restrictive language must be followed.

Weeds Controlled:

In addition to weeds controlled by this product alone, this product plus Clomazone (e.g., Command) will improve control of heavy infestations of Common ragweed, Jimsonweed and Velvetleaf.

Application Rate:

This Product + Clomazone (e.g., Command) – Broadcast Rate		
Soil Texture	This Product* (Pt./Ac.)	Command 3ME (Pt./Ac.)
COARSE Loamy sand or Sandy loam	1.2 to 1.5**	Labeled rates
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	1.5 to 2.1	Labeled rates
FINE Silty clay, Silty clay loam***, Clay, Clay loam	2.1 to 2.7	Labeled rates

*Use the higher specified rate on soils with more than 3% organic matter.
 **For southern and southeastern states in Coarse soils, see "THIS PRODUCT IN COARSE (LIGHT) SOILS" section for rates of this product.
 ***Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.

Use Precautions: On soils with pH above 7.0, Soybean injury caused by Metribuzin in this product occasionally occurs at rates higher than 1.5 pints per acre. To avoid injury, do not use this product at rates greater than 1.5 pints per acre on soils above pH 7.0. Do not plant Alfalfa, Barley, Oats, Rye or Wheat in the Fall or following Spring after application as crop injury may occur. Do not apply where weather conditions favor drift.

This product Plus Pendimethalin (e.g., PIN-DEE™ 3.3 EC, Prowl® 3.3 EC) Tank-Mix Application

This product may be applied with Pendimethalin (e.g., PIN-DEE 3.3 EC, Prowl 3.3 EC) herbicide as a pre-plant surface, pre-plant incorporated or pre-emergence broadcast application for the control of certain broadleaf weeds and grasses in Soybeans. Consult the Pendimethalin (e.g., PIN-DEE 3.3 EC, Prowl 3.3 EC) label for specific use directions, precautions, restrictions and any additional weeds not specified on this label. Where differences arise, the more restrictive language must be followed.

Weeds Controlled:

In addition to weeds controlled by this product alone, this product plus Pendimethalin (e.g., PIN-DEE 3.3 EC, Prowl 3.3 EC) will improve control of Triazine resistant weeds such as Common lambsquarters, Pigweed, spp., etc.

Application Rate:

This Product + Pendimethalin (e.g., PIN-DEE 3.3 EC or Prowl 3.3 EC) – Broadcast Rate		
Soil Texture	This Product* (Pt./Ac.)	PIN-DEE 3.3 EC or Prowl 3.3 EC (Pt./Ac.)
COARSE Loamy sand or Sandy loam	1.5 to 1.8**	Labeled rates
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	1.8 to 2.1	Labeled rates
FINE Silty clay, Silty clay loam***, Clay, Clay loam	2.1 to 2.7	Labeled rates
*Use the higher specified rate on soils with more than 3% organic matter. **For southern and southeastern states in Coarse soils, see "THIS PRODUCT IN COARSE (LIGHT) SOILS" section for rates of this product. ***Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.		

Use Precautions: On soils with pH above 7.0, Soybean injury caused by Metribuzin in this product occasionally occurs at rates higher than 1.5 pints per acre. To avoid injury, do not use this product at rates greater than 1.5 pints per acre on soils above pH 7.0.

HERBICIDES THAT MAY BE APPLIED POST-EMERGENCE FOLLOWING APPLICATION OF THIS PRODUCT

If required, application of this product alone or in tank-mixture may be followed by an application of a post-emergence herbicide to provide additional control of certain weeds. The following post-emergence herbicides may be applied:

Acifluorfen (e.g., Ultra Blazer®) Bentazon (e.g., Basagran®) Bentazon + Acifluorfen (e.g., Storm®) Carfentrazone (e.g., Aim®) Chloransulam (e.g., First Rate) Chlorimuron-ethyl (e.g., Classic®) Clethodim (e.g., Arrow™, Select®) Fluazifop (e.g., Fusilade®) Fluazifop + Fenoxaprop (e.g., Fusion®) Flumiclorac (e.g., Resource®) Fomesafen (e.g., Foma™, Flexstar®) Glufosinate ammonium (e.g., Liberty®)**	Glyphosate (e.g., Imitator, Roundup)* Glyphosate + Fomesafen (e.g., FomAsate™, Flexstar GT)* Glyphosate + Imazethapyr (e.g., Extreme®)* Glyphosate + Metolachlor (e.g., Sequence®)* Imazamox (e.g., Raptor®) Imazaquin (e.g., Scepter) Imazethapyr (e.g., Pursuit®) Lactofen (e.g., Cobra®) Quizalofop (e.g., Assure®) Sethoxydim (e.g., Poast®) Thifensulfuron (e.g., Harmony®) Thifensulfuron + Chlorimuron (e.g., Synchrony®)***
*Use on Roundup Ready or Glyphosate tolerant Soybean varieties only. **Use on LibertyLink® Soybean varieties only. ***Use on STS™ Soybean varieties only.	

Refer to the above information and the individual product labels for use directions, use rates and precautions and restrictions.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

Reduced Application Rate of Imazaquin (e.g., Scepter) Following Application of This Product

If required, application of this product alone or in tank-mixture may be followed by an early post-emergence application of a reduced rate of Imazaquin (e.g., Scepter) herbicide for improved control of Cocklebur. Apply specified rates of Scepter 70 DG. Use the lower specified rate of Imazaquin (e.g., Scepter) if Cockleburs are less than 3 inches tall or have fewer than 3 leaves and are actively growing and use the higher specified rate if Cockleburs are 3 to 6 inches tall and actively growing. Do not use Imazaquin (e.g., Scepter) when plants have been subjected to stress conditions. Use of non-ionic surfactant or Crop Oil Concentrate (COC) (e.g., Peptoil®) is recommended for Scepter applications. Refer to the Imazaquin (e.g., Scepter) label for additional use directions, precautions and restrictions. Where differences arise, the more restrictive language must be followed.

BURNDOWN WEED CONTROL

This product can be used as part of a burndown herbicide program for control of existing vegetation prior to Soybean emergence in conservation tillage (reduced-tillage/no-till) systems. This product may be tank-mixed with Clethodim (e.g., Select), 2,4-DB (e.g., 2,4-DB DMA 175 or 200), 2,4-D Low Volatile Ester (LVE) (e.g., De-Ester®), Fluazifop + Fenoxaprop (e.g., Fusion), Glyphosate (e.g., Imitator, Roundup), Paraquat (e.g., Quik-Quat®, Gramoxone®) or Sethoxydim (e.g., Poast) for control of emerged weeds prior to crop emergence. Burndown tank-mixes of this product can be applied before planting or prior to crop emergence.

Weeds Controlled:

This product in tank-mixtures with the herbicides in the above "TANK-MIX PARTNER BURNDOWN RATES" table will provide burndown control of the weeds listed below.

WEEDS CONTROLLED BY BURNDOWN RATES OF THIS PRODUCT									
WEEDS	This product +								
	De-Ester LV4	Poast Plus + De-Ester LV4	Select + De-Ester LV4	Fusion + De-Ester LV4	Imitator, Roundup or Touchdown	Imitator, Roundup or Touchdown + De-Ester LV4	Quik-Quat or Gramoxone	Quik-Quat or Gramoxone + De-Ester LV4	2,4-DB DMA 175
ANNUAL GRASSES MAXIMUM BURNDOWN HEIGHT (INCHES)									
Barley	Does not control these species	-	-	-	8		4 to 6		Does not control these species
Barnyardgrass		2 to 3	3 to 4	-	6		4 to 6		
Crabgrass, spp.		2 to 3	-	-	6		4 to 6		
Foxtail, spp.		2 to 3	3 to 4	2 to 6	8		4 to 6		
Johnsongrass, Seedling		2 to 3	-	-	8		4 to 6		
Panicum, Fall		2 to 3	3	2 to 6	6		4 to 6		
Sandbur, Field		-	-	-	8		4 to 6		
Shattercane		2 to 3	-	-	8		4 to 6		
Wheat, Volunteer		-	-	-	6		4 to 6		
Witchgrass		2 to 3	-	-	6		4 to 6		
BROADLEAVES MAXIMUM BURNDOWN HEIGHT (INCHES)									
Buffalobur		-			6	6	4 to 6	4 to 6	-
Chickweed, Common		6			6	6	4 to 6	4 to 6	2
Cocklebur, Common		6			6	8	4 to 6	4 to 6	6
Dandelion, Common		6 dia ^a			2 dia ^b	6 dia ^a	4 dia ^c	6 dia ^a	2 dia
Henbit		4			4	4	4 to 6	4 to 6	-
Horseweed / Marestail		6 ^a			4 ^b	6	3	6 ^a	2 ^c
Jimsonweed		6			6	6	4 to 6	4 to 6	2
Kochia		4 ^a			4	4	4	4	-
Ladysthumb		6			6	8	4 to 6	4 to 6	3
Lambsquarters, Common		6			6	8	4 to 6	4 to 6	2
Lettuce, Prickly		6			4	6	4 to 6	4 to 6	2
Mallow, Venice		6			6	6	4 to 6	4 to 6	-
Morningglory, spp.		6			2	4	2	4	4
Mustard, spp.		6			6	8	4 to 6	4 to 6	2
Pennycress, Field		6			6	6	4 to 6	4 to 6	2
Pigweed, spp. (Annual)		6			6	8	4 to 6	4 to 6	3
Ragweed, Common		6			6 ^b	8	4 to 6	4 to 6	2
Ragweed, Giant		6 ^a			4 ^b	6	4	6	2
Shepherdspurse		6			6	6	4 to 6	4 to 6	-
Sida, Prickly		6			4	4	4	4	1
Smartweed, Pennsylvania		6			6	8	4 to 6	4 to 6	3
Sunflower, Common		6			6	6	4 to 6	4 to 6	4
Thistle, Russian		4 ^a			2 to 4 ^b	6	4	4 to 6	3 ^c
Velvetleaf		6			6	8	4 to 6	4 to 6	3
Waterhemp, spp.		6			6	8	4 to 6	4 to 6	3
^a Use 2,4-D LVE (e.g., De-Ester LV4) at labeled rates.									
^b Use Imitator, Roundup or Touchdown at a minimum labeled rate.									
^c Suppression only.									

^a Use 2,4-D LVE (e.g., De-Ester LV4) at labeled rates.

^b Use Imitator, Roundup or Touchdown at a minimum labeled rate.

^c Suppression only.

Application:

This product may be applied up to 30 days before planting or pre-emergence. Apply only by ground equipment when this product is used for burndown of existing vegetation in conservation tillage systems. Use the high end of the rate range for applications of this product made 14 to 30 days before planting.

Refer to the following tables for rates of this product and rates of the tank-mix partners. Follow all label directions, precautions and restrictions for tank-mix partners. Where differences arise, the more restrictive language must be followed.

Refer to the "PRODUCT INFORMATION" section of this label for additional information, precautions and limitations.

Soil Texture	This Product (Pt./Ac.)*
COARSE** Loamy sand, Sandy loam	1.2 to 2.1
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	2.1 to 3.0
FINE Silty clay, Silty clay loam***, Clay, Clay loam	2.7 to 3.6

*Use the low rate range for low residue level or soils with less than 3% organic matter. Use the higher rate range for high residue level or soils with greater than 3% organic matter.

**Do not use on Sand soil. On Coarse textured soils, do not use on Loamy sand soils with less than 2% organic matter.

***Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.

TANK-MIX PARTNER BURNDOWN RATES		
This product * +	Rate/Ac.	Use Directions and Remarks
2,4-DB (e.g., 2,4-DB DMA 175 or 2,4-DB DMA 200)	Labeled rates	Apply pre-plant or before Soybean emergence. Include non-ionic surfactant (e.g., Surf-Ac® 820) at 2 qts./100 gals. (0.5% v/v) of spray solution.
2,4-D LVE** (e.g., De-Ester LV4)	Labeled rates	Observe the number of days for pre-planting when using 2,4-D LVE at the specified rates. Include Crop Oil Concentrate (COC) (e.g., Peptoil) at the rate of 1 gal./100 gal of spray solution (1% v/v).
Fusion + 2,4-D LVE** (e.g., De-Ester LV4)	Labeled rates	For this tank-mix, follow the planting restrictions under the above "Use Directions and Remarks" for "This product + 2,4-D LVE". Fusion at specified rates will control certain grasses up to 2, 4 and 6 inches in height. Include either Crop Oil Concentrate (COC) (e.g., Peptoil) at 1 gal./100 gals. (1.0% v/v) or non-ionic surfactant (e.g., Surf-Ac 820) at 1 to 2 qts./100 gals. (0.25 to 0.5% v/v) of spray solution. Refer to the Fusion label for additional information.
Imitator /Roundup or Touchdown	Labeled rates	Must be applied prior to crop emergence. Use the higher rates as weeds approach the maximum weed heights listed in the "Weeds Controlled" section below. Apply in 10 to 20 gals. of water per acre. With Imitator or Roundup, include non-ionic surfactant (e.g., Surf-Ac 820) at 2 qts./100 gals. (0.5% v/v) and ammonium sulfate (spray grade) at 17 lbs./100 gals. of spray solution. Any Glyphosate formulation registered and labeled for use in Soybeans may be tank-mixed with this product .
Imitator/Roundup or Touchdown + 2,4-D LVE** (e.g., De-Ester LV4)	Labeled rates	For this tank-mix, follow the "Use Directions and Remarks" for "This product + 2,4-D LVE" and "This product + Imitator/Roundup or Touchdown", paying special attention to planting restrictions with 2,4-D LVE. Use the adjuvant directions above for "This product + Imitator/Roundup or Touchdown" tank-mix. Do not use Crop Oil Concentrate (COC).
Poast Plus + 2,4-D LVE** (e.g., De-Ester LV4)	Labeled rates	For this tank-mix, follow the planting restrictions under the "Use Directions and Remarks" for "This product + 2,4-D LVE". Poast Plus at specified rates will control certain grasses up to 2 and 3 inches in height. Include either Crop Oil Concentrate (COC) (e.g., Peptoil) at the rate of 1 gal./ 100 gals. of spray solution (1% v/v) or non-ionic surfactant (e.g., Surf-Ac 820) as needed. Refer to the Poast Plus label for additional information.
(Continued)		

TANK-MIX PARTNER BURNDOWN RATES (Cont.)		
This product * +	Rate/Ac.	Use Directions and Remarks
Quik-Quat or Gramoxone	Labeled rates	Must be applied prior to crop emergence. See Quik-Quat or Gramoxone label for amount to use in relation to weed height. Apply in 20 to 60 gals. of water per acre. Include either non-ionic surfactant (e.g., Surf-Ac 820) at 1 qt./100 gals. (0.25% v/v) or Crop Oil Concentrate (COC) (e.g., Peptoil) at 1 gal./100 gals. (1% v/v) of spray solution.
Quik-Quat or Gramoxone + 2,4-D LVE** (e.g., De-Ester LV4)	Labeled rates	For this tank-mix, follow the "Use Directions and Remarks" for "This product + 2,4-D LVE" and "This product + Quik-Quat or Gramoxone" paying special attention to crop planting restrictions with 2,4-D LVE. Include either non-ionic surfactant (e.g., Surf-Ac 820) or Crop Oil Concentrate (COC) (e.g., Peptoil) in this tank-mix.
Select + 2,4-D LVE** (e.g., De-Ester LV4)	Labeled rates	For this tank-mix, follow the planting restrictions under the above "Use Directions and Remarks" for "This product + 2,4-D LVE". Select at specified rates will control certain grasses up to 3 and 4 inches in height. Include Crop Oil Concentrate (COC) (e.g., Peptoil) at the rate of 1 qt./Ac. and 28% Urea Ammonium Nitrate (UAN) at a rate of 1 to 2 qts./Ac. as needed. Refer to the Select label for additional information.
<p>*Refer to the preceding table for rates of this product to be used in the tank-mixture.</p> <p>**Apply only 2,4-D Low Volatile Ester (2,4-D LVE) formulations which are registered for pre-plant or burndown use.</p>		

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

Use Restrictions:

- Do not apply this product in tank-mixture with the above products for burndown weed control after crop emergence.
- Do not apply tank-mixtures containing 2,4-D LVE if wind is blowing toward desired susceptible plants (i.e., Cotton, Tobacco, Tomato, etc.) or when wind speeds exceed 6 miles per hour. Observe all precautions and limitations of all products used in the tank-mixture(s).

Feeding Restrictions:

Soybean plants or hay treated with this product may be grazed or fed to livestock 40 days after application. Follow the most restrictive Pre-Harvest Interval (PHI) of all products used in the tank-mixture.

THIS PRODUCT IN REDUCED AND NO-TILL SYSTEMS

Pre-plant Surface Application:

This product may be used in reduced-till and no-till systems. Applications may be made up to 30 days before planting or after planting, but before Soybean emergence. Residual herbicides such as Chloransulam (e.g., FirstRate), Clomazone (e.g., Command), Flumetsulam (e.g., Python), Imazaquin (e.g., Scepter), Metribuzin + Chlorimuron (e.g., Canopy) and Pendimethalin (e.g., PIN-DEE 3.3 EC, Prowl 3.3 EC) may be tank-mixed for additional weed control. If weeds are present at time of application, burndown herbicides may be added to the tank-mixes (see "BURNDOWN WEED CONTROL" section). Refer to the tank-mix product labels for specific rates and use directions.

Application Rate:

Reduced-Till and No-Till Systems – Broadcast Rate	
Soil Texture	This Product (Pt./Ac.)*
COARSE** Loamy sand, Sandy loam	1.2 to 2.1
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	2.1 to 3.0
FINE Silty clay, Silty clay loam***, Clay, Clay loam	2.7 to 3.6
*Use the low rate range for low residue level or soils with less than 3% organic matter. Use the higher rate range for high residue level or soils with greater than 3% organic matter.	
**Do not use on Sand soil. On Coarse textured soils, do not use on Loamy sand soils with less than 2% organic matter.	
***Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.	

THIS PRODUCT – SEQUENTIAL APPLICATION

An early pre-plant (surface-applied or shallow incorporated) application of this product, followed by a pre-emergence application of this product after planting but before Soybean emergence will provide more consistent control of broadleaf and grass weeds than a single application.

A sequential application will decrease the need for tillage and/or burndown herbicides for the control of existing vegetation before planting, while providing residual control of weeds after planting.

Weeds Controlled:

In addition to weeds controlled by this product alone, the sequential application improves control of the following annual broadleaf weeds.

Buffalobur	Ragweed, Common	Velvetleaf
Cocklebur	Sunflower	

Application:

An early pre-plant application may be made 15 to 30 days before planting Soybeans. Follow this application with a pre-emergence overlay application of this product after planting but before crop emergence. Follow directions on this label for sequential applications from 0 to 14 days before planting.

Where a rate range is directed, use the higher specified rates (a) in fields with a history of severe weed pressure, (b) when the time between early pre-plant and pre-emergence overlay applications approaches the maximum 30 days, (c) when the organic matter content of the soil is over 3%, and/or (d) when heavy crop residues are present on the soil surface.

When weeds exceed 1 to 1.5 inches in height or diameter at application, use a burndown herbicide such as Glyphosate (i.e., Imitator, Roundup or Touchdown), Paraquat (e.g., Quik-Quat, Gramoxone or 2,4-D LVE (e.g., De-Ester LV4).

Sequential Application – Broadcast Rate*			
Soil Texture	Early Pre-Plant Application This Product (Pt./Ac.)	- Followed By -	Pre-Emergence Overlay Application This Product (Pt./Ac.)
COARSE** Sand, Loamy sand or Sandy loam	1.2 to 1.8	- Followed By -	0.3 to 0.9
MEDIUM Loam, Silt loam, Silt, Sandy clay, Sandy clay loam	1.5 to 2.1	- Followed By -	0.6 to 1.2
FINE Silty clay, Silty clay loam***, Clay loam	1.8 to 2.4	- Followed By -	0.9 to 1.5
*Do not exceed a total of 3.9 pts./Ac. of this product per use season.			
**On Coarse textured soils, do not use on Sand soil with less than 1% organic matter. However, on Coarse textured soils with a calcareous surface area or a pH of 7.5 or higher, do not use on Sand soil with less than 2% organic matter or on Loamy sand or Sandy loam soils with less than 1% organic matter.			
***Silty clay loam soils are transitional soils and may be classified as Medium textured soils in some regions of the U.S. When using this product, treat this soil as Fine textured.			

THIS PRODUCT – POST-EMERGENCE DIRECTED APPLICATION

(AR, LA, MO - Bootheel only, MS, TN)

This product can be applied post-emergence directed to Soybeans to provide residual control of weeds that emerge after crop emergence in the states of Arkansas, Louisiana, Mississippi, Missouri (Bootheel only) and Tennessee. A post-emergence directed spray of this product can be applied to Soybeans in addition to a pre-emergence or pre-plant application of this product according to label directions. The total amount of this product applied must not exceed 3.9 pints per acre per year.

Weeds Controlled:

A post-emergence directed application of this product will provide residual pre-emergence weed control of the weeds listed in the table "WEEDS CONTROLLED BY THIS PRODUCT" under the "SOYBEANS" section.

Application:

Apply this product in 10 to 20 gallons of water per acre in a 6 to 8 inch band on each side of the row when Soybeans are at least 8 inches tall. Do not allow the directed spray to contact more than the lower one-fourth to one-third of the Soybean plant. Soybean leaves contacted by the spray will be killed or severely injured. Do not apply directly to Soybeans or serious injury will occur.

Post-Emergence Directed Application of THIS PRODUCT – Broadcast Rate (Pts./Ac.)		
Soil Texture	0.5% to < 3% Organic Matter	≥ 3% Organic Matter
COARSE Loamy sand, Sandy loam (> 2% organic matter)	1.3	1.5
MEDIUM	1.5 to 2	2
FINE	2	2
Mississippi Delta Only Silty clay, Clay	2	2

Use Precaution: If heavy rain occurs soon after application, crop injury may result especially in poorly drained areas where water stands for several days.

Post-Directed Application on Glyphosate Tolerant Soybeans Only

Post-emergence directed applications of this product can be made in tank-mixture with Glyphosate such as Imitator, Roundup or Touchdown in Glyphosate tolerant Soybeans only.

Refer to the tank-mix partner label for use directions, precautions and restrictions. The most restrictive product labeling applies.

Use Restrictions (THIS PRODUCT - Post-emergence Directed Application)

- Do not exceed a total of 3.9 pints of this product per acre per year.
- Do not graze or feed treated Soybean forage, hay or straw to livestock.
- Pre-Harvest Interval (PHI): Do not apply within 90 days of Soybean harvest.
- This product cannot be applied to *Sandy loam* or *Loamy sand* soils with less than 2% organic matter.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place. Store in original container. Keep container tightly closed.

PESTICIDE DISPOSAL: To avoid waste, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often, such programs are run by state or local governments or by industry).

CONTAINER HANDLING:

Nonrefillable Container (rigid material; ≤ 5 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid material; > 5 gallons up to < 250 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable Container (≥ 250 gallons & Bulk): Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically directed and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable laws, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable laws, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

ME-TOO-LACHLOR, DE-ESTER, FOMA, FOMASATE, IMITATOR, MIX, PEPTOIL, PIN-DEE, QUIK-QUAT, SURF-AC and the DREXEL logo are either trademarks or registered trademarks of Drexel Chemical Company. All other brand names, product names or trademarks belong to their respective holders.