

Agri Star®

SPECIMEN LABEL

METALAXYL

GROUP

4

FUNGICIDE

METALAXYL 265 ST

Seed Treatment Fungicide

ACTIVE INGREDIENTS:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester . . . 28.35%

OTHER INGREDIENTS: 71.65%

TOTAL: 100.00%

Contains 2.65 lbs. of metalaxyl per gallon.

EPA Reg. No. 42750-208

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

In case of an emergency call CHEMTREC toll free at 1-800-424-9300. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

See inside booklet for additional PRECAUTIONARY STATEMENTS.

A SEED TREATMENT CHEMICAL FOR CONTROL OF SEED ROT AND DAMPING-OFF DISEASES OF CERTAIN CROPS

Manufactured For:

ALBAUGH, LLC

1525 NE 36th Street
Ankeny, Iowa 50021

**FOR CHEMICAL SPILL, LEAK,
FIRE, OR EXPOSURE, CALL
CHEMTREC (800) 424-9300**



Albaugh®
Your Alternative™

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING – AVISO

Causes skin irritation. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants,
- Socks and chemical-resistant footwear and
- Chemical-resistant gloves (made of nitrile \geq 14 mils in thickness, neoprene \geq 14 mils in thickness, Viton \geq 14 mils in thickness and/or barrier laminate).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate bodies of water when disposing of planting equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: if the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls,
- socks and chemical resistant footwear,
- Chemical-resistant gloves (made of nitrile \geq 14 mils in thickness, neoprene \geq 14 mils in thickness, Viton \geq 14 mils in thickness and/or barrier laminate).
- protective eyewear.

RESTRICTION:

- In California – Only for use in commercial seed treating establishments. Not allowed for “on-farm” use.
- When applying both metalaxyl-containing and mefenoxam-containing products to the same crop within the same season, do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.
- When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

Seed Bag Label Requirements

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with METALAXYL 265 ST (or other product), a fungicide containing metalaxyl.
- Do not use treated seed for feed, food, or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with metalaxyl:

- Store treated seed away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and chemical-resistant gloves (made of nitrile \geq 14 mils in thickness, neoprene \geq 14 mils in thickness, Viton \geq 14 mils in thickness and/or barrier laminate) when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife.
- Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.
- Excess treated seed may be used for ethanol production if:
 1. by-products are not used for livestock feed and
 2. no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

METALAXYL 265 ST is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora spp.* For control of other soilborne diseases, combination of Gustafson Captan and VITA VAX® registered formulations are compatible.

Do not use with other seed treatment products unless previous experience assures compatibility.

METALAXYL 265 ST may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment. Follow more restrictive labeling requirements of tank mix products. Do not tank mix with any product that contains a prohibition on tank mixing.

RESISTANCE MANAGEMENT RECOMMENDATIONS

METALAXYL 265 ST is a systemic fungicide that contains:

Metalaxyl, a Group 4 mode of action fungicide that belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis.

Any fungal population may contain individuals naturally resistant to Group 4 mode of action fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Because resistance development cannot be predicted, the use of this product should follow appropriate resistance management strategies.

To delay fungicide resistance, use should be based on an IPM program that includes the following steps:

- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance- management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Albaugh, LLC at 1-800-247-8013.

You can also contact your pesticide distributor or university extension specialist to report resistance.

Rotate the use of METALAXYL 265 ST or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.

- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

NOTE: METALAXYL 265 ST is a systemic fungicide having a specific mode of action.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

CROP	APPLICATION INSTRUCTIONS (METALAXYL 265 ST contains 0.015 lbs. of active per fluid ounce)				
Canola* Mustard Seed* Rapeseed*	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.25–0.5 fl. oz. per 100 lbs of seed. <u>*Not registered for use in California.</u>				
Cotton	For <i>Pythium</i> seed rot and damping-off. Apply METALAXYL 265 ST as a seed treatment:				
	METALAXYL 265 ST Application rate			Maximum Planting rate – seeds/A*	
	seeds/lb	fl oz/CWT	mg AI/seed	@ 0.75 fl oz/CWT	@ 1.5 fl oz/CWT
	3600	0.75 - 1.5	0.019 - 0.038	96,320	48,160
	4400	0.75 - 1.5	0.015 - 0.031	117,700	58,860
	5200	0.75 - 1.5	0.013 - 0.026	139,100	69,560
	6000	0.75 - 1.5	0.011 - 0.023	160,500	80,270
*Do not apply by seed treatment more than 0.004 pounds of Metalaxyl (0.2 fl. oz. METALAXYL 265 ST) per acre per year.					

(continued)

CROP	APPLICATION INSTRUCTIONS (METALAXYL 265 ST contains 0.015 lbs. of active per fluid ounce)					
Soybeans	For <i>Pythium</i> damping off and early season <i>Phytophthora</i> control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75–1.5 fl. oz. per 100 lbs. of seed.					
		METALAXYL 265 ST Application rate			Maximum Planting rate – seeds/A*	
	seeds/lb	fl oz/CWT	Fl. Oz./140,000 seed unit	mg AI/seed	@ 0.75 fl oz/CWT	@ 1.5 fl oz/CWT
	2000	0.75 - 1.5	0.53 - 1.05	0.034 - 0.068	222,100	111,000
	2400	0.75 - 1.5	0.44 - 0.88	0.028 - 0.057	266,500	133,200
	2800	0.75 - 1.5	0.38 - 0.75	0.024 - 0.048	310,900	155,400
	3200	0.75 - 1.5	0.33 - 0.66	0.021 - 0.042	355,300	177,700
	3600	0.75 - 1.5	0.29 - 0.58	0.019 - 0.038	399,700	199,900
	*Do not apply by seed treatment more than 0.0166 pounds of Metalaxyl (0.83 fl. oz. METALAXYL 265 ST) per acre per year.					
Peanuts*	For <i>Pythium</i> seed rot and damping off: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75–1.5 fl. oz. per 100 lbs. of seed. *Not registered for use in California.					
Sorghum	For control of <i>Pythium</i> damping-off apply 0.375–0.75 fl. oz. of METALAXYL 265 ST per 100 lbs. of seed applied as a seed treatment. For control of systemic downy mildew (races susceptible to chemical control), apply 0.75–1.5 fl. oz. METALAXYL 265 ST per 100 lbs. of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.5–3.0 fl. oz. of METALAXYL 265 ST per 100 lbs. of seed as a seed treatment. Only high quality sorghum seed (90% germination and above) should be treated with METALAXYL 265 ST.					
Sunflowers	For control of systemic downy mildew: Apply METALAXYL 265 ST as a seed treatment at the rate of 1.5–3.0 fl. oz. per 100 lbs. of seed.					
Forage Legumes	Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage). For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75–1.5 fl. oz. per 100 lbs. of seed.					
Grain Crops	Wheat, barley, rye, oats, millet, milo, corn (field corn, sweet corn and popcorn), buckwheat and triticale. For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For systemic downy mildew control of sweet corn: Apply METALAXYL 265 ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.					
Rice*	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75–1.5 fl. oz. per 100 lbs. of seed. If using metalaxyl treated rice do not use Ridomil for control of <i>Pythium</i> in water-seeded rice. *Not registered for use in California.					
Seed and Pod Vegetables	Black-eyed peas, chickpeas (Garbanzos), cowpeas, dill, edible soybeans, field beans, field peas, garden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, snap beans, string beans, wax beans, and lentils. For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply METALAXYL 265 ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.					
Turf Grasses	For <i>Pythium</i> damping-off control on golf courses, home lawns reclamation and erosion control projects: Apply METALAXYL 265 ST as a seed treated at the rate of 1.5 fl. oz. per 100 lbs of seed.					

(continued)

CROP	APPLICATION INSTRUCTIONS (METALAXYL 265 ST contains 0.015 lbs. of active per fluid ounce)
Forage Grasses	Grasses grown for hay, grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage. For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.
Brassica (Cole) Leafy Vegetables*	Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa and mustard), cauliflower, cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. *Not registered for this use in California.
Cucurbit Vegetables*	Chayote (fruit), Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd (includes hyotan, cucuza, Chinese okra, and hechima), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon (includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash (includes butternut squash, calabaza, hubbard squash), Cucumis mixta, Cucumis pepo (includes acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties of Citrullus lanatus). For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed. *Not registered for this use in California.
Leafy Vegetables*	Amaranth (leafy, Chinese spinach, tampala), cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (finocchio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach* (including New Zealand and vine), and Swiss chard. *Not registered for use in California. NOTE: Do not apply METALAXYL 265 ST to spinach which is destined to be grown in the greenhouse. Containers or bags containing METALAXYL 265 ST treated spinach seed must be labeled with the restriction: "Do not grow spinach from treated seed in greenhouses." For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.
Fruiting Vegetables	Eggplants, ground cherry, pepino, pepper (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs of seed.
Onions	Onion (dry bulb and green) For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs of seed.
Root and Tuber Vegetables (except potatoes)*	Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet (garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley (turnip-rooted), parsnip, radish (includes oriental dalkon), rutabaga, salsify (includes black and Spanish), skirret, sweet potato, tanier, turmeric, turnip, yam bean (jicama, manico pea), yam For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz./100 lbs. of seed. NOTE: Do not apply METALAXYL 265 ST to potatoes. *Not registered for this use in California.

Directions for Use at Reduced Rates in Combination With Other Fungicides:

This product is a systemic fungicide seed dressing which may be applied in combinations with other EPA registered fungicides to aid in control of *Pythium*. This product may be applied as a water-based slurry through standard slurry or mist type commercial seed treatment equipment.

SEED	APPLICATION INSTRUCTIONS (METALAXYL 265 ST contains 0.015 lbs. of active per fluid ounce)
Corn (field) Wheat Oats Barley Rye	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 265 ST as a seed treatment at the rate of 0.10–0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of seed treatment fungicides.
Soybeans Field beans Kidney beans Navy beans Pinto beans Lentils Peas	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 265 ST as a seed treatment at the rate of 0.10–0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Sunflowers	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 265 ST as a seed treatment at the rate of 0.10–0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Peanuts*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 265 ST as a seed treatment at the rate of 0.10–0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides. *Not registered for use in California.
Rice*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 265 ST as a seed treatment at the rate of 0.10–0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides. *Not registered for use in California.

SEED TREATMENT – EXPORT USE

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation.

Seed treated in accordance with the export use application instructions must be labeled as follows:

“For export only – NOT for domestic sale or use.”

Refer to the Seed Bag Label Requirements section for labeling requirements on treated seed containers.

Directions for use at reduced rates in combination with other fungicides METALAXYL 265 ST is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. METALAXYL 265 ST may be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

CROP	APPLICATION INSTRUCTIONS (METALAXYL 265 ST contains 0.015 lbs. of active per fluid ounce)
Brassica (Cole) Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75–1.50 fl. oz. per 100 lbs. of seed.
Canola	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.50–1.70 fl. oz. per 100 lbs. of seed. (32–100 ml per 100 kg of seed)

(continued)

CROP	APPLICATION INSTRUCTIONS (METALAXYL 265 ST contains 0.015 lbs. of active per fluid ounce)					
Corn	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75–1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply METALAXYL 265 ST as a seed treatment at 3.0 fl. oz. per 100 lbs. of seed.					
		METALAXYL 265 ST Application rate			Maximum Planting rate – seeds/A*	
	seeds/lb	fl oz/CWT	Fl. Oz./80,000 seed unit	mg AI/seed	@ 0.1 fl oz/CWT	@ 0.375 fl oz/CWT
	1200	0.1 - 0.375	0.07 - 0.25	0.008 - 0.028	457,500	122,000
	1600	0.1 - 0.375	0.05 - 0.19	0.006 - 0.021	610,000	162,700
	2000	0.1 - 0.375	0.04 - 0.15	0.005 - 0.017	762,500	203,300
	2400	0.1 - 0.375	0.03 - 0.13	0.004 - 0.014	915,000	244,000
	*Do not apply by seed treatment more than 0.0076 pounds of Metalaxyl (0.46. fl. oz. METALAXYL 265 ST) per acre per year.					
Cucurbit Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.					
Fruiting Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75–1.50 fl. oz. per 100 lbs. of seed.					
Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75–1.50 fl. oz. per 100 lbs. of seed.					
Onions	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75–1.50 fl. oz. per 100 lbs. of seed.					
Millet	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75–1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply METALAXYL 265 ST as a seed treatment at 2.5 fl. oz. per 100 lbs. of seed.					
Peas	For <i>Pythium</i> damping-off and early season Phytophthora control: Apply METALAXYL 265 ST as a seed treatment at 3.5 fl. oz. per 100 lbs. of seed.					
Root and Tuber Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75–1.50 fl. oz. per 100 lbs. of seed. Do not apply METALAXYL 265 ST to potatoes.					
Sorghum	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75–1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply METALAXYL 265 ST as a seed treatment at 5.25 fl. oz. per 100 lbs. of seed.					
Sunflowers	For systemic downy mildew control: Apply METALAXYL 265 ST as a seed treatment at 9.0 fl. oz. per 100 lbs. of seed.					

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool dry location. Do not store in direct hot sunlight. Do not store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable > 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container:

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Albaugh, no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Albaugh warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. TO THE EXTENT PERMITTED BY APPLICABLE LAW, ALBAUGH SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, ALBAUGH NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. To the extent permitted by applicable law, Buyer assumes the responsibility to handle, use, and store this product in accordance with safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Albaugh. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.