



# Metalosate®

# Boron

Amino Acid Complex Liquid Foliar



**Net Weight: 10.3 lbs./4.6 kg (1 Gallon)**



07322-AL-GL0001

**Lot #:**



12000000

v02/2012

# METALOSATE® BORON

## GUARANTEED ANALYSIS (w/w):

Boron (B)..... 5.0%  
5.0% Complexed Boron

Derived from Sodium Tetraborate Amino Acid Complex.

## GENERAL INFORMATION:

**Metalosate® Boron** is designed for foliar application on plants to prevent or correct nutrient deficiencies that may limit crop growth and yields. It is water soluble and nontoxic to plants when applied as directed. For best results, apply **Metalosate® Boron** according to recommendations based on plant tissue or soil analysis.

## DIRECTIONS:

**Shake well before use. Keep from freezing.**

Dilute one part **Metalosate® Boron** with 20 or more parts water. Apply by a spray method and in an adequate amount of water that will provide complete coverage of the plants. Use of a non-ionic wetting agent may improve spray coverage of certain hard to wet plants. Do not apply undiluted.

**Metalosate® Boron** can be included in a regular spray program on crops. Consult with an **Albion®** representative on compatibility with other spray materials.

The rate of application will depend on the crop, stage of growth, and severity of deficiency. The maximum recommended rates are for mature, full-sized plants. Reduce the rates proportionately when spraying smaller plants.

**Metalosate® Boron** may be used on legumes, grain crops, root crops, cucurbits, cole crops, leafy vegetables, woody and herbaceous ornamentals, deciduous fruits, vine crops, tropical and subtropical fruits, and many other crops.

Albion, the Albion logo, and Metalosate are trademarks of Albion Laboratories, Inc.

**Manufactured by: Albion Laboratories, Inc.**  
101 North Main Street  
Clearfield, Utah 84015 U.S.A.

## RECOMMENDATIONS:

**Field Crops and Vegetables:** Apply 8 to 16 fluid ounces per acre (0.5 to 1.0 liter per hectare) during periods of rapid growth or nutritional stress. The application may be repeated 2 or more times through the growing season.

**Tree Crops:** Make an application of 12 to 24 fluid ounces per acre (0.75 to 1.5 liter per hectare) after the beginning of active growth. The application may be repeated at 2 to 4 week intervals through the growing season.

**Grapes and Berries:** Make an application of 8 to 16 fluid ounces per acre (0.5 to 1.0 liter per hectare) after active growth begins. The application may be repeated at intervals of one week or more through the vegetative-growth period.

**Turf:** Apply 0.5 to 1.0 fluid ounces per 1000 square feet (10 to 20 milliliters per 100 square meters) during periods of active growth.

**Ornamental Trees, Shrubs and Flowering Plants:** Dilute at a rate of 24 fluid ounces in 10 or more gallons (100 milliliters in 10 or more liters) of water and spray to complete coverage.

## LIMITED WARRANTY:

Seller's and Manufacturer's obligation is limited to replacement of defective product. **NEITHER SELLER NOR MANUFACTURER SHALL BE LIABLE FOR ANY INJURY, LOSS, OR DAMAGE DIRECTLY OR CONSEQUENTLY ARISING OUT OF THE MISUSE OR INABILITY TO USE THE PRODUCT. ALL OTHER WARRANTIES EXCEPT IN WRITING FROM THE MANUFACTURER, WHETHER EXPRESSED OR IMPLIED ARE HEREBY DISCLAIMED.**

General information regarding the contents and levels of metals in this product is available by calling 1 866 CHELATE (1 866 243 5283).

**CAUTION:** This material contains **boron**. Its misuse may cause serious injury to the crops. It should be used only under the direction of fertilizer consultants.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.