Micro-Plus Zinc



SPECIMEN

F2399

Z-MPZN Rev: 21C10

Density: 10.5 lb/gal (1.26 kg/L) @ 68°F (20°C)

NET CONTENT	NET WEIGHT
□ 2.50 gal (9.5 L)	26.3 lb (11.9 kg)
□ 2.64 gal (10.0 L)	27.7 lb (12.6 kg)
□ 5.00 gal (18.9 L)	52.5 lb (23.8 kg)
□ 5.28 gal (20.0 L)	55.4 lb (25.2 kg)
☐ 55.00 gal (208.2 L)	577.5 lb (262.3 kg)
□ 275.00 gal (1041.0 L)	2887.5 lb (1311.6 kg)
□ BULK	

CAUTION

KEEP OUT OF REACH OF CHILDREN; MAY BE HARMFUL IF SWALLOWED, INHALED OR IN CONTACT WITH SKIN; MAY CAUSE EYE IRRITATION. MAY CAUSE DAMAGE THROUGH PROLONGED EXPOSURE.

PRECAUTIONS:

Do not breathe dust/fume/gas/mist/vapors/spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in well-ventilated area. Avoid release to environment. Wear protective gloves/protective clothing/eye protection/face protection. Keep container tightly closed. Keep cool and protected from sunlight.

If medical advice is needed, have product container or label at hand. For more details regarding the safe use/handling, storage, and disposal of this product as well as additional hazards and first aid measures, please obtain the current Safety Data Sheet at www.stollerusa.com.

STORAGE AND DISPOSAL

MICRO-PLUS products are complexed with medium corrosivity. It is recommended that MICRO-PLUS ZINC be stored in fiberglass, polyethylene or polyolefin. We recommend storing pails a maximum of three pails high. Do not stack pallets. Store container tightly closed in a cool/well-ventilated dry place protected from sunlight above 40°F. Do not allow container to freeze. Store locked up. DISPOSE of contents/container at a permitted facility or as advised by your local regulatory authority. Do not contaminate water, food or feed by storage or disposal. Do not apply directly to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash water. Do not reuse empty container.

Information regarding the contents and levels of metals in this product is available on the internet at

http://www.aapfco.org/metals.html

Always read and follow label instructions.

GENERAL INFORMATION

MICRO-PLUS ZINC is for soil and foliar application. It is readily available and rapidly absorbed by the root system or foliage. MICRO-PLUS ZINC is formulated to be used with Stoller liquid fertilizers (Nitro-Plus, Ultra-Cal, Foli-Zyme) and nitrogen solutions. When used with polyphosphate containing fertilizers, dilute with nitrogen solution before addition. Broadcast or row applications are acceptable, although greater efficiency may be expected from row applications. Micronutrient deficiencies usually occur when plants are small; therefore, micronutrients should be added to the fertilizer at or before planting.

DIRECTIONS FOR USE

It may be a violation of law to use this product inconsistent with its labeling.

RECOMMENDED USES

FOR FERTILIZER IMPREGNATION:

Soil application of **MICRO-PLUS ZINC** can be incorporated in liquid fertilizer or applied separately. **MICRO-PLUS ZINC** can be coated on dry blend fertilizer to control dust in the blend plant and supply needed micronutrients. For best results, follow soil test and plant analysis recommendations. In absence of a soil test use the following as a guide:

FOR SOIL APPLICATION

- Maintenance: 1-2 quarts per acre
- Moderate deficiency: 2-4 quarts per acre
- Severe deficiency: 4-6 quarts per acre

FOR FOLIAR SPRAY APPLICATION

Row Crops: 1 quart per acre every 2-3 weeks with a maximum of 4 applications including but not limited to corn, wheat, soybean, barley and oats.

Vegetable Crops: 1 quart per acre every 10-14 days with a maximum of 4 applications including but not limited to: beans, broccoli, cauliflower, brussels sprouts, cabbage, carrot, spinach, lettuce, potato, onion and tomato.

Fruit and Nut Trees: 1 quart per acre as needed beginning with new growth thru pre bloom and post-harvest when fully leafed out, repeat 3-4 times.

Grapes (all types): 1 quart as needed per bloom and post bloom.

Berries (all types): 1 quart as needed starting with new growth thru pre bloom and prior to leaf fall.

Blueberries, Caneberries, Cranberries, Strawberries: 1 quart per acre as needed. Do not apply after fruit set.

Forage Crops: 1 quart per acre as needed.

*Do not exceed 1 quart per acre for foliar sprays per application.

Notice: The information contained herein is true and accurate to the best of our knowledge and belief, but is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturer nor the seller makes any warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instruction. Read and follow the label directions carefully.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Always read and follow label instructions.

9090 Katy Freeway, Suite 400 Houston, Texas 77024 | www.stollerusa.com | 1-800-539-5283 or 713-461-1493 | @2021 Stoller USA Inc. All rights reserved. Printed in the U.S.A.

Page 2 of 2 Micro-Plus Zinc