

PEEL HERE
TO OPEN ▶



FOLIAR NUTRITIONAL FORMULATION

GUARANTEED ANALYSIS:

Magnesium (Mg)	1.76%
1.76% Water Soluble Magnesium (Mg)	
Sulfur (S)	5.19%
Zinc (Zn)	5.47%

DERIVED FROM: Magnesium sulfate, Zinc sulfate, Kelp extract

KEEP OUT OF REACH OF CHILDREN
KEEP FROM FREEZING

For Product Use Information: 1-866-761-9397

SDS is available at www.arysta-na.com

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

Produced by:

Arysta Lifescience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

F-1721

AD033109V3
103692-C



Danger

Causes serious eye damage

Wear protective gloves, eye protection

If in eyes: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a doctor

Product of France

NET WEIGHT: 26.5 POUNDS (12.02 KILOGRAMS)
NET CONTENTS: 2.5 GALLONS (9.46 LITERS)
WEIGHT PER GALLON: 10.6 POUNDS (4.81 KILOGRAMS) @ 68° F



FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE
CALL PROPHARMA: 1-866-303-6952 or +1-651-603-3432.

FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident)
CALL CHEMTREC: 1-800-424-9300 or +1-703-527-3887.

1. FRUIT TREES

ALMONDS, PECANS, WALNUTS: Dosage per acre per application: 2 quarts. Two applications: 1) Pre-bloom to early bloom. 2) Full bloom to petal fall.

APPLE, PEAR: Dosage per acre per application: 2 quarts. Three applications: 1) Pre-bloom. 2) Early king bloom (for apples)/bloom (for pears). 3) Post bloom (petal fall – fruit set).

BLUEBERRIES & CANE BERRIES: Dosage per acre per application: 1 quart. Three applications: 1) At first bloom 2) At full bloom. 3) At fruit set.

CHERRIES, PEACHES, NECTARINES, APRICOTS, PLUMS: Dosage per acre per application: 2 quarts. Three applications: 1) At first white flower. 2) At petal fall. 3) At shuck fall.

CITRUS & TROPICAL FRUITS: Dosage per acre per application: 2 quarts. Three to six applications: 1) All fruits: A) At 2/3 petal fall. B) With the June spray. C) With the Fall miticide or scale spray. For varieties with heavy fruit set problems: Make an additional application at pre-bloom. 2) Navels and other varieties susceptible to premature fruit drop: A) June drop, spray 10 to 14 days before drop starts. B) Fall drop (from August to harvest), spray 10 to 20 days before drop starts. C) Grapefruits and other fruits to be held on the trees for later markets (export), spray 30 to 60 days prior to expected harvest. Do not apply after the end of January or when trees begin to break dormancy.

2. VEGETABLES

ARTICHOKE, BROCCOLI, CAULIFLOWER: Dosage per acre per application: 2 quarts. Three applications: 1) Within 7 days of transplant or 100% emergence. 2) When buds just become visible. 3) 14 days later.

ASPARAGUS: Dosage per acre per application: 3 pints. 1) Branches formation. 2) 15 days after first treatment. 3) 15 days after second treatment.

BEANS, PEAS & OTHER LEGUMES: Dosage per acre per application: 2 quarts. Three applications: 1) At 2-3 trifoliate leaves. 2) First bloom. 3) First pod.

CABBAGE, LETTUCE, SPINACH: Dosage per acre per application: 2 quarts. Three applications: 1) Within 7 days of transplant or 100% emergence. 2) When buds just become visible. 3) 14 days later.

CANTALOUPE, HONEYDEW, WATERMELON: Dosage per acre per application: 2 quarts. Four applications: 1) Within 7 days of transplant or 100% emergence. 2) Just prior to first blooming. 3) 7 to 10 days after second application. 4) 7 to 10 days after third application.

3. ROW CROPS

ALFALFA HAY: Dosage per application: 1 to 2 quarts. Three to four applications: Within 4-6 days after each hay cutting.

CORN: Dosage per acre per application: 2 quarts. Two applications: 1) At 5-6 leaf stage. 2) At 10-12 leaf stage.

COTTON: Dosage per acre per application: 1 quart. One application: 1) At early to mid-bloom.

RICE: Dosage per acre per application: 2 quarts. One application: 1) At early shooting stage (one visible joint).

SUGAR BEET: Dosage per acre per application: 2 quarts. Two applications: 1) 5-6 true leaf stage. 2) 10-12 true leaf stage.

SUNFLOWER: Dosage per application: 2 quarts. One application: At 6-8 true leaf stage.

WHEAT, BARLEY, OATS, SORGHUM, MILLET: Dosage per acre per application: 1 quart. Two applications: 1) Early shooting (one to two joints). 2) First visible heads.

1. FRUIT TREES

GRAPES: (Raisin, Table and Wine): Three applications: 1) 12" shoot growth, 2 quarts per acre. 2) At bloom stage, 1 quart per acre. 3) After fruit set, 1 quart per acre.

HAZELNUTS (FILBERTS): Dosage per acre per application: 2 quarts. Three applications: 1) At ovary development (late May). 2) Ovary fertilization (mid-June). 3) Young fruit development (late June, early July).

STRAWBERRIES: Dosage per acre per application: 1 quart. 1) Within 14 days of transplant. 2) At first bloom. 3) 14-21 days later. Subsequent applications: Every 14-21 days thereafter.

2. VEGETABLES

CARROTS, RADISHES: Dosage per acre per application: 2 quarts. Two applications: 1) When plants are 4-6 inches tall. 2) 14-20 days later.

CELERY: Dosage per acre per application: 3 pints. Five to six applications: 1) Within 14 days of transplant. 2) Every 14 days until harvest.

CUCUMBERS, SQUASH: Dosage per acre per application: 3 pints. Four to five applications: 1) Within 7 days of transplant. 2) Every 7-10 days thereafter.

ONIONS: Dosage per acre per application: 2 quarts. Two applications: 1) When plants are 4-6 inches tall. 2) 15 days after first application.

PEPPERS & TOMATOES (Fresh Market): Dosage per acre per application: 3 pints. Four applications: 1) Within 7 days of transplant. 2) At first flower. 3) 7-14 days later. 4) 7-14 days later.

POTATOES: Dosage per acre per application: 3 pints. Two applications: 1) 100% emergence. 2) 15 days later.

3. ROW CROPS

PEANUTS: Dosage per acre per application: 1 quart. Two applications: 1) At pegging. 2) 10-14 days later.

FOR APPLICATION TO ORNAMENTALS, NURSERIES AND COMMERCIAL LANDSCAPES.

Application Rate:

(ORNAMENTAL) NURSERY & GREENHOUSE: Use 2 quarts per 100 gallons of water on plants, flowers, shrubs and trees for direct spray to foliage. Apply 3 applications beginning at pre-bloom, mid-bloom and post bloom. Can be tank mixed with common fungicides and insecticides.

NOTICE: Do not tank mix with oil sprays.

KEEP FROM FREEZING

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Arysta LifeScience North America, LLC (“Arysta”), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

Arysta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Arysta, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA, MANUFACTURER, OR THE SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE BUYER AND THE USER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF ARYSTA IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER OR USER’S EXCLUSIVE REMEDY, AND ARYSTA’S, THE MANUFACTURER’S, OR THE SELLER’S TOTAL LIABILITY SHALL BE LIMITED TO DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT, OR AT ARYSTA’S ELECTION, THE REPLACEMENT OF THE PRODUCT.

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation. MZ’O’ is a trademark of Laboratoires Goëmar.



FOLIAR NUTRITIONAL FORMULATION

GUARANTEED ANALYSIS:

Magnesium (Mg)	1.76%
1.76% Water Soluble Magnesium (Mg)	
Sulfur (S)	5.19%
Zinc (Zn)	5.47%

DERIVED FROM: Magnesium sulfate, Zinc sulfate, Kelp extract

KEEP OUT OF REACH OF CHILDREN KEEP FROM FREEZING

For Product Use Information: 1-866-761-9397
SDS is available at www.arysta-na.com

FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL PROPHARMA: 1-866-303-6952 or +1-651-603-3432.
FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC: 1-800-424-9300 or +1-703-527-3887.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

Produced by:

Arysta Lifescience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

F-1721

AD033109V3
103692-C



Danger

Causes serious eye damage
Wear protective gloves, eye protection
If in eyes: Rinse cautiously with water for several minutes.
Remove contact lenses, if present and easy to do. Continue rinsing.
Immediately call a doctor

Product of France

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.
MZ'O' is a trademark of Laboratoires Goëmar.

NET WEIGHT: 26.5 POUNDS (12.02 KILOGRAMS)
NET CONTENTS: 2.5 GALLONS (9.46 LITERS)
WEIGHT PER GALLON: 10.6 POUNDS (4.81 KILOGRAMS) @ 68° F

