



## INSTRUCTIONS

**PREPARATION:** Adds naturally occurring materials containing humic and fulvic acids to soil that may enhance soil health for plant nutrient availability and uptake, water use efficiency, growth of beneficial microbes, and crop vigor to overcome abiotic stress. Should be applied when tissue and soil analysis identify the need and proper rate of application for this product. Can be applied in combination with compatible granular fertilizers. Avoid mixing with high-moisture fertilizers such as granular urea.

**APPLICATION:** Designed to be applied to the soil. Best results will be obtained when applied directly to the soil followed by shallow incorporation or irrigation. Apply by broadcast, whether through air or ground machine. Can be applied to all soil types for agricultural fields, ornamentals, lawns, gardens, and landscapes. Avoid spreading during high humidity. Soil moisture is required for maximum bioactivity: if soil is dry, moisture should be provided by irrigation. Applications can be made as often as every 30 days, as needed.

**CONDITIONS:** Application timing, intervals, and rates may vary according to individual plant requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the plants.

**CONSULT:** Consult your local Huma® Representative or qualified individual/institution for use-specific recommendations.

**STORAGE & DISPOSAL:** Keep product in original bag. Do not transfer into food or drink containers. Always dispose of bag in accordance with local, state, and/or federal regulations.

**DISCLAIMER:** This product is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by plants.

## GUARANTEED ANALYSIS

CONTAINS BENEFICIAL SUBSTANCES:

Total Humic Acid (HA) and Fulvic Acid (FA) ..... 70.0%  
(Analysis using the Colorimetric Test Method)

Other Test Methods:

Total Humic Acids (HA) ..... 40.0%  
(Analysis using the HPTA or ISO 19822 Method)

Total Humic Acids (HA) ..... 35.0%  
(Analysis using the CDFA Method)

Derived from leonardite; less than 3% dust and 18% moisture.

## SOIL AMENDMENT

Applications can be made as often as every 30 days, as needed.

## APPLICATION

METHOD	SUGGESTED RATE		
	Field Crops, Sod, and Specialty Crops	Tree or Vine Crops	
Soil banded or side dressed	Up to 75 lb/ac, 75 kg/ha	7 lb/1,000 ft <sup>2</sup> , 3 kg/100 m <sup>2</sup>	Up to 150 lb/ac, 150 kg/ha
Soil broadcast incorporated	Up to 150 lb/ac, 150 kg/ha	14 lb/1,000 ft <sup>2</sup> , 6 kg/100 m <sup>2</sup>	Up to 300 lb/ac, 300 kg/ha

Product Density: 45–48 lb/ft<sup>3</sup>

Net Wt. 50 lb  
Net Wt. 22.68 kg

## SAFETY

KEEP OUT OF REACH OF CHILDREN

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.html>

Ingestion of this product may cause gastrointestinal irritation or pain. Inhalation of dust may cause respiratory irritation.

**CLASSIFICATION:** RESPIRATORY TRACT IRRITATION—HAZARD CATEGORY 3

**SIGNAL WORD:** WARNING

**HAZARD STATEMENT:** H335; May cause respiratory irritation.

**PRECAUTIONARY STATEMENT:** P261, P271; Avoid breathing dust. Use only outdoors or in a well-ventilated area. P304+P340, P319; IF INHALED: Remove person to fresh air and keep comfortable for breathing. Get medical help if you feel unwell.

**PRODUCT CHARACTERISTICS:** A dark reddish brown to black particles/dust having no odor.

**CONDITIONS OF SALE:** The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, plant and soil conditions, and other factors are beyond the control of the seller.



Guaranteed by Huma, Inc.  
1331 W. Houston Ave., Gilbert, AZ 85233  
480.961.1220 © 2024  
F003537  
Publication No. HP&S-644-240612

[huma.us](http://huma.us)