

 ELEMENT
 %W/V

 N
 6.90

 B
 15.00

DIRECTIONS FOR USE: SHAKE WELL BEFORE USE

APPLICATION RATES:

Foliar Applications:

Several applications may be required, 7-10 days apart.

Apples:

Apply 2 x 1-2L/ha.

Apply at open cluster to early pink (Do not apply with spraying oil). Repeat at Post Harvest.

Stone Fruit: Apply 1-2L/ha up to early fruit set.

Avocado: Apply 2-3L/ha up to early fruit set.

Grape Vines Apply 2 x 1-5L/ha.

1st cluster visible, 2nd flower buds separated.

Vegetables: Apply 2-3L/ha.

Bananas: Apply 0.5-1L/ha. Apply monthly.

Canola: Apply 1-2L/ha.

Fertigation / Soil Applications

Apply as required but do not exceed 5L/ha per application. Several applications may be required.

When mixing with other liquid fertilizers, test for compatibility with a small quantity in a jar.

DO NOT EXCEED RECOMMENDED RATES



OMNIBOR

15% BORON, 6.9 NITROGEN

RAPIDLY AVAILABLE BORON
SUITABLE FOR FOLIAR AND FERTIGATION USE.



FOLIAR



OmniBor

OmniBor has the following benefits:

- High analysis flowable and available
- Addition of Nitrogen for enhanced uptake
- Ready to use formulation
- Ideal for foliar application and fertigation
- Effective treatment for all crops

CAUTION

Avoid contact with eyes. Keep out of reach of children. If swallowed seek medical advice.

Do not store in direct sunlight.

Omnia Specialities Australia Pty Ltd

(ACN 102 717 019) Tramway Road Morwell 3840

Victoria, Australia

Tel: +61-3-5133 9118 Fax: +61-3-5133 9114

CONDITIONS OF SALE:

www.omnia.com.au

Every care is taken during the manufacture of this product. However, as environmental conditions can vary considerably, no responsibility can be taken by the manufacturer, importer or his agents for any injury, damage, loss or other results following the use or misuse of this product.

PROUDLY MADE IN AUSTRALIA