

Orchard Clean[®]

Broadleaf Herbicide

THE 2,4-D AMINE WEED KILLER

FOR CONTROL OF MANY BROADLEAF WEEDS IN GRAPE VINEYARDS AND ORCHARD FLOOR (APPLE, PEAR, STONE FRUIT AND NUT).

ACTIVE INGREDIENT:

2,4-Dichlorophenoxyacetic acid, dimethylamine salt* 46.8%

OTHER INGREDIENTS: 53.2%

TOTAL: 100.0%

*2,4-Dichlorophenoxyacetic acid equivalent 38.9% by weight or 3.8 pounds per gallon. Isomer specific by AOAC method No. 978.05

**KEEP OUT OF REACH OF CHILDREN
DANGER / PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS

For Medical Emergencies, Call (877) 325-1840.

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.

EPA Reg. No. 71368-1

Manufactured for
Nufarm, Inc.
11901 S. Austin Avenue
Alsip, IL 60803



1 91662 01488 4

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIANS	
This product contains a phenoxy herbicidal chemical. There is no specific antidote. All treatments should be based on observed signs and symptoms of distress in the patient. Probable mucosal damage may contraindicate the use of gastric lavage.	

 **Nufarm**
Grow a better tomorrow

Net Contents
2.5 Gal.
(9.46 L)

Nonrefillable Container

12821000

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER / PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

All mixers, loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks, plus
- chemical-resistant gloves including barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, or Viton ≥ 14 mils, when applying with any handheld nozzle or equipment, mixing loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate, and
- protective eyewear.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-e)], the handler PPE (personal protective equipment) may be reduced or modified as specified in the WPS. Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.607(d-e)].

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinseate.

This product contains a chemical with properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

For Aquatic Uses: Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, apply the product in lanes separated by untreated strips that can be treated after vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Waters having limited and less dense weed infestations may not require partial treatments.

MIXING AND LOADING: Most cases of ground water contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent ground water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical-resistant gloves made of any water-proof material, shoes plus socks, protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

PRODUCT INFORMATION

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this product only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company, State Agricultural Extension Service or qualified crop consultant for advice.

Be sure that use of this product conforms to all applicable laws, rules and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator must become familiar with these laws, rules or regulations and follow them exactly.

USE RESTRICTIONS

Do not apply this product through any type of irrigation system. Do not use in or near a greenhouse. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not contaminate water used for irrigation or domestic purposes (except as specifically listed on this label) especially in areas where grapes, cotton, tomatoes or other susceptible plants are grown.

Do not treat irrigation ditches in areas where water will be used to overhead (sprinkler) irrigate susceptible crops especially grapes, tomatoes, tobacco, and cotton.

Do not apply this product directly to, or permit to drift onto cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants which are susceptible to 2,4-D herbicide. Do not apply near susceptible plants since very small quantities of the 2,4-D will cause severe injury during the growing or dormant periods. Crops contacted by this product sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

WEED RESISTANCE MANAGEMENT

For resistance management, this product contains a Group 4 herbicide –2,4-D. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and that considers mechanical control methods, cultural (e.g., timing to favor the desirable plants and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - (2) a spreading patch

of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest controls advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of plants and weed biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. Do not assume that each listed weed is being controlled by this mechanism of action. Co-formulated active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredients in this product.

MIXING INSTRUCTIONS

Add about one-half the water to the mixing tank, then add this product with agitation and finally the rest of water with continuing agitation.

NOTE: Adding oil, wetting agent, or other surfactants to the spray may increase effectiveness on weeds but also may reduce selectivity to crops, resulting in crop damage.

COMPATIBILITY

If this product is to be tank mixed with fertilizers or with other pesticides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing.

Read and follow all directions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

APPLICATION PROCEDURES

Apply by air or ground equipment in sufficient gallonage to obtain adequate coverage, except as otherwise directed on this label. Use 2 or more gallons of water per acre for aerial application and 10 or more gallons of water per acre for ground application.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASABE standard 572).

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASABE standard 572).

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

SMALL QUANTITY DILUTION TABLE

To spray small areas use the following dilution table.

If Dosage on Label Shows Following Rate Per Acre	Use this Amount for each Gallon of Water Per 1,000 Square Feet
2 Pints (1 quart)	0.72 ounces (4.3 teaspoons)
4 Pints (2 quarts)	1.4 ounces (2.8 tablespoons)
6 Pints (3 quarts)	2.2 ounces (4.4 tablespoons)

SPOT TREATMENTS

To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers using a fixed spray volume per 1,000 sq ft as indicated below. Hand-Held Sprayers: Hand-held sprayers may be used for spot applications of this product. Apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on the application rate for an area of 1,000 sq ft. Mix the amount of this product (fl oz or ml) corresponding to the desired broadcast rate in 1 to 3 gallons of spray. To calculate the amount of this product required for larger areas, multiply the table value (fl oz or ml) by the thousands of sq ft to be treated. An area of 1000 sq ft is approximately 10.5 x 10.5 yards (strides) in size.

Rate Conversion Table for Spot Treatment:

Label Broadcast Rate (pints per acre)						
1/2	2/3	3/4	1	2	3	4
Equivalent Amount of Orchard Clean® per 1000 sq ft						
1/5 fl oz* (5.5 ml)	1/4 fl oz (7.3 ml)	1/3 fl oz (8.3 ml)	3/8 fl oz (11 ml)	3/4 fl oz (22 ml)	1 fl oz (33 ml)	3 fl oz (88 ml)

* Conversion factors: 1 pt - 16 fl oz; 1 fl oz = 29.6 (30) ml

Band Application: Orchard Clean® may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated area.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre} = \text{Band rate per treated acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast volume per acre} = \text{Band volume per treated acre}$$

WEED LIST

Annual or Biennial Weeds

Beggarticks (1) Bittercress, smallflowered (3) Bitterweed Broomweed, common (1) Burdock, common Buttercup, smallflowered (1)(3) Carrot, wild Carpetweed Cinquefoil, common (3) Cinquefoil, rough (3) Cocklebur, common Coffeeweed Copperleaf, Virginia (1)(3) Croton, Texas Croton, woolly Fleabane, rough Flixweed Galinsoga Geranium, Carolina (3) Hemp, wild Horseweed (marestail) (3) Jewelweed Jimsonweed	Knotweed (1) Kochia Lambsquarter, common Lettuce, prickly (1) Lettuce, wild Lupines Mallow, little (1) Mallow, Venice (1) Marshelder Morningglory, annual Morningglory, common Morningglory, ivy Morningglory, woolly Mousetail (3) Mustards (except blue mustard) Parsnip, wild Pennycress (fanweed) Pepperweeds (<i>Lepidium</i> spp.)(1) Pigweeds (<i>Amaranthus</i> spp.)(2) Poorjoe Primrose, common evening Purslane, common (3) Pusley, Florida	Radish, wild Ragweed, common Ragweed, giant Rape, wild Rocket, yellow Salsify, common (1) Salsify, western (1) Shepherd's purse Sicklepod Smartweed (annual species)(1) Sneezeweed, bitter Sowthistle, annual Sowthistle, spiny Spanish needles Sunflower Sweetclover Tansy mustard Thistle, bull Thistle, musk(1) Thistle, Russian (tumbleweed)(1) Velvetleaf Vervains(1) Vetches
--	--	--

WEED LIST (continued)

Perennial Weeds

Alfalfa (1)(3)	Catnip	Ironweed (1)
Artichoke, Jerusalem (1)	Chicory	Ivy, ground (1)
Aster, many-flower (1)	Clover, red (1)(3)	Nettles (including stinging) (1)
Austrian fieldcress (1)	Coffeeweed	Onion, wild (1)
Bindweed, European (1)	Cress, hoary (1)	Pennywort
Bindweed, field (1)	Dandelion	Plantains
Bindweed, hedge (1)	Docks (1)	Ragwort, tansy (1)
Blue lettuce	Dogbanes (1)	Sowthistle, perennial
Blueweed, Texas	Eveningprimrose, cutleaf (3)	Speedwell
Broomweed	Garlic, wild (1)	Spotted catsear
Bull nettle (1)(3)	Goldenrod (1)	Thistle, Canada (1)
Carrot, wild	Healweed, orange (1)	Vervains (1)
	Healall	Wormwood

1- These species may require repeat applications and/or use of the higher specified rate even under ideal conditions for application.

2- Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product

3- This product may not be used to control this weed species in the State of California.

CROP SPECIFIC USE DIRECTIONS

APPLES, PEARS, STONE FRUIT AND NUT ORCHARDS (EXCEPT FILBERTS) (ORCHARD FLOOR)

APPLICATION TIMING	AMOUNT OF ORCHARD CLEAN® PER ACRE	DIRECTIONS
Postemergence		
Annual and biennial broadleaf weeds	1 to 2 pints	For application to orchard floors, use coarse, low-pressure sprays and sufficient water for thorough coverage of weeds. Apply to annual weeds when small and actively growing.
Perennial broadleaf weeds	4 pints	Apply to perennial weeds from bud to bloom stage. Because newly established trees or young orchards are more susceptible to 2,4-D injury, apply only to orchards that are at least one year old and well-established as indicated by vigorous plant growth.

RESTRICTIONS FOR USE IN APPLES, PEARS, STONE FRUIT AND NUT ORCHARDS (EXCEPT FILBERTS)

- Do not apply immediately before irrigation and withhold irrigation for 2 days before and for 3 days after treatment.
- Do not apply to newly established or young orchards. Trees must be at least 1 year old and in vigorous condition.
- Do not apply during bloom.
- Do not graze or feed cover crops from treated orchards.
- Do not make more than 2 applications per crop cycle. Maximum of 4.2 pints (2.0 lbs. ae) per acre per application.
- For apples, pears and stone fruits, allow at least 75 days between applications.
- For tree nuts, allow at least 30 days between applications.
- Do not cut orchard floor forage for hay within 7 days of application.
- Orchard Clean® contains 3.8 pounds a.e. of 2,4-D per gallon. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.
- **(PHI) Preharvest Intervals:**
 - **Apple and Pear:** Do not harvest within 14 days of application.
 - **Stone Fruit:** Do not harvest within 40 days of application.
 - **Nut Orchard and Pistachio:** Do not harvest within 60 days of application.

Use Precautions

- Do not apply to bare ground as injury may result.
- Do not allow spray to drift onto or contact foliage, fruit, stems, trunks of trees or exposed roots as injury may result.

FILBERTS (ORCHARD FLOOR)

WEEDS IN CROPS	AMOUNT OF ORCHARD CLEAN® PER ACRE	DIRECTIONS
Annual broadleaf weeds	2.1 pints	Apply a maximum of 2.1 pints (1.0 lb. ae) in 100 gallons of spray solution per acre.

RESTRICTIONS FOR USE IN FILBERTS

- Do not use on light sandy soil.

- Do not apply immediately before irrigation and withhold irrigation for 2 days before and for 3 days after treatment.
- Do not apply to newly established or young orchards. Trees must be at least 1 year old and in vigorous condition.
- Do not apply during bloom.
- Do not graze or feed cover crops from treated orchards.
- Do not make more than 4 applications per crop cycle. Maximum of 2.1 pints (1.0 lbs. ae) per acre per application.
- Orchard Clean® contains 3.8 pounds a.e. of 2,4-D per gallon. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 1.0 pounds of a.e. per acre per year.
- (PHI) Do not harvest filberts with 45 days of application.
- Allow at least 30 days between applications.
- Do not cut orchard floor forage for hay within 7 days of application.

Use Precautions

- Do not apply to bare ground as injury may result.
- Do not allow spray to drift onto or contact foliage, fruit, stems, trunks of trees or exposed roots as injury may result.

GRAPE VINEYARDS

Established at least 3 years to control Field Bindweed (Morning Glory), Canada Thistle and other 2,4-D susceptible broadleaf weeds.

APPLICATION TIMING	AMOUNT OF ORCHARD CLEAN® PER ACRE	DIRECTIONS
Apply when weeds are in the bud to early bloom stage and growing vigorously. Apply after shatter following bloom and before grape shoots reach the ground or during dormant season.	1.8 to 2.7 pints	Dilute in 10 to 100 gallons of water to treat one acre of ground to be sprayed. For band or spot treatment, calculate rates according to the actual portion of acre treated. Use a hooded boom and low pressure flooding nozzles to deliver coarse droplets.

RESTRICTIONS FOR USE ON GRAPES

- For use only in California, Oregon and Washington.
- The pre-harvest interval (PHI) is 100 days.
- Limited to 1 application per crop cycle.
- Maximum of 2.7 pints (1.36 lbs. ae) per acre per application.
- Orchard Clean® contains 3.8 pounds a.e. of 2,4-D per gallon. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 1.36 pounds of a.e. per acre per year.

Precaution: Grapes are extremely sensitive to 2,4-D. Use a direct application so no 2,4-D contacts grape leaves and young shoots or stems

TANK MIXES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Using this product and **Harmony®** (EPA Reg No. 279-9577, active ingredient thifensulfuron), **Treaty®** (EPA Reg No. 71368-74, active ingredient thifensulfuron), **Harmony® Extra** (EPA Reg No. 279-9583, active ingredient thifensulfuron), or **Treaty® Extra** (EPA Reg No. 71368-76, active ingredient thifensulfuron) for selective post-emergence control of certain weeds on wheat (including Durum) and barley: Use Harmony Extra plus 1/8 to 3/8 pound active ingredient 2,4-D. Surfactant may be added 0.125 to 0.25% vol/vol (1 pint to 1 quart per 100 gallons of spray volume); however, the addition of surfactant may increase the chance of crop injury. Use the 1 pint rate of surfactant with 1/4 to 3/8 pound active ingredient of 2,4-D. Higher rates of 2,4-D may be used, but do not exceed highest rate allowed on the label. Always mix Harmony, Treaty, Harmony Extra or Treaty Extra in water prior to adding 2,4-D. Always add surfactant last.

Using this product and **Ally®** (EPA Reg No. 279-9575, active ingredient metsulfuron) or **Purestand®** (EPA Reg No. 71368-38, active ingredient metsulfuron) for selective weed control in wheat (including Durum), barley and in grasses in acreage enrolled in the Conservation Reserve Programs (CRP) - also for resistant weed management: Use Ally at 1/10 ounce/A plus 1/4 to 1/2 pound active ingredient 2,4-D. Surfactant may be added at one to two pints per 100 gallons of spray volume; however, the addition of surfactant may increase the chance of crop injury. Always mix Ally or Purestand in water prior to adding 2,4-D and surfactant. Always add surfactant last.

Using this product and **Express®** (EPA Reg No. 279-9578, active ingredient tribenuron-methyl) or **Victory®** (EPA Reg No. 71368-75, active ingredient tribenuron-methyl) tank mixtures for weed control in wheat and barley: Use Express plus 1/8 to 3/8 pound active ingredient 2,4-D. Surfactant may be added at 0.125 to 0.25% vol/vol (1 to 2 pints per 100 gallons of spray volume); however, the addition of surfactant may increase the chance of crop injury. Use the 1 to 2 pint rate of surfactant with 1/8 pound active ingredient of 2,4-D. Use the 1 pint rate of surfactant with 1/4 to 3/8 pound active ingredient of 2,4-D. Higher rates of 2,4-D may be used, but do not exceed highest rate allowed on the label. Always mix Express or Victory in water prior to adding 2,4-D and surfactant.

Using this product and **Glean®** (EPA Reg No. 279-9600, active ingredient chlorsulfuron) for post-emergent weed control in wheat, barley and oats: Mixtures of 2,4-D and Glean are recommended when weeds are large and/or stressed due to adverse environmental conditions (cold temperature, low soil moisture, dry, dusty field conditions) or when dense crop canopy makes it difficult to obtain thorough spray coverage. Use 1/4 to 1/2 pound active ingredients 2,4-D plus 1/6 to 1/3 ounce/A of Glean. Surfactant may be added at 1/2, but not more than 1 quart/100 gallons of spray; however, the addition of surfactant may increase the chance of crop injury. Glean should be mixed in water with agitator running prior to adding 2,4-D. For resistant weed management, see Glean label rates for different regional applications.

Using this product and **Buctril® 4EC** (EPA Reg No. 264-540, active ingredient bromoxynil) or **Maestro® 4EC** (EPA Reg No. 71368-78, active ingredient bromoxynil) for weed control on cereal grains (wheat, barley and rye): Buctril 4EC Broadleaf Herbicide or Maestro 4EC Selective Herbicide will control some annual weeds that are resistant to

this product and may be tank mixed with this product for broader spectrum weed control on small grains. In cereal areas except Washington, Oregon and Idaho, use 1/2 to 1 pint of this product plus 1/2 to 3/4 pint of Buctril 4EC or Maestro 4EC per acre. In Washington, Oregon and Idaho: use 1/2 to 1 pint of this product plus 3/4 to 1 pint of Buctril 4EC or Maestro 4EC per acre. First mix this product in water then add the Buctril 4EC or Maestro 4EC. Use the higher rates for larger weeds or where weed growth is slow due to dry or cold weather. Apply before weeds are 6 inches high. Use 10 to 20 gallons total spray volume per acre with ground equipment or 5 to 10 gallons total spray volume with air application. Use higher volume on larger weeds. This product and Buctril 4EC or Maestro 4EC can also be tank mixed for field and popcorn. See both product labels for rates of application.

Using this product and **Relegate**® (EPA Reg No. 228-521, active ingredient triclopyr) or **Tahoe**® 3A (EPA Reg No. 228-517, active ingredient triclopyr) tank mixtures for Non-Crop Areas: **Broadleaf Weed Control** - Use 2 to 4 pints of this product plus 2 to 6 pints of Relegate (or 3 to 8 pints of Tahoe 3A) per acre. For wider spectrum control of broadleaf weeds and woody plants, apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing. **Woody Plant total Broadcast Foliar Spray** - Use 1 gallon of this product plus 1-1/2 to 3 quarts of Relegate (or 2 to 4 quarts of Tahoe 3A) per acre. Apply as a broadcast spray in enough water to adequately wet foliage. This may require 20 to 100 gallons of water per acre. Apply when woody plants are actively growing. **Woody Plant Control High Volume Leaf-Stem Treatment with Ground Equipment** - Use 1 gallon of this product per acre plus 1-1/2 to 12 pints of Relegate (or 2 to 16 pints of Tahoe 3A). Mix 1 to 3 quarts of this product per acre plus 1-1/2 to 3 pints of Relegate (or 2 to 4 pints of Tahoe 3A) in enough water to wet all parts of the brush foliage, stem and bark. This may require 100 to 400 gallons of water per acre depending on size and density of woody plants. Thoroughly wet all leaves, stems, and root collars of plants to be controlled. **Woody Plant Control Aerial Application (Helicopter only)** - Use 1 gallon of this product per acre plus 3 to 4 quarts of Relegate (or 4 to 6 quarts of Tahoe 3A). Thoroughly wet all leaves, stems, and root collars of plants to be controlled. This may require 10 to 30 gallons of water per acre using drift control equipment such as Microfoil boom or an effective drift control agent such as Lo-Drift Spray Additive. Use the higher rates and volumes when plants are dense or under drought conditions.

Using this product and **Diablo**® Herbicide (EPA Reg No. 228-379, active ingredient dicamba) tank mixtures for Non-Crop Areas: **Annual broadleaf weeds** - Use 2 to 4 pints of this product plus 1/2 to 1-1/2 pints of Diablo. For wider spectrum control of broadleaf weeds and woody plants - Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons of water per acre. Apply when broadleaf weeds are actively growing. Use the higher rates when treating dense or tall vegetative growth. **Perennial and Biennial Broadleaf Weeds** - Use 3 to 6 pints of this product per acre plus 1/2 to 6 pints Diablo. Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons of water per acre. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches rosette diameter. Use the higher rates for perennial weeds or for biennial weeds past the 3 inch rosette stage. **Woody Plant Control Broadcast, High Volume, Stem Foliage or Aerial Application** - Use 1 gallon of this product plus 2 to 8 quarts of Diablo. Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre or apply as a high volume stem foliage spray in enough volume to thoroughly wet leaves, stems and root collars (100 to 400 gallons per acre) or apply aerially in enough water to wet all parts of the brush foliage, stem and bark. This may require 10 to 30 gallons of water per acre using drift control equipment such as the Microfoil Boom or an effective drift control agent such as Lo-Drift Spray Additive. Use the higher rates and volumes when plants are dense or under drought conditions.

Tank mixing 2,4-D with Diablo Herbicide will improve control of late, post-emergent applications in corn, fall and spring seeded wheat, fall seeded barley, pastures, rangelands, non-crop use, turfs and lawns.

This product may be tank mixed with **Banvel SGF**® (EPA Reg No. 66330-287, active ingredient 2,4-D and dicamba) for control of grasses or additional broadleaf weeds in fall seeded wheat and barley. This product may be tank mixed with **Veteran 720** (EPA Reg No. 228-295, active ingredient 2,4-D and dicamba) for broader spectrum control for non-crop uses (e.g. railroad, highway, pipelines, etc. including forest management applications). Add water to spray tank prior to addition of tank mix products. Do not pre-mix concentrates. Since Veteran 720 contains 2,4-D, do not exceed 16 lbs. total 2,4-D acid equivalent per treated acre per growing season.

For control of Quackgrass and listed weeds in annual cropping systems, pastures and sods, this product may be mixed with **Ranger**® - refer to specific product label for use rates.

This product may be tank mixed with **Tordon**® 22K (EPA Reg No. 62719-6, active ingredient picloram-potassium) or **Trooper**® 22K (EPA Reg No. 228-535, active ingredient picloram-potassium). For use on areas having mixed species in non-cropland range, pasture wheat, barley, oats and fallow cropland.

Using this product and **Razor**® (EPA Reg No. 228-336, active ingredient glyphosate-isopropylammonium) will control annual grasses and broadleaf weeds listed for Razor alone plus the following broadleaf weeds: Lambsquarter, Prickly lettuce, Red root, Pigweed, Russian thistle, Velvet leaf. Fallow and reduced tillage areas only. Apply 12 to 16 ounces of Razor plus 1/2 pound acid equivalent of this product plus 1/2 to 1% nonionic surfactant by total spray volume per acre to control dense populations of the aforementioned weeds when less than 6 inches in height. Follow use directions as given in the "low-volume broadcast application" section of the Razor label.

For high-volume broadcast applications:

When weeds are less than 6 inches tall, increase the quantity of Razor to 1 quart; when weeds are over 6 inches tall, use 1-1/2 quarts of Razor per acre. In both instances, water volumes should be 10 to 40 gallons per acre for ground applications. If weeds have been mowed, graded, or cut, allow adequate time for new growth to recommended stages prior to treatment. These rates will also provide control of weeds listed in the low-volume broadcast application section in addition to the following: Fivehook bassia, Broom fiddleneck, Flaxleaf fleabane, Fleabane, Kochia, Prickly lettuce, Panicum, Common ragweed, Giant ragweed, Pennsylvania smartweed, Annual sowthistle, Sunflower, Russian thistle, Velvetleaf. For Balsam Apple, apply with hand-held equipment only. A tank mix of Razor and this product will also control most perennial weeds. See Razor label for specifics. For additional tank mixes in fallow and reduced tillage systems for control of annual weeds prior to emergence of crops, please see Round Up RT label.

For additional non-crop weed control benefits, up to 1-1/2 quarts per acre of this product may be added to tank mixes of Razor plus Telar®, Razor plus Spyder®, Razor plus Patriot® for the suppression of tall Fescue growth and seed heads and control, or partial control, of some annual weeds. For the suppression of Smooth brome growth and seed heads and control or partial control of some annual weeds, 1-1/2 quarts per acre of this product may be added to a tank mix of Razor plus Spyder.

Using this product and **Acclaim**®1 EC Herbicide (EPA Reg No. 432-950, active ingredient fenoxaprop-p-ethyl) may be added to provide broadleaf weed and annual grassy weed control in Turfgrass including sod farms, commercial and residential turf. Apply before grassy weed tillering at a rate of 32 ounces of Acclaim per acre (or 0.73 ounces per 1,000 square feet) when mixing with this product at a rate of 3/4 to 3 pints per acre (or 0.9 - 1.1 ounces per 1,000 square feet). Apply by means of a pressurized hydraulic sprayer using 30 to 60 psi and 30 to 60 gpa. Thorough spray coverage is important. Flat fan nozzles are recommended. Always follow use directions in accordance with respective labels. No label dosage rates should be exceeded.

Using this product and **Patriot®** (EPA Reg No. 228-391, active ingredient metsulfuron) **Spyder®** (EPA Reg No. 228-408, active ingredient sulfometuron) and **Telar®** (EPA Reg No. 432-1561, active ingredient chlorsulfuron) To improve control of some target species, this product may also be tank mixed with Patriot, Spyder, and Telar herbicides for non-crop, post-emergent weed control. Tank mixes have shown improved control where resistant bio-types are present.

NOTE: All intended tank mix combinations should be used only in recommended areas on the same broadleaf weed species found on both labels. For application methods and other use specifications, use the most restricted limitations from labeling of both products.

This product will either kill, control or suppress the weeds listed in the label booklet for this product. Some of these species may require repeat spot applications even under ideal conditions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry, secured storage area. Keep container tightly closed when not in use. Store at temperature above 32°F. If allowed to freeze, warm to at least 40°F and remix before using. Freezing does not alter this product.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

NOTE: This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type / size.

Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

Nonrefillable Containers Larger than 5 Gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

Refillable Container Larger than 5 Gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

RV012621

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Orchard Clean is a Registered Trademark of Nufarm Americas Inc.

All other trademarks are the property of their respective owners.

Orchard Clean®

Broadleaf Herbicide

THE 2,4-D AMINE WEED KILLER

FOR CONTROL OF MANY BROADLEAF WEEDS IN GRAPE VINEYARDS AND ORCHARD FLOOR (APPLE, PEAR, STONE FRUIT AND NUT).

ACTIVE INGREDIENT:

2,4-Dichlorophenoxyacetic acid, dimethylamine salt* 46.8%

OTHER INGREDIENTS: 53.2%

TOTAL: 100.0%

*2,4-Dichlorophenoxyacetic acid equivalent 38.9% by weight or 3.8 pounds per gallon.

Isomer specific by AOAC method No. 978.05

KEEP OUT OF REACH OF CHILDREN DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

For Medical Emergencies, Call (877) 325-1840.
For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER / PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIANS	
This product contains a phenoxy herbicidal chemical. There is no specific antidote. All treatments should be based on observed signs and symptoms of distress in the patient. Probable mucosal damage may contraindicate the use of gastric lavage.	

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry, secured storage area. Keep container tightly closed when not in use. Store at temperature above 32°F. If allowed to freeze, warm to at least 40°F and remix before using. Freezing does not alter this product.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

NOTE: This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type / size.

Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

Nonrefillable Containers Larger than 5 Gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

Refillable Container Larger than 5 Gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

EPA Reg. No. 71368-1

Manufactured for Nufarm, Inc.
11901 S. Austin Avenue Alsip, IL 60803