



OROCIT

PENETRANT-SPREADER-WETTING AID FOR CITRUS CROPS

PRINCIPAL FUNCTIONING AGENTS

Alcohol Ethoxylate.....	8.15%
Constituents Ineffective as Spray Adjuvants.....	91.85%
Total.....	100.00%

All ingredients are exempt from tolerance according to 40 CFR 180

Surfactant Content: 8.15%

APENPE
FREE

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

HAZARD STATEMENTS: HARMFUL IF INHALED. CAUSES SERIOUS EYE IRRITATION.

PRECAUTIONARY STATEMENTS: Wash hands thoroughly after handling. Wear eye protection/face protection. Avoid breathing fume/mist/vapors/spray. Use only outdoors or in a well-ventilated area.

ENVIRONMENTAL HAZARDS: Do not contaminate water sources by cleaning of equipment or disposal of wastewaters.



CA Reg. No. 72662-50007-AA

MANUFACTURED BY:

Oro Agri, Inc.
2788 S. Maple Ave., Fresno, CA 93725

Copyright © Jun. 2017

Oro Agri, Inc. All Rights Reserved - v20170606

ORO AGRI

GROW WITH CONFIDENCE

Call a **POISON CENTER** or **doctor/physician** if you feel unwell.

For Chemical Emergency (spill, leak, fire or accident), call **CHEMTREC (800) 424-9300**

EYE CONTACT - IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Have the product container with you when calling a poison control center or doctor, or going for treatment. If eye irritation persists: Get medical advice or attention.

SKIN CONTACT - IF ON SKIN OR CLOTHING: Take off contaminated clothing and wash it before reuse. Wash with plenty of soap and water for several minutes. Call a poison center, doctor, chemical manufacturer or distributor to specify the appropriate source of emergency.

INGESTION - IF SWALLOWED: Do not give anything by mouth to an unconscious person. Call a poison center, doctor, chemical manufacturer or distributor to specify the appropriate source of emergency medical advice.

INHALATION - IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a poison center/doctor if you feel unwell.

Dispose of contents and containers in accordance with local, regional, and national laws.

NET CONTENTS:

2.5 US GAL (9.463 L)

GENERAL INFORMATION

OROCIT is a penetrating, wetting and spreading adjuvant to improve fungicide, miticide, insecticide, nutrient, herbicide, and PGR distribution and performance in citrus crops.

DIRECTIONS FOR USE

Prior to use, read all the directions on this label and on the labels of the manufacturers' products being used with OROCIT. The manufacturers' labels for specific usage and restrictions has precedence over directions for use referred to on the OROCIT label. Test for compatibility prior to mixing and application with other chemicals. Always test new tank-mix combinations before widespread use. If unsure, check with the manufacturers or contact your local State Extension Service if you have questions. To ensure thorough mixing, it is suggested that you recirculate mini tanks and bulk tanks or shake containers before using. Pre-rinse all tanks, spray lines or containers thoroughly before adding OROCIT as OROCIT contains surfactants. OROCIT should not be added to tanks, spray lines, or containers which may contain additional surfactant adjuvant products. Coatings on inner tank and spray lines, from previous chemicals sprayed, may be dissolved by OROCIT, thereby releasing it into the mix about to be sprayed. If in doubt, add 20 ounces of OROCIT to 20 gallons of water circulate through system for 20 minutes, spray to waste. Increasing tank mix compounds will increase precipitation and phytotoxicity risk. Extended high temperatures (>92° F) with some tank mixes may increase phytotoxicity risk and should be tested and observed before widespread use. Thereafter, mix as prescribed for regular spraying under similar conditions. Spray up to the point before runoff occurs as excessive run-off may accumulate tank mix chemicals underneath fruit. Treat a small area and observe results. If the results are satisfactory, proceed with a further application.

RECOMMENDED RATES

HERBICIDES: Use 16-100 fl.oz. OROCIT per 100 gallons of water. Use the higher rates on weeds with waxy or hairy leaf surfaces that are hard to penetrate.

INSECTICIDES, FUNGICIDES, MITICIDES, FOLIAR NUTRIENTS:

- Use 16-96 fl.oz. OROCIT per acre in sufficient water at rates no greater than 0.3% v/v.
- Do not exceed 64 fl.oz. OROCIT per acre during June to September high volume (>400 gpa) water sprays.

PLANT GROWTH REGULATORS: Use 10-20 fl.oz. OROCIT per acre in sufficient water to adequately wet target.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE: Store in original container only and keep sealed. Handle in accordance with Precautionary Statements. In the event of spillage or leakage, soak up the material with absorbent clay, sand, sawdust or other absorbent materials. Scrape up and dispose in accordance with Product Disposal.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) during mixing and loading. Recycling decontaminated containers is the best option of container disposal.

NOTICE OF WARRANTY

CONDITIONS OF SALE-LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES: To be read prior to using this product. The directions should be followed carefully together with good application practice. Failure to do so may cause injury to crops, animals, man or the environment. Extraordinary or unusual weather conditions may affect product performance and/or crop injury which are beyond the control of Oro Agri, Inc. (the "Company") or seller. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the Directions For Use. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law. The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Company's election, ONE of the following: 1. Refund of the purchase price paid by buyer or user for product bought. 2. Replacement of the product used. To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The company/seller offer this product and the buyer/user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

Patented and or patent/s pending. For more information contact legal@oroagri.com - OROCIT and ORO AGR1 are registered trademarks of Oro Agri, Inc. - v20170606

PRECAUTIONS

- **WARNING:** Do not apply to fruiting stages of sensitive or stressed crops without first consulting your Oro Agri distributor or representative.

- Test for compatibility when combining in tank mixes with EC products, copper and/or chlorpyrifos on sensitive crops.

NOTE: OROCIT reduces dew formation for up to 7 days and therefore may increase the risk of frost damage in frost prone areas.

NOTE: **NOT FOR AQUATIC USE.**

MIXING

Fill tank one-half full with water. Add the required amount of other tank mix products while agitating. Add remainder of the water. Add the recommended amount of OROCIT last and continue agitation until completion of spraying.

Antifoam and buffering agents may be used if required.