

GROUP U15 FUNGICIDE

PULL HERE TO OPEN 



Orondis® Ultra A

syngenta®

Fungicide

KEEP OUT OF REACH OF CHILDREN.

CAUTION/PRECAUCIÓN

Active Ingredient:

Oxathiapiprolin* 10.2%

Other Ingredients: 89.8%

Total: 100.0%

* CAS No. 1003318-67-9

Orondis® Ultra A is formulated as an oil dispersion and contains 0.83 pounds of oxathiapiprolin per gallon of product.

EPA Reg. No. 100-1572

EPA Est. No. 072344-MO-004

Product of France

SCP 1572A-L1A 0915

(Continued on booklet attached to back of container.)

16.4 fluid ounces Net Contents



TABLE OF CONTENTS

- 1.0 FIRST AID**
- 2.0 PRECAUTIONARY STATEMENTS**
 - 2.1 Hazards to Humans and Domestic Animals
 - 2.2 Personal Protective Equipment (PPE)
 - 2.3 Environmental Hazards

DIRECTIONS FOR USE

- 3.0 PRODUCT INFORMATION**
 - 3.1 Integrated Pest Management (IPM)
 - 3.2 Resistance Management
- 4.0 APPLICATION DIRECTIONS**
 - 4.1 Methods of Application
 - 4.2 Application Equipment
 - 4.3 Application Volume and Spray Coverage
 - 4.4 Mixing Directions
 - 4.5 Application through Irrigation Systems (Chemigation)
- 5.0 ROTATIONAL CROP RESTRICTIONS**
- 6.0 RESTRICTIONS AND PRECAUTIONS**
 - 6.1 Use Restrictions
 - 6.2 Spray Drift Precautions

continued...

TABLE OF CONTENTS *(continued)*

DIRECTIONS FOR USE

7.0 CROP USE DIRECTIONS

7.1 Brassica, Head and Stem

7.2 Bulb Vegetables

7.3 Cucurbit Vegetables

7.4 Fruiting Vegetables

7.5 Ginseng

7.6 Leafy Greens

7.7 Tobacco

7.8 Tuberous and Corm Vegetables

8.0 STORAGE AND DISPOSAL

9.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

10.0 APPENDIX

1.0 FIRST AID

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

CAUTION/PRECAUCIÓN

Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear chemical-resistant gloves made of barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or Viton® \geq 14 mils. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

2.2 Personal Protective Equipment (PPE)

Mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes and socks
- Chemical-resistant gloves made of barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or Viton \geq 14 mils.

2.2.1 USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.2.2 ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.3 Environmental Hazards

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Orondis Ultra A must be used only in accordance with instructions on this label, in separately issued labeling or exemptions under FIFRA (Supplemental Labels, Special Local Need Registration, FIFRA Section 18 exemptions), or as otherwise permitted by FIFRA. Always read the entire label, including the Conditions of Sale and Limitation of Warranty and Liability.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), and restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes and socks
- Chemical-resistant gloves (made of any waterproof material)

3.0 PRODUCT INFORMATION

Read all label directions before use. All applications must be made according to the use directions that follow.

- Orondis Ultra A is an oil dispersion containing oxathiapiprolin and is recommended for use by foliar application for the control or suppression of the diseases listed on this label.
- Orondis Ultra A is active against selective Oomycete diseases listed on this label.
- Orondis Ultra A is a systemic fungicide and moves systemically in the plant xylem. Uptake into the leaf tissue allows good translaminar movement and protection of new plant growth.
- Orondis Ultra A must be applied in a regularly scheduled protective spray program in rotation with other fungicides.
- See **Section 7.0** for specific crop/disease recommendations.

3.0.1 RAINFASTNESS

Orondis Ultra A rapidly penetrates into plant tissues and is rain-fast within 30 minutes after spray residues have dried.

3.0.2 MODE OF ACTION

Oxathiapiprolin, the active ingredient in Orondis Ultra A, acts as an oxysterol-binding protein modulator in fungal cells.

3.0.3 CROP TOLERANCE

Not all crops within a crop group, and not all varieties, cultivars, or hybrids of crops, have been individually tested for crop safety. It is not possible to evaluate for crop safety all applications of Orondis Ultra A on all crops within a crop group, on all varieties, cultivars, or hybrids of those crops, or under all environmental conditions and growing circumstances. To test for crop safety, apply the product in accordance with the label instructions to a small area of the target crop to ensure that a phytotoxic response will not occur, especially where the application is a new use of the product by the applicator.

3.1 Integrated Pest Management (IPM)

Syngenta recommends the use of Integrated Pest Management (IPM) programs to control pests. Orondis Ultra A may be used as part of an IPM program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when disease forecasting models reach locally determined action levels. Consult your state cooperative extension service, professional consultants, or other qualified authorities to determine the appropriate management, cultural practice and treatment threshold levels for the specific crop, geography and diseases.

3.2 Resistance Management

GROUP	U15	FUNGICIDE
-------	-----	-----------

Orondis Ultra A contains the active ingredient oxathiapiprolin, which has been assigned Group U15 by the Fungicide Resistance Action Committee (FRAC). Oxathiapiprolin modulates an oxysterol-binding protein (OSBP) in fungal cells. Repeated use of products for control of specific plant pathogens may lead to selection of resistant strains of fungi and result in a reduction of disease control. A disease management program for Orondis Ultra A that includes rotation and tank mixing with fungicides with a different mode of action is essential to reduce the risk of fungicide resistance development.

As part of a resistance management strategy:

- Do not tank-mix Orondis Ultra A with any fungicide for which resistance to the target disease has developed.
- Make no more than 2 sequential applications of Orondis Ultra A before rotating to a fungicide with a different mode of action.
- Do not follow soil applications of Orondis with foliar applications of Orondis Ultra A.
- Do not use Orondis Ultra A for more than 33% of the total fungicide applications per season per crop.

- For guidance on a particular crop and disease control situation, consult your state extension specialist for official state recommendations.

4.0 APPLICATION DIRECTIONS

4.1 Methods of Application

4.1.1 FOLIAR APPLICATION (INCLUDING AERIAL APPLICATION)

See **Section 7.0** for specific foliar application instructions. Orondis Ultra A may be used with adjuvants, for example, non-ionic surfactants, crop oils, methylated seed oils, and blends at typical agricultural use rates for these adjuvants.

4.2 Application Equipment

Orondis Ultra A can be applied with commonly used ground equipment, hose-end, pressurized, greenhouse and hand-held sprayers, air or chemigation equipment, except as otherwise directed, using sufficient water to obtain thorough coverage of plants. Maintain agitation during mixing and application to assure uniform product suspension.

4.2.1 SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce the effects of wind.
- However, it is the responsibility of the applicator to verify that the shields are minimizing drift potential, and not interfering with uniform deposition of the product.

4.2.2 AIR-ASSISTED (AIR-BLAST) FIELD CROP SPRAYERS

- Air-assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result.
- It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, that it is configured properly, and that drift potential has been minimized.
- **Note:** Air-assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Read the specific crop use and application equipment instructions to determine if an air-assisted field crop sprayer can be used.

4.2.3 SPRAY TANK CLEAN-OUT

- Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.
- Drain application equipment. Thoroughly rinse and flush all application equipment with clean water.
- Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

4.3 Application Volume and Spray Coverage

See **Section 7.0** for application volume information.

4.4 Mixing Directions

4.4.1 ORONDIS ULTRA A ALONE

1. Fill clean spray tank 1/2 - 2/3 full of water.
2. While agitating, add the required amount of Orondis Ultra A, continuing agitation until the product is completely dispersed.
3. Continue filling the tank, with agitation. Spray immediately after preparation, continuing agitation during spraying.

4.4.2 TANK-MIX PRECAUTIONS

- The crop safety of all tank mixtures with Orondis Ultra A which may include physically compatible pesticides, fertilizers, adjuvants, and/or additives, has not been tested.
- When using a tank mixture with Orondis Ultra A, it is important to understand crop safety.
- To test for crop safety prepare a small volume of the intended tank mixture, apply it to an area of the target crop as directed by both this label and the tank-mix partner product labels, and observe the treated crop to ensure that a phytotoxic response does not occur.
- Some materials including oils, surfactants, adjuvants, and pesticide formulations when applied individually, sequentially, or in tank mixtures may solubilize the plant cuticle, facilitate penetration into plant tissue, and increase potential for crop injury.

4.4.3 TANK-MIX COMPATIBILITY TEST

Orondis Ultra A is physically compatible with many commonly used fungicides, herbicides, insecticides, biological control products, liquid fertilizers, non-ionic surfactants, crop oils, methylated seed oils and drift control additives. However, since the formulations of products change, it is important to test the physical compatibility of desired tank mixes and check for undesirable physical effects, including settling out or flocculation. A jar compatibility test is recommended prior to tank mixing with other pesticides and/or adjuvants/additives, in order to ensure the compatibility of Orondis Ultra A with other tank-mixed pesticide, adjuvant or fertilizer partners. The recommended procedure for conducting jar tank-mix compatibility tests is as follows:

Compatibility Test: Since pesticides, adjuvants and fertilizers can vary in quality, always **check tank-mix compatibility with tank-mixed partners each time before use.** Be especially careful when using **complete** suspension or fluid fertilizers as carriers, as serious compatibility problems are more likely to occur with these products. Commercial application equipment may improve tank-mix compatibility in some instances. The following test assumes a spray volume of 25 gallons/A. For other spray volumes, make appropriate changes in the components. Check tank-mix compatibility using this procedure:

1. Add 1 pt of carrier (either the water or liquid fertilizer to be used in the spray operation) to each of two clear 1-qt jars with tight lids.
2. To **one** of the jars, add 1/4 teaspoon or 1.2 ml of a commercially available tank-mix compatibility agent approved for this use

(1/4 teaspoon is equivalent to 2 pt/100 gallons of spray). Invert the jar, shake or stir gently to ensure thorough mixing.

3. To **both** jars, add the appropriate amount of each tank-mix partner. If more than one tank-mix partner is to be used, add them separately with dry formulations (wetttable powders or water dispersible granules) first, followed by liquid flowables, capsule suspensions, emulsifiable concentrates and finally adjuvants. After each addition, invert the jar, shake or stir gently to thoroughly mix. The appropriate amount of each tank-mix partner for this test, is as follows:

Dry formulations: For each pound to be applied per acre, add 1.5 level teaspoons to each jar.

Liquid formulations: For each pint to be applied per acre, add 1/2 teaspoon or 2.5 milliliters to each jar.

4. After adding all ingredients, put lids on and tighten, then invert each jar 10 times to fully mix. Let the mixtures stand for 15-30 minutes and then assess by looking for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if a compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry dry formulations in water before addition, or (B) add the compatibility agent directly into liquid formulations, before addition to the tank-mixture. If these procedures are followed but incompatibility is still observed, do not use the tank-mixture.

4.4.4 ORONDIS ULTRA A IN TANK MIXTURES

- Always follow the tank mix instructions of the product label that is most restrictive.
- Apply at least the minimum labeled rate of each fungicide in the tank mix.
- Consult a Syngenta representative or local agricultural authorities for more information concerning tank mixtures.
- When using in a tank-mix, add different formulation types in the sequence indicated below. Allow time for complete mixing and dispersion after addition of each product.
 1. Water-soluble bag (WSB)
 2. Water-soluble granules (SG)
 3. Water-dispersible granules (WG)
 4. Wettable powders (WP)
 5. Water-based suspension concentrates (SC)
 6. Capsule suspension (CS)
 7. Suspo-emulsion (SE)
 8. Oil dispersion (OD) (Orondis Ultra A)
 9. Emulsion in water (EW)
 10. Emulsifiable concentrates (EC)
 11. Water-soluble concentrates (SL)
 12. Adjuvants, surfactants, oils
 13. Soluble fertilizers
 14. Drift retardants

4.5 Application through Irrigation Systems (Chemigation)

- Apply Orondis Ultra A only through sprinkler irrigation systems (such as center pivot, lateral move, end tow, side

(wheel) roll, traveler, big gun, solid set or hand move irrigation systems).

- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Do not connect any irrigation system (including greenhouse systems) used for pesticide applications to a public water system unless the pesticide label-prescribed safety devices for public water systems below are in place. Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals at least 60 days out of the year.
- See **Required System Safety Devices for All Chemigation and Public Water Systems (Section 4.5.1)**.
- **Preparation:** A pesticide tank is recommended for the application of Orondis Ultra A in drip chemigation systems. Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. With the mix tank 1/4 to 1/2 full with water and the agitator running, measure the required amount of Orondis Ultra A and add it to the tank. Then add additional water to bring your total pesticide mixture up to the desired volume for your application. **Note:** Always add the Orondis Ultra A to water; never put Orondis Ultra A into a dry tank or other mixing equipment without first adding water. See **Section 4.4.2** for tank-mixing sequence. Continue to agitate the mixture throughout the application process. Use mechanical or hydraulic agitation; do not use air agitation.

- **Injection into Chemigation Systems:** Inject the proper amount of Orondis Ultra A into the irrigation water flow using a positive displacement injection pump or a Venturi injector. Injection should occur at a point in the main irrigation water flow to ensure thorough mixing with the irrigation water.
- **Uniform Water Distribution:** The irrigation system used for application of Orondis Ultra A must provide for uniform distribution of Orondis Ultra A-treated water. Non-uniform distribution can result in crop injury, lack of effectiveness, or illegal pesticide residues in or on the crop being treated. Ensure the drip chemigation system is operating properly to uniformly distribute the chemigation application to the crop. Contact the equipment manufacturer, the local University Extension agent or other experts if you have questions about achieving uniform distribution of the application.
- **Monitoring of Chemigation Applications:** A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of a responsible person, shall shut the system down and make necessary adjustments should the need arise. Wear the personal protective equipment as defined in the PPE section of the label for applicators and other handlers when making adjustments or repairs on the chemigation system when Orondis Ultra A is in the irrigation water.
- **Operation:** Start the water pump and let the system achieve the desired pressure before starting the injector. Start the injector. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

- **Cleaning the System:** Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. Consult your owner's manual or your local equipment dealer for cleanout procedures for your injection system.

4.5.1 REQUIRED SYSTEM SAFETY DEVICES AND INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering device, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
8. Good agitation is required in the injection tank.
9. In moving systems, apply specified dosage of Orondis Ultra A fungicide as a continuous injection. In non-moving systems, inject Orondis Ultra A for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.
10. Mix the amount of Orondis Ultra A needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems, inject into the system

continuously for one complete revolution of the field. For non-moving systems, inject into system for the time established during calibration.

11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all Orondis Ultra A is flushed from system.

5.0 ROTATIONAL CROP RESTRICTIONS

The following crops may be planted at the specified interval following application of Orondis Ultra A.

Crop, Crop Group, or Subgroup	Plant-back Restriction (in Days) following Last Application of Orondis Ultra A
Tuberous and Corm Vegetables (Subgroup 1C)	0
Bulb Vegetables (Group 3-07)	0
Leafy Greens (Subgroup 4A)	0
Brassica, Head and Stem (Subgroup 5A)	0
Peas, Succulent Shelled	0
Peas, Edible-Podded	0
Fruiting Vegetables (Group 8-10)	0
Cucurbit Vegetables (Group 9)	0
Strawberries	0

continued...

Crop, Crop Group, or Subgroup	Plant-back Restriction (in Days) following Last Application of Orondis Ultra A
Herbs and Spices (Group 19)	0
Oilseed (Group 20)	0
Ginseng	0
Tobacco	0
Cereals (Group 15,16)	30
Grass animal feeds (Group 17)	30
Legume Vegetables, except succulent shelled and edible-podded peas	180
Non-grass Animal feed (Group 18)	180
Peanuts	180
All other crops not listed	180

6.0 RESTRICTIONS AND PRECAUTIONS

See **Section 7.0** for crop-specific restrictions and precautions.

6.1 Use Restrictions

- Use this product only in commercial and farm plantings.
- Do not use for home plantings.

- May be used in greenhouse production of tomatoes. Do not use in greenhouses on any other crops.
- Orondis Ultra A must be used only in accordance with this label.
- Do not formulate this product into other end-use products.

6.2 Spray Drift Precautions

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

6.2.1 IMPORTANCE OF DROPLET SIZE

- The most effective drift management strategy is to apply the largest droplets which are consistent with pest control objectives.
- The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage.
- Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions.
- A droplet size classification system describes the range of droplet sizes produced by spray nozzles. The American Society of Agricultural and Biological Engineers (ASABE) provides a Standard that describes droplet size spectrum

categories defined by a number of reference nozzles (fine, coarse, etc.). Droplet spectra resulting from the use of a specific nozzle may also be described in terms of volume mean diameter (VMD). Coarser droplet size spectra have larger VMD's and lower drift potential.

6.2.2 AERIAL APPLICATION SPRAY DRIFT MANAGEMENT

- **Nozzle Type** – Solid-stream or other low-drift nozzles produce the coarsest droplet spectra.
- **Number of Nozzles** – Using the minimum number of nozzles with the highest flow rate that provide uniform coverage will produce a coarser droplet spectra.
- **Nozzle Orientation** – Orienting nozzles in a manner that minimizes the effects of air shear will produce the coarsest droplet spectra. For some nozzles such as solid stream, pointing the nozzles straight back parallel to the airstream will produce a coarser droplet spectra than other orientations.
- **Pressure** – Selecting the pressure that produces the coarsest droplet spectrum for a particular nozzle and airspeed reduces spray drift potential. For some nozzle types such as solid streams, lower pressures can produce finer droplet spectra and increase drift potential.
- **Boom Length** – Using shorter booms decreases drift potential. Boom lengths are expressed as a percentage of an aircraft's wingspan or a helicopter's rotor blade diameter. Shorter boom length and proper positioning can minimize drift caused by wingtip or rotor vortices.
- **Application Height** – Applications made at the lowest height that are consistent with pest control objectives and

the safe operation of the aircraft will reduce the potential for spray drift.

6.2.3 GROUND APPLICATION SPRAY DRIFT MANAGEMENT

- **Nozzle Type** – Select a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. The use of low-drift nozzles will reduce drift potential.
- **Pressure** – The lowest spray pressures recommended for the nozzle produce the largest droplets. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, using a higher-capacity nozzle instead of increasing pressure results in the coarsest droplet spectra.
- **Flow Rate/Orifice Size** – Using the highest flow rate nozzles (largest orifice) that are consistent with pest control objectives reduces the potential for spray drift. Nozzles with higher rated flows produce coarser droplet spectra.
- **Application Height** – Applications made at the lowest height consistent with pest control objectives, and that allow the applicator to keep the boom level with the application site and minimize bounce, will reduce the exposure of spray droplets to evaporation and wind, and reduce spray drift potential.

6.2.4 WIND

- Drift potential is lowest when applications are made in light to gentle sustained winds (2-10 mph), which are blowing in a constant direction.
- Many factors, including droplet size and equipment type also determine drift potential at any given wind speed.

- AVOID GUSTY OR WINDLESS CONDITIONS.
- Local terrain can also influence wind patterns.
- Every applicator is expected to be familiar with local wind patterns and how they affect spray drift.

6.2.5 TEMPERATURE AND HUMIDITY

- Setting up equipment to produce larger droplets to compensate for droplet evaporation can reduce spray drift potential.
- Droplet evaporation is most severe when conditions are both hot and dry.

6.2.6 SURFACE TEMPERATURE INVERSIONS

- Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which may cause small suspended droplets to remain close to the ground and move laterally in a concentrated cloud.
- Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning.
- Mist or fog may indicate the presence of an inversion in humid areas. Inversions may also be identified by producing smoke and observing its behavior. Smoke that remains close to the ground, or moves laterally in a concentrated cloud under low wind conditions indicates a surface inversion. Smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

6.2.7 SENSITIVE AREAS

This pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

6.2.8 DRIFT CONTROL ADDITIVES

- Using product compatible drift control additives can reduce drift potential.
- When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label.
- If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution.
- Preferred drift control additives have been certified by the Council of Producers and Distributors of Agrotechnology.

*continued in booklet attached to
back of container...*

Orondis®, the ALLIANCE FRAME 
the SYNGENTA Logo 
and the PURPOSE ICON 
are Trademarks of a
Syngenta Group Company

Viton® is a trademark of E.I. DuPont de Nemours, Inc.

©2015 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1572A-L1A 0915

GROUP U15 FUNGICIDE

 **Orondis® Ultra A**

Fungicide

Active Ingredient:
Oxathiapiprolin*: 10.2%
Other Ingredients: 89.8%
Total: 100.0%

* CAS No. 1003318-67-9

Orondis™ Ultra A is formulated as an oil dispersion and contains 0.83 pounds of oxathiapiprolin per gallon of product.

KEEP OUT OF REACH OF CHILDREN.

**CAUTION/
PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

See additional precautionary statements and directions for use inside booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1572

EPA Est. No. 072344-MO-004

Orondis® and the Syngenta logo are trademarks of a Syngenta Group Company

SCP 1572A-L1A 0915

16.4 fluid ounces
Net Contents



GROUP U15 FUNGICIDE

PULL HERE
TO OPEN 

 **Orondis[®] Ultra A**

syngenta[®]

Fungicide

*(Continued from booklet attached
to front of container.)*

KEEP OUT OF REACH OF CHILDREN.

CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1572

SCP 1572A-L2A 0915

16.4 fluid ounces

Net Contents



7.0 CROP USE DIRECTIONS

7.1 Brassica, Head and Stem Brassica, Crop Subgroup 5A

Crops (including all cultivars, varieties, and/or hybrids of these)		
Broccoli		Cabbage, Chinese mustard
Broccoli, Chinese (gai lon)		(gai choy)
Brussels sprouts		Cauliflower
Cabbage		Cavalo broccolo
Cabbage, Chinese (Napa)		Kohlrabi
Target Disease	Rate (fl oz/A)	Application Timing
Downy mildew (<i>Peronospora parasitica</i>)	2.0 - 4.8	Begin foliar applications prior to disease development and continue on a 5- to 10-day interval.
Use Directions		
<p>Use the higher rates when disease is present, for longer application intervals, or for susceptible varieties.</p> <p>For conventional ground application, apply at least 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of the foliage.</p> <p>For air-assisted ground application, apply at least 10 gallons per acre.</p> <p>For aerial application, apply at least 2 gallons per acre.</p>		

continued...

7.1 Brassica, Head and Stem Brassica, Crop Subgroup 5A (continued)

Resistance Management: <ul style="list-style-type: none"> Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action.
USE RESTRICTIONS
1) Maximum Single Application Rate: Do not exceed 4.8 fl oz per acre per application. 2) Maximum Annual Rate: Do not exceed 19.2 fl oz per acre per year. 3) Maximum Number of Applications: <ol style="list-style-type: none"> Do not use for more than 33% of the total foliar fungicide applications or make more than four applications per crop, whichever is more restrictive. Do not exceed six foliar applications per acre per year for the same crop. 4) Minimum Application Interval: 5 days 5) Pre-harvest Interval (PHI): 0 days

7.2 Bulb Vegetables, Crop Group 3-07

Crops (including all cultivars, varieties, and/or hybrids of these)		
Chive, fresh leaves		Onion, Beltville bunching
Chive, Chinese, fresh leaves		Onion, bulb
Daylily, bulb		Onion, Chinese, bulb
Elegans hosta		Onion, fresh
Fritillaria, bulb		Onion, green
Fritillaria, leaves		Onion, macrostem
Garlic, bulb		Onion, pearl
Garlic, great-headed, bulb		Onion, potato, bulb
Garlic, serpent, bulb		Onion, tree, tops
Kurrat		Onion, Welsh, tops
Lady's leek		Shallot, bulb
Leek		Shallot, fresh leaves
Leek, wild		
Lily, bulb		
Target Disease	Rate (fl oz/A)	Application Timing
Downy mildew (<i>Peronospora destructor</i>)	2.0 - 4.8	Begin foliar applications prior to disease development and continue on a 5- to 10-day interval.

continued...

7.2 Bulb Vegetables, Crop Group 3-07 (continued)

Use Directions
Use the higher rates when disease is present, for longer application intervals, or for susceptible varieties. For conventional ground application, apply at least 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of the foliage. For air-assisted ground application, apply at least 10 gallons per acre. For aerial application, apply at least 2 gallons per acre.
Resistance Management:
<ul style="list-style-type: none"> Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action.
USE RESTRICTIONS
<ol style="list-style-type: none"> Maximum Single Application Rate: Do not exceed 4.8 fl oz per acre per application. Maximum Annual Rate: Do not exceed 19.2 fl oz per acre per year. Maximum Number of Applications: <ol style="list-style-type: none"> Do not use for more than 33% of the total foliar fungicide applications or make more than four applications per crop, whichever is more restrictive. Do not exceed six foliar applications per acre per year for the same crop. Minimum Application Interval: 5 days Pre-harvest Interval (PHI): 0 days

7.3 Cucurbit Vegetables, Crop Group 9

Crops (including all cultivars, varieties, and/or hybrids of these)																																												
<table> <tr> <td>Chayote (fruit)</td> <td>Muskmelon (continued)</td> </tr> <tr> <td>Chinese waxgourd (Chinese preserving melon)</td> <td>Honey balls</td> </tr> <tr> <td>Citron melon</td> <td>Mango melon</td> </tr> <tr> <td>Cucumber (field)</td> <td>Persian melon</td> </tr> <tr> <td>Gherkin</td> <td>Pineapple melon</td> </tr> <tr> <td>Gourd, edible</td> <td>Santa Claus melon</td> </tr> <tr> <td>Hyotan</td> <td>Snake melon</td> </tr> <tr> <td>Cucuzza</td> <td>True cantaloupe</td> </tr> <tr> <td>Hechima</td> <td>Pumpkin</td> </tr> <tr> <td>Chinese okra</td> <td>Squash, summer (field)</td> </tr> <tr> <td>Momordica spp.</td> <td>Crookneck squash</td> </tr> <tr> <td>Balsam apple</td> <td>Scallop squash</td> </tr> <tr> <td>Balsam pear</td> <td>Straightneck squash</td> </tr> <tr> <td>Bittermelon</td> <td>Vegetable marrow</td> </tr> <tr> <td>Chinese cucumber</td> <td>Zucchini</td> </tr> <tr> <td>Muskmelon</td> <td>Squash, winter</td> </tr> <tr> <td>Cantaloupe</td> <td>Acorn squash</td> </tr> <tr> <td>Casaba</td> <td>Butternut squash</td> </tr> <tr> <td>Crenshaw melon</td> <td>Calabaza</td> </tr> <tr> <td>Golden pershaw melon</td> <td>Hubbard squash</td> </tr> <tr> <td>Honeydew melon</td> <td>Spaghetti squash</td> </tr> <tr> <td></td> <td>Watermelon</td> </tr> </table>	Chayote (fruit)	Muskmelon (continued)	Chinese waxgourd (Chinese preserving melon)	Honey balls	Citron melon	Mango melon	Cucumber (field)	Persian melon	Gherkin	Pineapple melon	Gourd, edible	Santa Claus melon	Hyotan	Snake melon	Cucuzza	True cantaloupe	Hechima	Pumpkin	Chinese okra	Squash, summer (field)	Momordica spp.	Crookneck squash	Balsam apple	Scallop squash	Balsam pear	Straightneck squash	Bittermelon	Vegetable marrow	Chinese cucumber	Zucchini	Muskmelon	Squash, winter	Cantaloupe	Acorn squash	Casaba	Butternut squash	Crenshaw melon	Calabaza	Golden pershaw melon	Hubbard squash	Honeydew melon	Spaghetti squash		Watermelon
Chayote (fruit)	Muskmelon (continued)																																											
Chinese waxgourd (Chinese preserving melon)	Honey balls																																											
Citron melon	Mango melon																																											
Cucumber (field)	Persian melon																																											
Gherkin	Pineapple melon																																											
Gourd, edible	Santa Claus melon																																											
Hyotan	Snake melon																																											
Cucuzza	True cantaloupe																																											
Hechima	Pumpkin																																											
Chinese okra	Squash, summer (field)																																											
Momordica spp.	Crookneck squash																																											
Balsam apple	Scallop squash																																											
Balsam pear	Straightneck squash																																											
Bittermelon	Vegetable marrow																																											
Chinese cucumber	Zucchini																																											
Muskmelon	Squash, winter																																											
Cantaloupe	Acorn squash																																											
Casaba	Butternut squash																																											
Crenshaw melon	Calabaza																																											
Golden pershaw melon	Hubbard squash																																											
Honeydew melon	Spaghetti squash																																											
	Watermelon																																											

continued...

7.3 Cucurbit Vegetables, Crop Group 9 (continued)

Target Disease	Rate (fl oz/A)	Application Timing
Downy mildew (<i>Pseudo-peronospora cubensis</i>)	2.0 - 4.8	Begin foliar applications prior to disease development and continue on a 5- to 14-day interval.
Use Directions		
Use the higher rates when disease is present, for longer application intervals, or for susceptible varieties. For conventional ground application, apply at least 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of the foliage. For air-assisted ground application, apply at least 10 gallons per acre. For aerial application, apply at least 2 gallons per acre.		
Target Disease	Rate (fl oz/A)	Application Timing
Phytophthora Blight (<i>Phytophthora capsici</i>)	2.0 - 4.8	Begin foliar applications prior to disease development, and continue on a 3- to 14-day interval. For pickle fruit protection, apply with a copper fungicide starting at 1 inch fruit on 3- to 5-day intervals.

Use Directions
Use the higher rates when disease is present, for longer application intervals, or for susceptible varieties. For conventional ground application, apply at least 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of the foliage. For air-assisted ground application, apply at least 10 gallons per acre. For aerial application, apply at least 2 gallons per acre.
Resistance Management: <ul style="list-style-type: none"> • Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action. • Do not follow soil applications of Orondis with foliar applications of Orondis Ultra A. Use either soil applications or foliar applications but not both for disease control.
USE RESTRICTIONS
<ol style="list-style-type: none"> 1) Maximum Single Application Rate: Do not exceed 4.8 fl oz per acre per application. 2) Maximum Annual Rate: Do not exceed 19.2 fl oz per acre per year. 3) Maximum Number of Applications: <ol style="list-style-type: none"> a. Do not exceed six foliar applications per acre per year for the same crop. b. Do not use for more than 33% of the total foliar fungicide applications. 4) Minimum Application Interval: 3 days 5) Pre-harvest Interval (PHI): 0 days

7.4 Fruiting Vegetables, Crop Group 8-10

Crops (including all cultivars, varieties, and/or hybrids of these)		
African eggplant	Pea eggplant	
Bush tomato	Pepino	
Cocona	Pepper, bell (field)	
Currant tomato	Pepper, non-bell (field)	
Eggplant	Roselle	
Garden huckleberry	Scarlet eggplant	
Goji berry	Sunberry	
Groundcherry	Tomatillo	
Martynia	Tomato (field and greenhouse)	
Naranjilla	Tree tomato	
Okra		
Target Disease	Rate (fl oz/A)	Application Timing
Buckeye Rot (<i>Phytophthora parasitica</i>) Late Blight (<i>Phytophthora infestans</i>) Pepper Downy Mildew (<i>Peronospora tabacina</i>) Phytophthora Blight (<i>Phytophthora capsici</i>)	2.0 - 4.8	Begin foliar applications prior to disease development and continue on a 5- to 14-day interval.

Use Directions		
Use the higher rates when disease is present, for longer application intervals, or for susceptible varieties. For conventional ground application, apply at least 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of the foliage. For air-assisted ground application, apply at least 10 gallons per acre. For aerial application, apply at least 2 gallons per acre.		
Target Disease	Rate (fl oz/A)	Application Timing
Buckeye Rot Late Blight Phytophthora Blight (foliar)	2.0 - 4.8	Begin foliar applications prior to disease development and continue on a 5- to 14-day interval.
Use Directions		
Greenhouse Production of tomatoes: Use a rate range of 0.07 - 0.167 fl oz (0.42 tsp - 1 tsp) per gallon of spray per 1518 sq ft.		
Resistance Management:		
<ul style="list-style-type: none"> • Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action. • Do not follow soil applications of Orondis with foliar applications of Orondis Ultra A. Use either soil applications or foliar applications but not both for disease control. 		

continued...

7.4 Fruiting Vegetables, Crop Group 8-10 (continued)

USE RESTRICTIONS	
1)	Maximum Single Application Rate: Do not exceed 4.8 fl oz per acre per application.
2)	Maximum Annual Rate: Do not exceed 19.2 fl oz per acre per year.
3)	Maximum Number of Applications: <ol style="list-style-type: none"> Do not exceed six foliar applications per acre per year for the same crop. Do not use for more than 33% of the total foliar fungicide applications.
4)	Minimum Application Interval: 5 days
5)	Pre-harvest Interval (PHI): 0 days

7.5 Ginseng

Target Disease	Rate (fl oz/A)	Application Timing
Phytophthora Root Rot (<i>Phytophthora cactorum</i>)	4.8 - 38.6	Begin foliar applications prior to disease development, and continue on a 14-day interval.

Use Directions
Use the higher rates for heavy disease pressure conditions and susceptible varieties. For conventional ground application, apply at least 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of the foliage. For air-assisted ground application, apply at least 10 gallons per acre. For aerial application, apply at least 2 gallons per acre.
USE RESTRICTIONS
1) Maximum Single Application Rate: Do not exceed 38.6 fl oz per acre per application.
2) Maximum Annual Rate: Do not exceed 77.2 fl oz per acre per year.
3) Maximum Number of Applications: Do not make more than 4 applications per year.
4) Minimum Application Interval: 14 days
5) Not for use on Ginseng in California.
6) Pre-harvest Interval (PHI): 14 days

7.6 Leafy Greens (Crop Subgroup 4A)

Crops (including all cultivars, varieties, and/or hybrids of these)		
Amaranth		Endive
Arugula		Lettuce, head and leaf
Chervil		Orach
Chrysanthemum, edible-leaved		Parsley
Chrysanthemum, garland		Purslane, garden
Corn salad		Purslane, winter
Cress, garden		Radicchio
Cress, upland		Spinach
Dandelion		Spinach, New Zealand
Dock		Spinach, vine
Target Disease	Rate (fl oz/A)	Application Timing
Downy Mildew (<i>Bremia lactucae</i>)	2.0 - 4.8	Begin foliar applications prior to disease development, and continue on a 3- to 14-day interval.
Downy Mildew (<i>Peronospora farinosa</i>)	2.4 - 4.8	Begin applications prior to disease development, and continue on a 3- to 10-day interval.
Use Directions		
Use the higher rates when disease is present, for longer application intervals, or for susceptible varieties.		

Use Directions (continued)
For conventional ground application, apply at least 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of the foliage.
For air-assisted ground application, apply at least 10 gallons per acre.
For aerial application, apply at least 2 gallons per acre.
Resistance Management:
<ul style="list-style-type: none"> • Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action. • Do not follow soil applications of Orondis with foliar applications of Orondis Ultra A. Use either soil applications or foliar applications but not both for disease control.
USE RESTRICTIONS
1) Maximum Single Application Rate: Do not exceed 4.8 fl oz per acre per application.
2) Maximum Annual Rate: Do not exceed 19.2 fl oz per acre per year.
3) Maximum Number of Applications: <ol style="list-style-type: none"> Do not exceed six foliar applications per acre per year for the same crop. Do not use for more than 33% of the total foliar fungicide applications.
4) Minimum Application Interval: 3 days
5) Pre-harvest Interval (PHI): 0 days

7.7 Tobacco

Target Disease	Rate (fl oz/A)	Application Timing
Blue Mold (<i>Peronospora tabacina</i>)	2.0 - 4.8	Begin applications prior to disease development, and continue on a 7- to 10-day interval.
Use Directions		
<p>Use the higher rates when disease is present, for longer application intervals, or for susceptible varieties.</p> <p>For conventional ground application, apply at least 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of the foliage.</p> <p>For air-assisted ground application, apply at least 10 gallons per acre.</p> <p>For aerial application, apply at least 2 gallons per acre.</p>		
Resistance Management:		
<ul style="list-style-type: none"> • Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action. • Do not follow soil applications of Orondis with foliar applications of Orondis Ultra A. Use either soil applications or foliar applications but not both for disease control. 		

USE RESTRICTIONS

- 1) **Maximum Single Application Rate:** Do not exceed 4.8 fl oz per acre per application.
- 2) **Maximum Annual Rate:** Do not exceed 19.2 fl oz per acre per year.
- 3) **Maximum Number of Applications:**
 - a. Do not exceed six foliar applications per acre per year for the same crop.
 - b. Do not use for more than 33% of the total foliar fungicide applications.
- 4) **Minimum Application Interval:** 7 days
- 5) Not for use on Tobacco in California
- 6) **Pre-harvest Interval (PHI):** 7 days

7.8 Tuberos and Corm Vegetables, Crop Subgroup 1C

Crops (including all cultivars, varieties, and/or hybrids of these)

Arracacha	Dasheen (taro)
Arrowroot	Ginger
Artichoke, Chinese	Potato
Artichoke, Jerusalem	Sweet potato
Canna, edible	Tanier
Cassava, bitter and sweet	Turmeric
Chayote (root)	Yam bean
Chufa	Yam, true

continued...

7.8 Tuberos and Corm Vegetables, Crop Subgroup 1C (continued)

Target Disease	Rate (fl oz/A)	Application Timing
Late Blight (<i>Phytophthora infestans</i>)	1.6 - 4.8	Begin applications prior to disease development, and continue on a 5- to 14-day interval.
Use Directions		
<p>Use the higher rates when disease is present, for longer application intervals, or for susceptible varieties.</p> <p>For conventional ground application, apply at least 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of the foliage.</p> <p>For air-assisted ground application, apply at least 10 gallons per acre.</p> <p>For aerial application, apply at least 2 gallons per acre.</p>		
Resistance Management:		
<ul style="list-style-type: none"> • Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action. 		

USE RESTRICTIONS
1) Maximum Single Application Rate: Do not exceed 4.8 fl oz per acre per application.
2) Maximum Annual Rate: Do not exceed 27.2 fl oz per acre per year.
3) Maximum Number of Applications: <ol style="list-style-type: none"> Do not use for more than 33% of the total foliar fungicide applications or make more than four applications per crop, whichever is more restrictive. Do not exceed six foliar applications per acre per year for the same crop.
4) Minimum Application Interval: 5 days
5) Not for use on Tuberos and Corm Vegetables in California
6) Pre-harvest Interval (PHI): 5 days

8.0 STORAGE AND DISPOSAL

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Keep container closed when not in use. Always store pesticides in the original container only, away from other pesticides, food, pet food, feed, seed, fertilizers, and veterinary supplies. If a leaky container must be contained within another, mark the outer container to identify the contents. Storage areas must be locked and secure from vandalism, with precautionary signs posted. The storage area must be dry, well-lit, and well-ventilated. Keep pesticide storage areas clean. Clean up any spills promptly. Protect pesticide containers from extreme heat and cold. Store herbicides, insecticides and fungicides in separate areas within the storage unit. Place liquid formulations on lower shelves and dry formulations above. Maintaining a spill kit and fire extinguisher on hand and having emergency phone numbers posted will allow you to be prepared for emergencies. If spill cleanup PPE is stored nearby, but outside the pesticide storage area, it will be accessible when needed.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons – mini-bulk]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

continued...

STORAGE AND DISPOSAL (continued)

Container Handling [greater than 5 gallons – bulk]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean the container before final disposal, empty the remaining contents from this container into application or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

9.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

10.0 APPENDIX

10.1 Orondis Ultra A Use Summary Table

IMPORTANT: The table below is a summary of the Crop Use Directions for Orondis Ultra A. However, it is important for the user to read and follow the complete instructions contained within this label.

Crop or Crop Group or subgroup with examples	Maximum Rate per Application (fl oz/A)	Maximum Rate per Application (lb ai/A)	Minimum Application Interval (days)
Brassica, Head and Stem: cabbage, broccoli, cauliflower	4.8	0.03	5
Bulb Vegetables (Crop Group 3-07)	4.8	0.03	5
Cucurbit Vegetables (Crop Group 9): cucumber, cantaloupe, watermelon, squash	4.8	0.03	3
Fruiting Vegetables (Crop Group 8-10): tomato, pepper	4.8	0.03	5
Ginseng*	38.6	0.25	14
Leafy Greens (Crop Subgroup 4A): lettuce, spinach	4.8	0.03	3
Tobacco*	4.8	0.03	7
Tuberous and Corm vegetables* (Crop Subgroup 1C): potato	4.8	0.04	5

continued...

Crop or Crop Group or subgroup with examples	Pre-Harvest Interval (PHI days)	Maximum Rate per Year (fl oz/A)	Maximum Rate per Year (lb ai/A)
Brassica, Head and Stem: cabbage, broccoli, cauliflower	0	19.2	0.12
Bulb Vegetables (Crop Group 3-07)	0	19.2	0.12
Cucurbit Vegetables (Crop Group 9): cucumber, cantaloupe, watermelon, squash	0	19.2	0.12
Fruiting Vegetables (Crop Group 8-10): tomato, pepper	0	19.2	0.12
Ginseng*	14	77.2	0.50
Leafy Greens (Crop Subgroup 4A): lettuce, spinach	0	19.2	0.12
Tobacco*	7	19.2	0.12
Tuberous and Corm vegetables* (Crop Subgroup 1C): potato	5	27.2	0.18

* Not for use on these crops in California.

Orondis®, the ALLIANCE FRAME
the SYNGENTA Logo
and the PURPOSE ICON
are Trademarks of a
Syngenta Group Company



Viton® is a trademark of E.I. DuPont de Nemours, Inc.

©2015 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1572A-L2A 0915

FIRST AID If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **HOT LINE NUMBER:** For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call **1-800-888-8372.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION/PRECAUCIÓN

Environmental Hazards: For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

STORAGE AND DISPOSAL

Refer to attached booklet for Pesticide Storage and Pesticide Disposal instructions.

Container Handling: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

©2015 Syngenta
Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419-8300

SCP 1572A-L2A 0915